



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SUEBBY HILLS

2007

**TRIAL HIGHER SCHOOL
CERTIFICATE EXAMINATION**

Mathematics Extension 1

General Instructions

- Reading Time – 5 Minutes
- Working time – 2 Hours
- Write using black or blue pen. Pencil may be used for diagrams.
- Board-approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown in every question.
- Each Question is to be returned in a separate bundle.

Total Marks – 84

- Attempt Questions 1 – 7.
- All questions are of equal value.

Examiner: *A. Fuller*

Total marks – 84
Attempt Questions 1-7
All questions are of equal value

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

Marks

Question 1 (12 marks) Use a SEPARATE writing booklet.

- (a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin 4x}{5x}$. 1
- (b) Calculate the acute angle (to the nearest minute) between the lines $2x + y = 4$ and $x - 3y = 6$. 2
- (c) (i) Show that $x + 1$ is a factor of $x^3 - 4x^2 + x + 6$. 1
- (ii) Hence, or otherwise factorise $x^3 - 4x^2 + x + 6$ fully. 2
- (d) The point $P(5, 7)$ divides the interval joining the points $A(-1, 1)$ and $B(3, 5)$ externally in the ratio $k:1$. Find the value of k . 2
- (e) Find the horizontal asymptote of the function $y = \frac{3x^2 - 4x + 1}{2x^2 - 1}$. 1
- (f) Find a primitive of $\frac{1}{\sqrt{4 - x^2}}$. 1
- (g) Solve the equation $|x + 1|^2 - 4|x + 1| - 5 = 0$. 2

Question 2 (12 marks)

- (a) Let $f(x) = \frac{1}{2} \cos^{-1}\left(\frac{x}{3}\right)$.
- (i) State the domain and range of the function $f(x)$. 2
- (ii) Show that $y = f(x)$ is a decreasing function. 2
- (iii) Find the equation of the tangent to the curve $y = f(x)$ at the point where $x = 0$. 2
- (b) Find the derivative of $y = \ln(\sin^3 x)$. 2
- (c) (i) Write $\cos x - \sqrt{3} \sin x$ in the form $A \cos(x + \alpha)$, where $A > 0$ and $0 < \alpha < \frac{\pi}{2}$. 2
- (ii) Hence, or otherwise, solve $\cos x - \sqrt{3} \sin x + 1 = 0$ for $0 \leq x \leq 2\pi$. 2

Marks

Question 3 (12 marks) Use a SEPARATE writing booklet.

- (a) (i) Show that the equation $e^x - x - 2 = 0$ has a solution in the interval $1 < x < 2$. 1
- (ii) Taking an initial approximation of $x = 1.5$ use one application of Newton's method to approximate the solution, correct to three decimal places. 2
- (b) The normal at $P(2ap, ap^2)$ on the parabola $x^2 = 4ay$ cuts the y -axis at Q and is produced to a point R such that $PQ = QR$.
- (i) Show that the equation of the normal at P is $x + py = 2ap + ap^3$. 2
- (ii) Find the coordinates of Q . 1
- (iii) Show that R has coordinates $(-2ap, ap^2 + 4a)$. 1
- (iv) Show that the locus of R is a parabola, and find its vertex. 3
- (c) If $\int_1^5 f(x) dx = 3$, find $\int_1^5 (2f(x) + 1) dx$. 2

Marks


Marks

Question 4 (12 marks) Use a SEPARATE writing booklet.

(a) Using the substitution $u = e^x$, or otherwise, find $\int e^{(e^x+x)} dx$

3

(b) The velocity-time graph below shows the velocity of a lift as it travels from the first floor to the twentieth floor of a tall building during the T seconds of its motion.



The velocity v m/s at time t s for $0 \leq t \leq 2$ is given by $v = t^2(3-t)$. After the first two seconds, the lift moves with a constant velocity of 4 m/s for a time, and then decelerates to rest in the final two seconds.

The velocity-time graph is symmetrical about $t = \frac{1}{2}T$.

(i) Express the acceleration in terms of t for the first two seconds of the motion of the lift.

1

(ii) Hence, find the maximum acceleration of the lift during the first two seconds of its motion.

2

(iii) Given that the total distance travelled by the lift during its journey is 41 metres, find the exact value of T .

2

(c) A solid is formed by rotating about the y -axis the region bounded by the curve $y = \cos^{-1} x$, the x -axis and the y -axis.

(i) Show that the volume of the solid is given by $V = \pi \int_0^{\frac{\pi}{2}} \cos^2 y dy$.

1

(ii) Calculate the volume of this solid.

3


Marks

Question 5 (12 marks) Use a SEPARATE writing booklet.

- (a) Use mathematical induction to prove that $\sum_{r=1}^n r \times r! = (n+1)! - 1$. 3
- (b) In the expansion of $\left(2x + \frac{1}{x^2}\right)^{15}$, determine the coefficient of the term that is independent of x . 3
- (c) The acceleration of a particle P is given by the equation $a = 8x(x^2 + 1)$, where x is the displacement of P from the origin in metres after t seconds, with movement being in a straight line. Initially, the particle is projected from the origin with a velocity of 2 metres per second in the negative direction.
- (i) Show that the velocity of the particle can be expressed as $v = 2(x^3 + 1)$. 2
- (ii) Hence, show that the equation describing the displacement of the particle at time t is given by $x = \tan 2t$. 2
- (iii) Determine the velocity of the particle after $\frac{\pi}{8}$ seconds. 2

Marks

Question 6 (12 marks) Use a SEPARATE writing booklet.



(a)

A, B, C and D are points on the circumference of a circle with centre O . EF is a tangent to the circle at C and the angle ECD is 60° .

3


Find the value of $\angle BAC$ giving reasons.

- (b) (i) By considering the expansion of $(1+x)^n$ in ascending powers of x , where n is a positive integer, and differentiating, show that $\binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \dots + n\binom{n}{n} = n(2^{n-1})$. 1
- (ii) Hence, find an expression for $2\binom{n}{1} + 3\binom{n}{2} + 4\binom{n}{3} + \dots + (n+1)\binom{n}{n}$. 2
- (c) If $f(x+2) = x^2 + 2$, find $f(x)$. 2
- (d) At a particular dinner, each rectangular table has nine seats, five facing the stage and four with their backs to the stage. In how many ways can 9 people be seated at the table if
- (i) John and Mary sit on the same side? 2
- (ii) John and Mary sit on opposite sides? 2

Marks


Question 7 (12 marks) Use a SEPARATE writing booklet.

- (a) A skier accelerates down a slope and then skis up a short jump (see diagram). The skier leaves the jump at a speed of 12 m/s and at an angle of 60° to the horizontal. The skier performs various gymnastic twists and lands on a straight line section of the 45° down-slope T seconds after leaving the jump. Let the origin O of a Cartesian coordinate system be at the point where the skier leaves the jump. Displacements are measured in metres and time in seconds. Let $g = 10 \text{ m s}^{-2}$ and neglect air resistance.



- (i) Derive the cartesian equation of the skier's flight as a function of y in terms of x . 3
- (ii) Show that $T = \frac{6}{5}(\sqrt{3} + 1)$. 3
- (iii) At what speed, in metres per second does the skier land on the down-slope? Give your answer correct to one decimal place. 2

(b)



ABC is a horizontal, right-angled, isosceles triangle where $AB = BC$ and $\angle ABC = 90^\circ$. P is vertically above B ; Q is vertically above C .

The angle of elevation of P from A , and Q from P are α and β respectively.

- (i) If the angle of elevation of Q from A is θ , prove that 2

$$\tan \theta = \frac{\tan \alpha + \tan \beta}{\sqrt{2}}$$
- (ii) If $\angle APQ = \phi$, prove that $\cos \phi = -\sin \alpha \sin \beta$. 2

End of Paper

2007 THSC Mathematics Extension 1: Solutions— Question 1

1. (a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin 4x}{5x}$.

1

Solution: $\lim_{x \rightarrow 0} \frac{\sin 4x}{5x} = \lim_{x \rightarrow 0} \frac{\sin 4x}{4x} \times \frac{4}{5},$
 $= \frac{4}{5} \times \lim_{x \rightarrow 0} \frac{\sin 4x}{4x},$
 $= \frac{4}{5}.$

(b) Calculate the acute angle (to the nearest minute) between the lines $2x + y = 4$ and $x - 3y = 6$.

2

Solution: $\tan \alpha = \frac{|-2 - 1/3|}{1 + (-2) \times (1/3)},$
 $= 7.$
 $\therefore \alpha = \tan^{-1} 7,$
 $= 81.86989765^\circ$ by calculator,
 $= 81^\circ 52'.$

(c) i. Show that $x + 1$ is a factor of $x^3 - 4x^2 + x + 6$.

1

Solution: Putting $P(x) = x^3 - 4x^2 + x + 6;$
 $P(-1) = -1 - 4 - 1 + 6,$
 $= 0.$
 $\therefore x + 1$ is a factor.

ii. Hence or otherwise factorise $x^3 - 4x^2 + x + 6$ fully.

2

Solution: Possible factors of 6 are 1, 2, 3 or 1, -2, -3.
 $P(-2) = -8 - 16 - 2 + 6 \neq 0,$
 $P(2) = 8 - 16 + 2 + 6,$
 $= 0.$
 $\therefore x^3 - 4x^2 + x + 6 = (x + 1)(x - 2)(x - 3).$

(d) The point $P(5, 7)$ divides the interval joining the points $A(-1, 1)$ and $B(3, 5)$ externally in the ratio $k : 1$. Find the value of k .

2

Solution: $\frac{5 - -1}{5 - 3} = \frac{k}{1},$
 $6 = 2k,$
 $k = 3.$

(e) Find the horizontal asymptote of the function $y = \frac{3x^2 - 4x + 1}{2x^2 - 1}$.

1

Solution: $\lim_{x \rightarrow \pm\infty} \frac{3 - 4/x + 1/x^2}{2 - 1/x^2} = \frac{3}{2}.$
 $\therefore y = 3/2$ is the horizontal asymptote.

(f) Find a primitive of $\frac{1}{\sqrt{4 - x^2}}$.

1

Solution: From the table of standard integrals,

$$\int \frac{dx}{\sqrt{4 - x^2}} = \sin^{-1} \frac{x}{2} + c.$$

(g) Solve the equation $|x + 1|^2 - 4|x + 1| - 5 = 0$.

2

Solution: Putting $y = |x + 1|;$
 $y^2 - 4y - 5 = 0,$
 $(y - 5)(y + 1) = 0,$
 $\therefore y = 5$ or $-1.$
 But $|x + 1| \geq 0,$
 hence $x + 1 = 5$ or $x + 1 = -5,$
 so $x = 4, -6.$

Solutions Q.20 3 unit anal MSC 2001

(a) $f(x) = \frac{1}{2} \cos^{-1}(\frac{x}{3})$

$y = \cos^{-1} x$ has $-1 \leq x \leq 1$
 $0 \leq y \leq \pi$

Domain $-1 \leq x \leq 1$
 $-1 \leq \frac{x}{3} \leq 1$
 $-3 \leq x \leq 3$ ①

(ii) $f'(x) = \frac{1}{2} \times \frac{-1}{\sqrt{1-\frac{x^2}{9}}} \times \frac{1}{3}$

$= -\frac{1}{6} \times \frac{1}{\sqrt{1-\frac{x^2}{9}}} = -\frac{1}{6} \times \frac{3}{\sqrt{9-x^2}} = -\frac{1}{2} \cdot \frac{1}{\sqrt{9-x^2}}$

So for $-3 < x < 3$, $f'(x) < 0$ always. ②

(iii) when $x=0$, $f(x) = \frac{1}{2} \cos^{-1}(\frac{0}{3})$
 $= \frac{1}{2} \cos^{-1}(0)$
 $= \frac{1}{2} \times \frac{\pi}{2} = \frac{\pi}{4}$

$f(x) = \frac{1}{2\sqrt{9-x^2}}$

at $x=0$, $m = \frac{-1}{2 \times 3} = -\frac{1}{6}$

$(y-y_1) = m(x-x_1)$
 $(y-\frac{\pi}{4}) = -\frac{1}{6}(x-0)$

$y = -\frac{1}{6}x + \frac{\pi}{4}$
 $\frac{1}{6}x + y - \frac{\pi}{4} = 0$

or $12 \times \frac{1}{6}x + 12y - 12 \times \frac{\pi}{4} = 0$
 $2x + 12y - 3\pi = 0$ ③

12.

Range: $0 \leq \cos^{-1} x \leq \pi$

$\frac{1}{2} \times 0 \leq \frac{1}{2} \cos^{-1}(\frac{x}{3}) \leq \frac{1}{2} \times \pi$

$0 \leq f(x) \leq \frac{\pi}{2}$ ①

(c) (i) $\cos x - \sqrt{3} \sin x = 2$

$= 2(\frac{1}{2} \cos x - \frac{\sqrt{3}}{2} \sin x)$

$= A(\cos x \cos \alpha - \sin x \sin \alpha)$

$\Rightarrow A = 2$ ①

$\cos \alpha = \frac{1}{2}$
 $\sin \alpha = \frac{\sqrt{3}}{2}$ } $\alpha = 60^\circ = \frac{\pi}{3}$ ① (1st quad $0 < \alpha < \frac{\pi}{2}$)

So $\cos x - \sqrt{3} \sin x = 2 \cos(x + \frac{\pi}{3})$

(ii) Now $\cos x - \sqrt{3} \sin x + 1 = 0$

$\cos x - \sqrt{3} \sin x = -1$

$2 \cos(x + \frac{\pi}{3}) = -1$

$\cos(x + \frac{\pi}{3}) = -\frac{1}{2}$

$\cos \frac{\pi}{3} = \frac{1}{2}$

2nd / 3rd quad.

now $x + \frac{\pi}{3} = \pi - \frac{\pi}{3}$ 2nd quad.

and $x + \frac{\pi}{3} = \pi + \frac{\pi}{3}$ 3rd quad.

So $x = \pi - \frac{2\pi}{3} = \frac{3\pi - 2\pi}{3} = \frac{\pi}{3}$ ✓ ②

and $x = \pi + \frac{\pi}{3} - \frac{\pi}{3} = \pi$ ✓

Q3 (a) let $f(x) = e^x - x - 2$.

Now $f(1) = e - 1 - 2 = e - 3 < 0$. (≈ -0.28)

\checkmark $f(2) = e^2 - 2 - 2 = e^2 - 4 > 0$. (≈ 3.38)

Since $f(x)$ changes sign in $1 < x < 2$ (\checkmark)

$f(x) = 0$ has a solution in $1 < x < 2$.


(i) Now $x_1 = 1.5$, $\frac{f(x_1)}{f'(x_1)}$ \checkmark $f(x_1) = e^{1.5} - 1.5 - 2$
 $f'(x_1) = e^{1.5} - 1$

$\therefore x_2 = 1.5 - \frac{f(1.5)}{f'(1.5)}$
 $= 1.5 - \frac{(e^{1.5} - 1.5 - 2)}{e^{1.5} - 1}$

$\approx 1.5 - \frac{0.98168}{3.481689}$

$\approx 1.5 - 0.281915$

≈ 1.218 (\checkmark)



(i) $y = \frac{1}{4a} x^2$

$y' = \frac{1}{2a} x$

$\therefore m_T = \frac{2ap}{2a} = p$

$\therefore m_N = -\frac{1}{p}$

H: $\frac{y - ap^2}{x - 2ap} = -\frac{1}{p}$
 $p(y - ap^2) = -x + 2ap$
 $\left[\frac{x + py = 2ap + ap^3}{x + py = 2ap + ap^3} \right]$ (\checkmark)

(ii) Co-ord of Q. $x = 0$ $\therefore py = 2ap + ap^3$
 $y = 2a + ap^2$ (\checkmark)
 $\therefore Q(0, 2a + ap^2)$

(iii) Q is the mid-pt of PR.

Let R be (x_1, y_1)

$\therefore \frac{x_1 + 2ap}{2} = 0$

$\therefore x_1 = -2ap$

$\frac{y_1 + ap^2}{2} = 2a + ap^2$

$y_1 + ap^2 = 4a + 2ap^2$

$y_1 = 4a + ap^2$

$\therefore R(-2ap, 4a + ap^2)$ (\checkmark)

(iii) To find the Area of R. $x = -2ap \therefore p = \frac{x}{-2a}$

$\therefore y = a \left(\frac{x}{-2a} \right)^2 + 4a$

$y = \frac{x^2}{4a} + 4a$

$$4ay = z^2 + 16az$$

$$z^2 = 4ay - 16az$$

$$z^2 = 4z(z - 4a)$$

Part 180-4
 15/25/2018
 (0, 4a)

(c)

$$\int_1^5 (2ax + 1) dx = 2 \int_1^5 fax dx + \int_1^5 1 dx$$

$$= 2 \times 3 + [x]^5_1$$

$$= 6 + (5-1)$$

$$= 10$$

(c)

The max. acceleration occurs when $f = 1$
 $a_{max} = 3 \times 1 = 3 \text{ m/s}^2$

(ii) Let $d(t)$ be the total distance travelled
 (where is $t \text{ m}$)
 $\therefore d(t) = 2 \int_0^t v(t) dt$
 $= 2 \int_0^t (3t^2 - t^3) dt$
 $= [4t^3 - \frac{1}{4}t^4]_0^t$
 $\therefore d = 4t^3 - \frac{1}{4}t^4$
 $\therefore d = 2(8-4) + 4T - 16$
 $41 = 4T - 8 \Rightarrow T = 12 \frac{1}{4}$

(iii) For $0 \leq t \leq 2$
 (i) $v = 3t^2 - t^3$
 $a = \frac{dv}{dt} = 6t - 3t^2 = 3t(2-t)$

Question (f)

(a) Let $u = e^x$
 $du = e^x dx$
 $\int e^{(x+k)} dx = \int e^x \cdot e^k dx = e^k \int e^x dx = e^k e^x + c = e^{x+k} + c$

(b) For $0 \leq t \leq 2$
 (i) $v = 3t^2 - t^3$
 $a = \frac{dv}{dt} = 6t - 3t^2 = 3t(2-t)$

From the graph of a versus t .

EX1 QUESTIONS

(a) If $n=1$, $1 \times 1! = (1+1)! - 1$

$\therefore P(1)$ is true

Assume $P(k)$ is true $1 \times 1! + 2 \times 2! + \dots + k \times k! = (k+1)! - 1$

If $P(k+1)$ is $1 \times 1! + 2 \times 2! + \dots + k \times k! + (k+1) \times (k+1)! = (k+2)! - 1$

LHS is $(k+1)! - 1 + (k+1)(k+1)!$ using assumption

$= (k+1)! (1 + k+1) - 1$

$= (k+1)! (k+2) - 1 = (k+2)! - 1 = \text{RHS}$

$\therefore P(k+1)$ is true if $P(k)$ is true. $P(1)$ is true, by Mathematical

induction $\sum_{r=1}^n r \times r! = (n+1)! - 1$

(b) $T_{k+1} = {}^{15}C_k (2x)^{n-k} (x^{-2})^k$

for term independent of x , $n-k-2k=0$, $k=5$

$\therefore {}^{15}C_5 \times 2^{10} = 3075072$

(c) (i) $d\left(\frac{1}{2}v^2\right) = 8x(x^2+1) = 8x^3+8x$

$\frac{1}{2}v^2 = 2x^4 + 4x^2 + C$

$v = -2$, $x = 0$, $C = 2$

$v^2 = 4x^4 + 8x^2 + 4 = 4(x^4 + 2x^2 + 1)$

$v = \pm 2(x^2+1)^{-1/2}$

(ii) if $\frac{dx}{dt} = 2(x^2+1)$

$\frac{dt}{dx} = \frac{1}{2(x^2+1)}$

$t = \frac{1}{2} \tan^{-1} x + C$

$t = 0$, $x = 0$, $C = 0$

$2t = \tan^{-1} x$

$x = \tan 2t$

(iii)

$t = \frac{\pi}{8}$, $x = \tan \frac{\pi}{4}$

$= 1$

$v = 2(1+1)$ from (i)

$v = 4 \text{ m/s}$

Sydney Boys' High School
Trial HSC 2007 - Mathematics Extension 1

Question 6

(a) $\angle CBD = 60^\circ$ (alternate segment theorem)

$\angle BCD = 90^\circ$ (angle in semicircle)

$\therefore \angle CDB = 30^\circ$ (angle sum of triangle)

$\therefore \angle CAB = 30^\circ$ (angles at circumference on same arc)

(b) (i) $(1+x)^n = 1 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_{n-1} x^{n-1} + x^n$

Differentiating with respect to x :

$n(1+x)^{n-1} = {}^nC_1 + 2{}^nC_2 x + 3{}^nC_3 x^2 + \dots + n{}^nC_n x^{n-1}$

Let $x = 1$:

$n2^{n-1} = {}^nC_1 + 2{}^nC_2 + 3{}^nC_3 + \dots + n{}^nC_n$

QED

(ii) Multiplying $(1+x)^n$ by x :

$x(1+x)^n = {}^nC_0 x + {}^nC_1 x^2 + {}^nC_2 x^3 + \dots + {}^nC_n x^{n+1}$

Differentiating with respect to x :

$n(1+x)^{n-1} + (1+x)^n = {}^nC_0 + 2{}^nC_1 x + 3{}^nC_2 x^2 + \dots + (n+1){}^nC_n x^n$

Let $x = 1$:

$n(2)^{n-1} + (2)^n = 1 + 2{}^nC_1 + 3{}^nC_2 + \dots + (n+1){}^nC_n$

Thus

$2^n C_1 + 3^n C_2 + \dots + (n+1)^n C_n = n(2)^{n-1} + (2)^n - 1$
 $= (n+2)2^{n-1} - 1$


(c) $f(x+2) = x^2 + 2$

$f(x) = (x-2)^2 + 2$

$= x^2 - 4x + 4 + 2$

$= x^2 - 4x + 6$

(d)



- (i) If J&M sit on the short side, they can be arranged in 12 ways, and the other guests in 7! ways. Thus $12 \times 7!$ ways.
 If J&M sit on the long side they can be arranged in 20 ways, and the other guests in 7! Ways. Thus $20 \times 7!$

Hence there are $32 \times 7! = 161\,280$ ways.


- (ii) If John sits on the short side he has four seats available, and Mary (on the long side) has 5, thus $20 \times 7!$

But Mary may be the one on the short side.

Thus the total is $40 \times 7! = 201\,600$

2007 EXT 1

Q7



$$(i) \quad \tan \alpha = \frac{BP}{x} = \frac{BP}{AB}$$

$$\therefore BP = x \tan \alpha$$

$$\tan \beta = \frac{BQ}{x} = \frac{BQ}{AB}$$

$$\therefore BQ = x \tan \beta$$

$$QC = x \tan \alpha + x \tan \beta$$

$$= x (\tan \alpha + \tan \beta)$$

$$\tan \theta = \frac{QC}{AP}$$

$$= \frac{x (\tan \alpha + \tan \beta)}{x}$$

$$= \frac{\sqrt{2} x}{\sqrt{2}} \tan \beta$$

GIVEN

2

$$(ii) \quad \cos \theta = \frac{AP^2 + QC^2 - AC^2}{2 \cdot AP \cdot QC}$$

$$= \frac{x^2 + x^2 (\tan \alpha + \tan \beta)^2 - 2x^2}{2 \cdot x \cdot x (\tan \alpha + \tan \beta)}$$

$$= \frac{\sec^2 \alpha + \sec^2 \beta - 2 (1 + \tan \alpha \tan \beta)}{2 \sec \alpha \sec \beta}$$

$$= \frac{\sec^2 \alpha + \sec^2 \beta - (2 + \tan^2 \alpha + 2 \tan \alpha \tan \beta + \tan^2 \beta)}{2 \sec \alpha \sec \beta}$$

$$= \frac{\sec^2 \alpha + \sec^2 \beta - \sec^2 \alpha - 2 \tan \alpha \tan \beta - \sec^2 \beta}{2 \sec \alpha \sec \beta}$$

$$= -\sin \alpha \sin \beta$$

GIVEN

2

$$\ddot{y} = -10$$

$$\dot{y} = 10t + 12.5 \sin 60^\circ$$

$$= 6$$

$$y = -5t^2 + 6\sqrt{3}t$$

$$y = -5 \left(\frac{2}{5}\right)^2 + 6\sqrt{3} \times \frac{2}{5}$$

$$= -\frac{5 \times 2^2}{5 \times 5} + \sqrt{3} \times 2$$

3

If $x = -y$, $-x = -\frac{5x^2}{5} + \sqrt{3}x$

$$\therefore \frac{5x^2}{5} - (\sqrt{3}+1)x = 0$$

$$\therefore x(5x - 36(\sqrt{3}+1)) = 0$$

$$\therefore x = 0 \text{ or } x = \frac{36(\sqrt{3}+1)}{5}$$

DIVIDE

$$\therefore \text{IF} = \frac{6(\sqrt{3}+1)}{5}$$

3

$$x = 6$$

$$\dot{y} = -10 \times \frac{6(\sqrt{3}+1)}{5} + 6\sqrt{3}$$

$$= -12(\sqrt{3}+1) + 6\sqrt{3} = -6\sqrt{3} - 12$$

$$\text{Speed} = \sqrt{36 + (-6\sqrt{3} - 12)^2}$$

$$= \sqrt{36 + 108 + 144 + 144\sqrt{3}}$$

$$= \sqrt{288 + 144\sqrt{3}}$$

$$= \sqrt{144(2 + \sqrt{3})} = 12\sqrt{2 + \sqrt{3}}$$

$$= 23.2$$

2