



SYDNEY BOYS HIGH
MOORE PARK, SURRY HILLS
AUGUST 2007
TRIAL HSC
YEAR 12

Mathematics Extension 2

General Instructions:

- Reading time—5 minutes.
- Working time—3 hours.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may NOT be awarded for messy or badly arranged work.


Total marks—120 Marks

- Attempt questions 1–8.
- All questions are of equal value.
- Start each question in a separate answer booklet.

Examiner: Mr P. Bigelow

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

Question 1 (15 marks)



Copy the diagram onto your answer booklet.

Carefully indicate the position of the following:

- | | |
|--|--------------------------|
| (i) R representing \bar{z} , | <input type="checkbox"/> |
| (ii) Q representing $-\frac{1}{2}z$, | <input type="checkbox"/> |
| (iii) S representing $\frac{1}{z}$, | <input type="checkbox"/> |
| (iv) T representing \sqrt{z} . | <input type="checkbox"/> |
|
 | |
| (b) The complex number z is given by $z = -1 + i\sqrt{3}$. | <input type="checkbox"/> |
| (i) Show that $z^2 = 2\bar{z}$. | <input type="checkbox"/> |
| (ii) Evaluate $ z $ and $\arg z$. | <input type="checkbox"/> |
| (iii) Show that z is a root of the equation $z^3 - 8 = 0$. | <input type="checkbox"/> |
|
 | |
| (c) On an Argand diagram, shade the region where the inequalities $0 \leq \operatorname{Re}(z) \leq 4$ and $ z - 1 + i \leq 4$ both hold. | <input type="checkbox"/> |
|
 | |
| (d) If $z^2 = i$, find z in the form $a + ib$ where a and b are real. | <input type="checkbox"/> |
|
 | |
| (e) Give reasons why each of the following statements is true or false.
It is not necessary to evaluate the integrals. | <input type="checkbox"/> |
| (i) $\int_{-1}^1 \frac{e^x - e^{-x}}{2} dx = 0$. | <input type="checkbox"/> |
| (ii) $\int_0^1 x^6 dx < \int_0^1 x^7 dx$. | <input type="checkbox"/> |
| (iii) $\int_0^\pi \sin^4 x dx > \int_0^\pi \sin 4x dx$. | <input type="checkbox"/> |

Question 2 (15 marks)

(Start a new writing booklet)

(a) Evaluate $\int_0^1 \frac{dx}{(x+1)\sqrt{x+1}}$.

(b) (i) Find a , b , and c such that

$$\frac{x^2 + 5x - 4}{(x-1)(x^2+1)} \equiv \frac{a}{x-1} + \frac{bx+c}{x^2+1}.$$

(ii) Hence find $\int \frac{x^2 + 5x - 4}{(x-1)(x^2+1)} dx$.

(c) Find $\int \frac{dx}{x\sqrt{x^2-1}}$, using $x = \sec \theta$.

(d) (i) Show that $\sqrt{\frac{4-x}{4+x}} = \frac{4-x}{\sqrt{16-x^2}}$.

(ii) Hence or otherwise find $\int_{-2}^2 \sqrt{\frac{4-x}{4+x}} dx$.

(e) Find $\int_0^1 2x \tan^{-1} x dx$.

Marks

[2]

[2]

[2]

[3]

[1]


[2]

[3]

Question 3 (15 marks)

(Start a new writing booklet)

(a)



(i) Find, in terms of z , the complex number represented by the point Q .

(ii) Find, in terms of z , the complex number which represents the vector \vec{QR} .

(iii) If R represents the complex number w , show that $w^3 + z^3 = 0$.

(b) (i) Given that $y = x - \ln(\sec x + \tan x)$, $0 < x < \frac{\pi}{2}$. Show that $\frac{dy}{dx} = 1 - \sec x$.

(ii) Hence show that $x < \ln(\sec x + \tan x)$ for $0 < x < \frac{\pi}{2}$.

(c) It is known that $2+i$ is a root of the equation $x^6 - 7x^4 + 31x^2 - 25 = 0$.

(i) Give a reason why $2-i$ is also a root of the equation.

(ii) Give a reason why $-(2+i)$ is also a root of the equation.

(iii) Find the other three roots, giving reasons (it should not be necessary to use long division).

Marks

[1]

[2]

[2]

[1]

[3]

[1]

[2]

[3]

Question 4 (15 marks)

(Start a new writing booklet)

- (a) The ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ (where $a > b > 1$) has eccentricity $e = 1/2$.

The point $(2, 3)$ lies on the ellipse.

- (i) Find the values of a and b . 2

- (ii) Sketch the graph of the ellipse, showing clearly the intercepts on the axes, the coordinates of the foci, and the equations of the directrices. 2

- (b) (i) Show that $P(2\sqrt{2}\cos\theta, 3\sqrt{2}\sin\theta)$ lies on the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$. 1

- (ii) Show that the slope of the tangent at P is $-\frac{3\cos\theta}{2\sin\theta}$. 2

- (iii) Find the equation of the normal to the ellipse at P . 2

- (iv) Find the value of θ to the nearest degree if the normal passes through the point $(-2\sqrt{2}, 0)$. 2

- (c) At a dinner party there are twelve people, consisting of six married couples. Each of the women wears a different coloured scarf. The husband of each woman has a matching colour tie.

- (i) The dinner takes place at a circular table. Find how many seating arrangements are possible if the women and men are in alternate positions. 2

- (ii) A committee of six is to be formed from the women and their partners, where not more than one of the six colours can be represented. How many such committees are possible?

Marks

2

2

1

2

2

2

Question 5 (15 marks)


(Start a new writing booklet)

Marks

- (a) (i) Prove that for any polynomial $P(x)$, if k is a zero of multiplicity 2, then k is also a zero of $P'(x)$. 2

- (ii) Show that $x = 1$ is a double root of $x^{2n} - nx^{n+1} + nx^{n-1} - 1 = 0$. 2

- (b) The region shown in the diagram, bounded by the curve $y = \frac{x^2}{x^2 + 1}$, the x -axis, and the line $x = 2$, is rotated about the line $x = 4$.



- (i) Using the method of cylindrical shells, show that the volume δV of a shell distant x from the origin is given by: $\delta V \approx 2\pi(4-x) \left(1 - \frac{1}{1+x^2}\right) \delta x$. 2

- (ii) Hence find the volume of the solid. 3

- (c) An object of mass m kg is thrown vertically upwards. Air resistance is given by $R = 0.05mv^2$ where R is in Newtons and v ms⁻¹ is the speed of the object. (Take $g = 9.8$ ms⁻².)

- (i) Explain why the equation of motion is $\ddot{x} = -\left(\frac{196 + v^2}{20}\right)$ where x is the height of the object t seconds after it is thrown. 2

- (ii) If the velocity of projection is 50 ms⁻¹, find the time taken to reach the highest point. 4

Question 6 (15 marks)

(Start a new writing booklet)

- (a) Consider the curve $x^2 - xy + y^2 = 3$.

(i) Show that $\frac{dy}{dx} = \frac{2x-y}{x-2y}$.

Marks

[2]


(ii) Hence find the two stationary points on the curve.

[2]

(iii) Find any values of x where there are vertical tangents.

[1]

(b)



The sketch shows the graph of $y = f(x)$. There is a horizontal asymptote at $y = -1$ and vertical asymptotes at $x = 0$ and $x = -4$. Draw separate sketches of the following:

(i) $y = |f(x)|$

[1]

(ii) $y = \frac{1}{f(x)}$

[2]

(iii) $|y| = f(x)$

[2]

(iv) $y = [f(x)]^2$

[1]

(c) (i) By considering the perfect square $\left(\sqrt{x} - \frac{1}{\sqrt{x}}\right)^2$, show that $x + \frac{1}{x} \geq 2$.

[2]

(ii) For all $a > 0$, $b > 0$, and $c > 0$, find the smallest possible values of

(α) $(a+b)\left(\frac{1}{a} + \frac{1}{b}\right)$

[1]

(β) $(a+b+c)\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)$

[1]

Question 7 (15 marks)

(Start a new writing booklet)

Marks

- (a) Let w be a non-real root of $z^7 - 1 = 0$.

(i) Show that $1 + w + w^2 + w^3 + w^4 + w^5 + w^6 = 0$.

[1]

(ii) Show that $(1+w)(1+w^2)(1+w^4) = 1$.

[1]

(iii) Form a quadratic equation with roots $(w+w^2+w^4)$ and $(w^6+w^5+w^3)$.

[2]

(iv) Sketch on an Argand diagram all seven roots of $z^7 - 1 = 0$.

[1]

- (b) (i) Show that if n is any even positive integer,

then $(1+x)^n + (1-x)^n = 2 \sum_{k=0}^{n/2} \binom{n}{2k} x^{2k}$.

[2]


- (ii) An alphabet consists of the three letters A , B , and C .

(α) Show that the number of words of five letters containing exactly two A s is given by $\binom{5}{2} \times 2^3$.

[1]

(β) Using (b)(i) and (ii)(α), or otherwise, show that if n is an even positive integer, then the number of words of n letters with zero or an even number of A s is given by $\frac{1}{2}(3^n + 1)$.

[3]



A solid has top and bottom faces which are parallel rectangles of dimensions 9×4 units and 3×2 units respectively. The altitude of the solid is 10 units.

- (c) (i) A rectangle of dimensions x and y units is h units from the base. Assuming that x and y are linear functions of h , or otherwise, show that $x = \frac{3h}{5} + 3$ and $y = \frac{h}{5} + 2$.

[2]


- (ii) By considering a thin slice of volume δV , thickness δh and dimensions $x \times y$ units, show that $\delta V = \left(\frac{3h}{5} + 3\right) \left(\frac{h}{5} + 2\right) \delta h$. Hence by integration find the volume V of the solid.

[2]

Question 8 (15 marks)
 (Start a new writing booklet)

Marks

(a)




$ABCD$ is a cyclic quadrilateral. The diagonals AC and BD intersect at right-angles at X . M is the mid-point of BC . MX produced meets AD at N .

- (i) Copy the diagram showing the above information.
- (ii) Show that $\widehat{M}B\widehat{X} = \widehat{M}\widehat{X}B$. 2
- (iii) Show that MN is perpendicular to AD . 3
- (b) Consider the integral $I_n = \int_0^1 x^{2n+1} e^{-x^2} dx$.
 It is given that $0 \leq x^{2n+1} e^{-x^2} \leq 1$ for $0 \leq x \leq 1$.
 - (i) Briefly explain why $0 \leq I_n \leq 1$. 1
 - (ii) Use integration by parts to show that $I_n = -\frac{1}{2e} + nI_{n-1}$, for $n \geq 1$. 3
 - (iii) Show that $I_0 = \frac{1}{2} - \frac{1}{2e}$. 2
 - (iv) Prove by induction that, for all $n \geq 1$,

$$1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!} = e - \frac{2eI_n}{n!}$$
3
 - (v) Deduce that $1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots = e$. 1

End of Paper

XI QUESTION 1



$$(b) |z|^2 = 1 - 2\sqrt{3}i - 3 \\ = -2 - 2\sqrt{3}i \\ 2\bar{z} = -2 - 2\sqrt{3}i = z^2$$


$$(ii) |z| = \sqrt{(-1)^2 + (\sqrt{3})^2} = 2$$

$$\text{Ang } z = \tan^{-1} -\sqrt{3} \\ = 2\pi/3$$

$$(iii) z^3 = 2^3 (\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}) \\ = 8$$

$z^3 - 8 = 0$
z is a root of the eqn

$$(e) \sqrt{(x-1)^2 + (y+1)^2} \leq 4$$



Circle Centre $(1, -1)$ $r = 4$.

$$(d) i = (a+bi)^2 = a^2 + 2abi - b^2 \\ a^2 - b^2 = 0 \quad ab = 1 \text{ equating LHS} \\ (a+bi)^2 = (a^2 - b^2)^2 + 4a^2b^2$$

$$\begin{cases} a^4b^2 = 1 \\ a^2 - b^2 = 0 \end{cases}$$

$$\begin{aligned} \textcircled{1} + \textcircled{2} & 2a^2 = 1 \\ a &= \pm \frac{1}{\sqrt{2}} \end{aligned}$$

$$b = \pm \frac{1}{\sqrt{2}}$$

$$z = \pm \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i \right)$$


(e) (i) True

$f(x)$ is odd with limits $a \rightarrow -a$

(ii) False

for $0 < x \leq \frac{1}{2}, > \frac{1}{2}$

(iii) True



Sydney Boys' High School
Trial HSC 2007 - Mathematics Extension 2

Question 2

$$(a) I = \int_0^1 \frac{dx}{(x+1)\sqrt{x+1}}$$

Let $u = x+1$; when $x=1, u=2$ and $X=0, u=1$
 $du = dx$

$$I = \int_1^2 u^{-3/2} du$$

$$= \left[\frac{u^{-1/2}}{-\frac{1}{2}} \right]_1$$

$$= \left[-2 \left(\frac{1}{\sqrt{u}} - 1 \right) \right]$$

$$= -2\sqrt{2}$$

$$(b) (i) \frac{x^2 + 5x - 4}{(x-1)(x^2+1)} = \frac{a}{x-1} + \frac{bx+c}{x^2+1}$$

Multiplying both sides by the LHS denominator:

$$x^2 + 5x - 4 = a(x^2 + 1) + (bx + c)(x - 1)$$

$$= ax^2 + a + bx^2 - bx + cx - c$$

$$= (a+b)x^2 + (c-b)x + (a-c)$$

Equating coefficients of corresponding powers:

$$x^2: 1 = a+b \quad \text{--- (1)}$$

$$x^1: 5 = -b+c \quad \text{--- (2)}$$

$$x^0: -4 = a - c \quad \text{--- (3)}$$

$$(1)+(2): 6 = a+c \quad \text{--- (4)}$$

$$(3)+(4): 2 = 2a \quad \text{--- (5)}$$

$$\therefore a = 1$$

$$\text{In (1): } 1 = 1+b$$

$$b = 0$$

$$\text{In (3): } -4 = 1 - c$$

$$c = 5$$

$$(ii) \int \frac{x^2 + 5x - 4}{(x-1)(x^2+1)} dx = \int \left[\frac{1}{x-1} + \frac{5}{x^2+1} \right] dx \\ = \ln|x-1| + 5 \tan^{-1} x + C$$

(c)

$$\begin{aligned}
 I &= \int \frac{dx}{x\sqrt{x^2-1}} && \text{Let } x = \sec \theta, \quad dx = \sec \theta \tan \theta d\theta \\
 I &= \int \frac{\sec \theta \tan \theta d\theta}{\sec \theta \sqrt{\sec^2 \theta - 1}} && \text{Alternatively} \\
 &= \int \frac{\sec \theta \tan \theta d\theta}{\sec \theta |\tan \theta|} && I = -\tan^{-1}\left(\frac{1}{\sqrt{x^2-1}}\right) + D \\
 &= \int (\pm 1) d\theta && \text{or} \\
 &= \pm \theta + C && I = \cos^{-1}\left(\frac{1}{|x|}\right) + E \\
 &= \pm \tan^{-1}\sqrt{x^2-1} + C
 \end{aligned}$$

(d) (i) RTP $\sqrt{\frac{4-x}{4+x}} = \frac{4-x}{\sqrt{16-x^2}}$

$$\begin{aligned}
 \text{LHS} &= \sqrt{\frac{(4-x)(4-x)}{(4+x)(4-x)}} \\
 &= \sqrt{\frac{(4-x)^2}{16-x^2}} \\
 &= \frac{4-x}{\sqrt{16-x^2}} \\
 &= \text{RHS (QED)}
 \end{aligned}$$


(ii)

$$\begin{aligned}
 \int_{-2}^2 \sqrt{\frac{4-x}{4+x}} dx &= \int_{-2}^2 \frac{4-x}{\sqrt{16-x^2}} dx \\
 &= \frac{1}{2} \int_{-2}^2 \frac{8-2x}{\sqrt{16-x^2}} dx \\
 &= \frac{1}{2} \int_{-2}^2 \frac{8}{\sqrt{16-x^2}} dx + \frac{1}{2} \int_{-2}^2 \frac{-2x}{\sqrt{16-x^2}} dx \\
 &= \left[\frac{1}{2} \times 8 \times \sin^{-1}\left(\frac{x}{4}\right) \right]_{-2}^2 + \left[\sqrt{16-x^2} \right]_{-2}^2 \\
 &= 4 \left(\sin^{-1}\left(\frac{1}{2}\right) - \sin^{-1}\left(-\frac{1}{2}\right) \right) + (\sqrt{12} - \sqrt{12}) \\
 &= 4 \left(\frac{\pi}{6} - \left(-\frac{\pi}{6}\right) \right) \\
 &= \frac{4\pi}{3}
 \end{aligned}$$

(e)

$$\begin{aligned}
 \int_0^1 2x \tan^{-1} x dx &= \int_0^1 \frac{d}{dx} (x^2) \tan^{-1} x dx \\
 &= \left[x^2 \tan^{-1} x \right]_0^1 - \int_0^1 x^2 \frac{d}{dx} \tan^{-1} x dx \\
 &= \frac{\pi}{4} - \int_0^1 \frac{x^2}{1+x^2} dx \\
 &= \frac{\pi}{4} - \int_0^1 \frac{1+x^2-1}{1+x^2} dx \\
 &= \frac{\pi}{4} - \int_0^1 1 dx + \int_0^1 \frac{1}{1+x^2} dx \\
 &= \frac{\pi}{4} - [x]_0^1 + [\tan^{-1} x]_0^1 \\
 &= \frac{\pi}{4} - 1 + \frac{\pi}{4} \\
 &= \frac{\pi}{2} - 1
 \end{aligned}$$

1007 Ext 2 Q3



$$(i) \theta = i\beta$$

$$(ii) R = \sqrt{\beta \cos \frac{\pi}{3}}$$

$$\therefore R = \beta \cos \frac{\pi}{3} - i\beta \sin \frac{\pi}{3} = \beta \left(\cos \frac{\pi}{3} - i \right)$$

$$(iii) \omega^3 + \beta = \beta^3 \cos \pi + \beta^3$$

$$= -\beta^3 + \beta^3$$

$$= 0.$$


$$(iv) y = x - \ln(\sec x + \tan x)$$

$$\frac{dy}{dx} = 1 - \frac{1}{\sec x + \tan x} \times (\sec x \tan x + \sec^2 x)$$

$$= 1 - \frac{1}{\sec x + \tan x} \sec x (\tan x + \sec x)$$

(1)

$$= 1 - \sec x.$$



$$\theta < x < \frac{\pi}{2}$$

$\omega = \ln(\sec x + \tan x) < 0$ for $0 < x < \frac{\pi}{2}$ (3)

(c) If the polynomial has real coefficients,

$$\frac{1}{2+i} = 2-i \text{ is also a root.}$$

(ii) At the time it was, if x is a root, $-x$ is also a root.

$$\therefore -(2-i) \text{ is also a root.}$$

$$(iii) \text{Sum of roots} = (2+i)(2-i) + (-2+i)(-2-i) + \alpha + \beta.$$

$$\alpha + \beta.$$

$$= 0$$

$$\therefore \alpha = -\beta.$$

$$(2+i)(2-i)(-2+i)\alpha \beta = -25$$

$$\therefore 25 \cdot \alpha \beta = -25$$

$$\therefore \alpha \beta = -1$$

$$\therefore \alpha^2 = -1$$

$$\therefore \alpha^2 = 1$$

$$\therefore \alpha = 1 \quad \text{and } \beta = -1.$$

$$\therefore \text{Other roots are } -(2-i), 1, -1.$$

$$\therefore \omega = \ln(\sec x + \tan x) \text{ is discontinuous.}$$

$$\text{When } x = 0 \quad \sec x = 1 \quad \tan x = 0$$

$$\sec x + \tan x = 1 \quad (\sec x - 1) = 0$$

$$\therefore \omega = 0$$

QUESTION 5

(a) (i)

$$\text{Let } P(x) = (x-k)^2 \cdot Q(x)$$

$$P'(x) = (x-k)^2 \cdot Q'(x) + 2(x-k)Q(x)$$

$$\therefore P'(x) = (x-k) \left[(x-k)Q'(x) + 2Q(x) \right]$$

$$P'(k) = 0 \quad [0 + 2Q(k)] = 0$$

$\Rightarrow k$ is a zero of $P'(x)$.

(ii) Let $P(x) = x^{2n} - nx^{n+1} + nx^{n-1} + 1$

$$\text{Now } P(1) = 1 - n + n + 1 = 0$$

$\therefore x=1$ is a root of $P(x) = 0$

$$P'(x) = 2nx^{2n-1} - n(n+1)x^n + n(n-1)x^{n-2}$$

$$P'(1) = 2n - n^2 - n + n^2 - n$$

$$= 0$$

$$\Rightarrow P'(1) = P'(1) = 0$$

$\therefore x=1$ is a double root of $P(x) = 0$

Vol. solid = $\lim_{\Delta x \rightarrow 0} \sum_{x=0}^2 2\pi(4-x)(1-\frac{1}{1+x^2})\Delta x$

$$V = 2\pi \int_0^2 (4-x) \left(1 - \frac{1}{1+x^2}\right) dx$$

$$V = 2\pi \int_0^2 (4-x)dx - 2\pi \int_0^2 \frac{4-x}{1+x^2} dx$$

$$= 2\pi \left[4x - \frac{x^2}{2} \right]_0^2 - 2\pi \left[4 \tan^{-1} x - \frac{1}{2} \ln(1+x^2) \right]_0^2$$

$$= 2\pi [8-2] - 2\pi \left[4 \tan^{-1} 2 - \frac{1}{2} \ln 5 \right] \text{ using } \int \frac{1}{1+x^2} dx = \tan^{-1} x$$

$$(c) (i) m\ddot{x} = -mg + R$$

$$\therefore m\ddot{x} = -gm - \frac{mv^2}{20}$$

$$\ddot{x} = -9.8 - \frac{v^2}{20}$$

$$\ddot{x} = - \left(\frac{196+v^2}{20} \right)$$

(ii) $m\ddot{x} = -gm - \frac{mv^2}{20}$

$$\therefore \ddot{x} = -9.8 - \frac{v^2}{20}$$

$$\ddot{x} = - \left(\frac{196+v^2}{20} \right)$$

(a) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$\therefore (2, 3)$ is on the ellipse.

$$\therefore \frac{4}{a^2} + \frac{9}{b^2} = 1$$

$$\therefore \frac{9}{b^2} = 1 - \frac{4}{a^2}$$


$$\therefore b^2 = a^2 \left(1 - \frac{4}{a^2}\right)$$

$$\therefore b^2 = \frac{3}{4}a^2$$

$$\therefore a^2 = \frac{3}{4}a^2 - 3$$

$$\therefore a^2 = 16 \Rightarrow a = 4$$

$$b = \sqrt{12} = 2\sqrt{3}$$



(b) $\frac{x^2}{4} + \frac{y^2}{9} = 1$

(ii) $\frac{x}{2} + \frac{2y}{9} \frac{dy}{dx} = 0$

$$\frac{\partial}{\partial x} \left(\frac{x^2}{4} + \frac{y^2}{9} \right) + \frac{\partial}{\partial y} \left(\frac{2y}{9} \frac{dy}{dx} \right) = 0$$

$$\therefore \frac{2\sin\theta}{3} \frac{du}{dx} = -4\cos\theta$$

$$\Rightarrow \frac{du}{dx} = -\frac{3\cos\theta}{2\sin\theta}$$

(i) $\frac{x^2}{4} + \frac{y^2}{9} = 1$

$$\left(\frac{2\sqrt{4\cos^2\theta}}{4} \right)^2 + \left(\frac{3\sqrt{2\sin\theta}}{9} \right)^2$$

$$= \frac{8\cos^2\theta}{4} + \frac{18\sin^2\theta}{9}$$

$$= 2(\cos^2\theta + \sin^2\theta) = 2$$

$$\therefore P(2\sqrt{4\cos^2\theta}, 3\sqrt{2\sin\theta})$$

$$\text{lies on } \frac{x^2}{4} + \frac{y^2}{9} = 1$$

\therefore Coordinate satisfy the equation.

(iii) $m' = \frac{2\sin\theta}{3\cos\theta}$

\therefore equation of normal

$$y - 3\sqrt{2}\sin\theta = \frac{2\sin\theta}{3\cos\theta}(x - 2\sqrt{4\cos^2\theta})$$

(iv) \therefore (3) passes through

$$(-2\sqrt{2}, 0)$$

$$-3\sqrt{2}\sin\theta = -\frac{4\sqrt{2}\sin\theta}{3\cos\theta}(1+4\cos\theta)$$

$$-9\cos\theta = -4 - 4\cos\theta$$

$$\therefore 4\cos\theta = 0.8$$

$$\theta = 37^\circ$$

$$0.646^\circ$$

$$720 \times 120$$

$$86400$$

$$6! \times 5!$$

$$(c)$$

$$(i)$$

$$(ii)$$

$$(iii)$$

$$(iv)$$

$$(v)$$

$$(vi)$$

X2 QUESTION 6

$$(i) 2x - y - x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$$

$$\frac{dy}{dx}(-x+2y) = y - 2x \\ \frac{dy}{dx} = \frac{y-2x}{-x+2y} = \frac{2x-y}{x-2y}$$

$$(ii) \text{ stat. pts } \frac{dy}{dx} = 0 \quad 2x = y$$

$$\text{Sub into eqn } x^2 - 2x^2 + 4x^2 = 3$$

$$x^2 = 1 \quad x = \pm 1$$

$$y = 2x$$

$$\text{stat. pts } (1, 2), (-1, -2)$$

(iii) Vert. tangents when $\frac{dy}{dx}$ undefined.



$$2x - 2y = 0$$

$$x^2 - x^2 + x^2 = 3$$


$$\frac{3x^2}{4} = 3$$

$$x = \pm 1$$

$$(iv) \quad y = |f(x)|$$



$$(c) \quad (y) = f(x)$$



$$(d) \quad y = [f(x)]^2$$



$$(c) \quad \left(\sqrt{x} - \frac{1}{\sqrt{x}}\right)^2 \geq 0$$

$$x - 2 + \frac{1}{x} \geq 0$$

$$(ii) \quad \left(\frac{1}{a} + \frac{1}{b}\right)(a+b)$$

$$= 1 + \frac{a}{b} + \frac{b}{a} + 1$$

$$\frac{a}{b} + \frac{b}{a} \geq 2 \quad \text{from (i)}$$

$$\therefore 2 + \frac{a}{b} + \frac{b}{a} \geq 4$$

$$\text{least value} = 4$$

$$(iii) \quad (a+b+c)\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)$$

$$= 1 + \frac{b}{a} + \frac{c}{a} + \frac{a}{b} + 1 + \frac{a}{c} + \frac{b}{c} + 1$$

$$= 3 + \frac{b}{a} + \frac{a}{b} + \frac{c}{a} + \frac{a}{c} + \frac{b}{c} + \frac{c}{b}$$

$$\geq 3 + 2 + 2 + 2 \quad \text{from (i)}$$

$$\text{least value} = 9$$

Sydney Boys' High School
Trial HSC 2007 - Mathematics Extension 2

Question 7

$$(a) \quad z^7 - 1 = 0$$

(i) $1 + w + w^2 + w^3 + w^4 + w^5 + w^6$ is a GP where $a = 1, r = w, n = 7$.

$$S_7 = \frac{a(1-r^7)}{1-r} \\ = \frac{1-w^7}{1-w} \\ = \frac{1-1}{1-w} \\ = 0$$

$$(ii) \quad (1+w)(1+w^2)(1+w^4) = (1+w^2+w+w^3)(1+w^4) \\ = 1+w^2+w+w^3+w^4+w^5+w^7 \\ = (1+w+w^2+w^3+w^4+w^5+w^6) + w^7 \\ = 0+1 \\ = 1$$

$$(iii) \quad \text{One such equation is the monic quadratic} \\ (z - (w+w^2+w^4))(z - (w^6+w^5+w^3)) = 0$$


Sum of roots = -1 (See Part (i))

$$\begin{aligned} \text{Product of roots} &= (w+w^2+w^4)(w^6+w^5+w^3) \\ &= w^7+w^6+w^4+w^8+w^7+w^5+w^{10}+w^9+w^7 \\ &= 1+w^6+w^4+w+1+w^5+w^3+w^2+1 \\ &= 2+(1+w+w^2+w^3+w^4+w^5+w^6) \\ &= 2 \end{aligned}$$

Thus the equation is $z^2+z+2=0$

(iv)

All angles $\frac{2\pi}{7}$



(b) (i) RTP: $(1+x)^n + (1-x)^n = 2 \sum_{k=0}^{\frac{n}{2}} {}^n C_{2k} x^{2k}$

$$\begin{aligned} LHS &= 1 + {}^n C_1 x + {}^n C_2 x^2 + \dots + {}^n C_{\frac{n}{2}-1} x^{\frac{n}{2}-1} + x^n + 1 - {}^n C_1 x + {}^n C_2 x^2 - \dots - {}^n C_{\frac{n}{2}-1} x^{\frac{n}{2}-1} + x^n \\ &= 2 + 2 {}^n C_2 x^2 + 2 {}^n C_4 x^4 + \dots + 2 x^n \\ &= RHS \quad (\text{QED}) \end{aligned}$$

- (ii) (α) Two As can be arranged in ${}^5 C_2$ ways. The remaining three letters can be chosen from 2 letters each, in 2^3 ways.
Hence the no. of ways overall is ${}^5 C_2 \times 2^3 = 80$.

(β) For 0 As: 2^n
For 2 As: ${}^n C_2 \times 2^{n-2}$
For 4 As: ${}^n C_4 \times 2^{n-4}$
For $n-2$ As: ${}^n C_{n-2} \times 2^2$
For n As: ${}^n C_n \times 2^0 = 1$

\therefore Total

$$\begin{aligned} &= \sum_{k=0}^{\frac{n}{2}} {}^n C_{2k} 2^{2k} \\ &= \frac{1}{2} [(1+2)^n + (1-2)^n] \\ &= \frac{1}{2} [3^n + (-1)^n] \\ &= \frac{1}{2} (3^n + 1) \quad \text{since } n \text{ is even.} \end{aligned}$$


(c) (i) $x = ah + b$
When $h = 0$, $x = 3$ and when $h = 10$, $x = 9$, so
 $3 = b$
 $9 = 10a + b$
 $= 10a + 3$
 $6 = 10a$
 $a = \frac{3}{5}$
 $\therefore x = \frac{3}{5}h + 3$

$$\begin{aligned} y &= ch + d \\ \text{When } h = 0, y = 2 \text{ and when } h = 10, y = 4, \text{ so} \\ 2 &= d \\ 4 &= 10c + d \\ &= 10c + 2 \\ 2 &= 10c \\ c &= \frac{1}{5} \\ \therefore y &= \frac{h}{5} + 2 \end{aligned}$$

(ii) $\delta V = xy\delta h$
Clearly $= \left(\frac{3}{5}h + 3\right)\left(\frac{h}{5} + 2\right)\delta h$
Thus

$$\begin{aligned} V &= \int_0^{10} \left(\frac{3}{5}h + 3\right)\left(\frac{h}{5} + 2\right)dh \\ &= \frac{1}{25} \int_0^{10} (3h+15)(h+10)dh \\ &= \frac{1}{25} \int_0^{10} (3h^2 + 30h + 15h + 150)dh \\ &= \frac{1}{25} \left[h^3 + \frac{45h^2}{2} + 150h \right]_0^{10} \\ &= \frac{1}{25} \left[1000 + \frac{4500}{2} + 1500 \right] \\ &= 190 \text{ unit}^3 \end{aligned}$$

S(1)



D

$$\angle BAC = 90^\circ \text{ (angle subtended at right-angle)}$$

\therefore B, X and C lie on a circle with diameter BC

M is mid-point of BC, M is the centre of the circle

$$\therefore BM = MX \text{ (radius)}$$

$$\therefore MX = CX \quad (\text{base angles of isosceles } \triangle BXC)$$

$$(iii) \quad \text{But } \angle BXM + \angle AXN = 180^\circ \quad (\text{Supplementary})$$

$$\therefore \theta + 90^\circ + \angle AXN = 180^\circ.$$

$$\therefore \angle AXN = (90 - \theta)$$

$$\angle XAN = \angle CHD = \angle CXB \quad (\text{Angle of alternate} \\ \text{chords on same side})$$

$$\angle XAN + \angle AXN = 180^\circ \quad (\text{angle } B + \angle D)$$

$$\therefore \theta + \angle AXN + 90 - \theta = 180^\circ.$$

$$\therefore \angle AXN = 90^\circ$$

$\therefore MN \perp AD$

$$b) \quad I_1 = \int_0^{\pi/2} xe^{-2x} dx$$

$$0 \leq xe^{-2x} \leq 1$$

$$0 \leq \int_0^{\pi/2} xe^{-2x} dx \leq \frac{1}{2}$$

$$\therefore 0 \leq \int_0^{\pi/2} xe^{-2x} dx \leq \frac{1}{2}$$

$$(i) \quad I_1 = \int_0^{\pi/2} xe^{-2x} dx$$

$$= \int_0^{\pi/2} \frac{x}{e^{2x}} dx$$

$$= \left[\frac{1}{2} e^{-2x} \right]_0^{\pi/2} + n \int_0^{\pi/2} x e^{-2x} dx$$

$$(ii) \quad I_1 = \int_0^{\pi/2} xe^{-2x} dx$$

$$= \int_0^{\pi/2} x e^{-2x} dx$$

$$= \int_0^{\pi/2} x e^{-2x} dx + n \int_0^{\pi/2} x e^{-2x} dx$$

$$= \left[\frac{1}{2} e^{-2x} \right]_0^{\pi/2} + n \int_0^{\pi/2} x e^{-2x} dx$$

$$(iii) \quad I_1 = \int_0^{\pi/2} xe^{-2x} dx$$

$$= \int_0^{\pi/2} x e^{-2x} dx$$

$$= \left[\frac{1}{2} e^{-2x} \right]_0^{\pi/2} + n \int_0^{\pi/2} x e^{-2x} dx$$

$$= \left[\frac{1}{2} e^{-2x} \right]_0^{\pi/2} + n \int_0^{\pi/2} x e^{-2x} dx$$

$$(iv) \quad S(1) \equiv I_1 = e^{-\frac{\pi^2}{8}}$$

$$RHS = e^{-\pi^2/8}$$

$$= e + 1 - e$$

$$\approx 1$$

$$\text{Also } e = \lim_{n \rightarrow \infty} e^{\frac{1}{n}} \leq \lim_{n \rightarrow \infty} (1 + \frac{1}{n})^{1/n} \dots + \frac{1}{n}) \leq e.$$

Show $\ln(1 + \frac{1}{k}) \leq \frac{1}{k}$

$$e^{\ln(1 + \frac{1}{k})} = 1 + \frac{1}{k} + \dots + \frac{1}{k} + \frac{1}{k(k+1)} = e - \frac{2e^{-\frac{1}{k+1}}}{k+1}$$

$$\text{L.H.F} = e - \frac{2e^{-\frac{1}{k+1}}}{k+1} + \frac{1}{k(k+1)}$$

$$= e - \frac{1}{k+1} (2e \cdot (\frac{1}{k+1} + \frac{1}{2k+1} \times \frac{1}{k+1} + \frac{1}{(k+1)k}))$$

$$= e - \frac{1}{(k+1)!} (2e \cdot \frac{1}{k+1} + 1) + \frac{1}{(k+1)!}$$

$$= e - \frac{1}{(k+1)!} (2e \cdot \frac{1}{k+1} + 1 - 1)$$

$$= e - \frac{2e \cdot \frac{1}{k+1}}{(k+1)!}$$

$$= L.H.F.$$

If $S(k)$ is true, $S(k+1)$ is true.

So it is true and $\{S(k)\}$ is true if $S(k)$ is true.

By the process of mathematical induction
for all natural $n \geq 0$

$$(V) \quad 0 \leq T_n \leq 1$$

$$\therefore 0 \leq \frac{T_n}{n!} \leq \frac{1}{n!}$$

$$\therefore -\frac{T_n}{n!} \leq -\frac{2e}{n!} \leq 0$$

$$\therefore e - \frac{2e}{n!} \leq 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} \leq e$$

$$\therefore e - \frac{2e}{n!} \leq 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} \leq e$$