

**GIRRAWEEEN HIGH SCHOOL
MATHEMATICS**

Year 12 Extension 2 Task 4

Thursday 9th June 2005

- Instructions: a) Write all your answers on your own paper.
 b) Show all necessary working.
 c) Marks may be deducted for careless or badly arranged work.

Time Allowed: 90 minutes

Question 1 (25 marks)

Marks

Find the following integrals:

- | | |
|--|---|
| (i) $\int \frac{1}{x \log x} dx$ | 3 |
| (ii) $\int \frac{x^2}{1+x} dx$ | 3 |
| (iii) $\int \sin^4 x \cos^3 x dx$ | 4 |
| (iv) $\int_0^{\frac{\pi}{2}} \frac{dx}{1 + \sin x + \cos x}$ | 4 |
| (v) $\int x \tan^{-1} x dx$ | 4 |
| (vi) $\int \frac{3x+1}{x^2+2x+2} dx$ | 3 |
| (vii) $\int \frac{x^2 - x - 21}{(2x-1)(x^2+4)} dx$ | 4 |


Question 2 (13 marks)

- a) (i) If $I_n = \int \tan^n x dx$ for $n \geq 0$ show that $I_n = \frac{1}{n-1} \tan^{n-1} x - I_{n-2}$ for $n \geq 2$ 3
- (ii) Evaluate $\int_0^{\frac{\pi}{4}} \tan^7 x dx$ 3
- b) (i) Prove that $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ 2
- (ii) Consider $f(x) = \frac{1}{1 + \tan x}$ where $0 \leq x \leq \frac{\pi}{2}$ and $f\left(\frac{\pi}{2}\right) = 0$ 2
- Show that $f(x) + f\left(\frac{\pi}{2} - x\right) = 1$
- (iii) Hence evaluate $\int_0^{\frac{\pi}{2}} \frac{1}{1 + \tan x} dx$ 3

Question 3 (22 marks)


Marks

- a) The region between the curve $y = \sin x$, the line $y = 1$ and the y axis is rotated about the line $y = 1$. Using a slicing technique find the volume formed. 5
- b) 5



The diagram shows a cross-sectional slice of a solid whose base is the region enclosed by the circle $x^2 + y^2 = 4$. Each such cross-section of the solid is an equilateral triangle. Find the volume of the solid.

- c) The region bounded by the curve $y = \frac{x^2}{x^2 + 1}$, the x axis and $0 \leq x \leq 2$, is rotated about the line $x = 4$ to form a solid.



- (i) Using the method of cylindrical shells, explain why the volume ΔV of a typical shell distant x units from the origin and with thickness Δx is given by 3

$$\Delta V = 2\pi(4 - x) \left(1 - \frac{1}{1 + x^2}\right) \Delta x$$

- (ii) Hence, find the total volume of the solid formed. 3

- d) (i) Show that the area of a regular hexagon of side s is given by $A = \frac{3\sqrt{3}s^2}{2}$ 2

- (ii) The diagrams below illustrate a dome tent. When erected, the base is a regular hexagon which measures 2 metres from corner to adjacent corner. 4

Flexible exterior poles extend between opposite corners in semi-circle arcs to support the tent.

By taking slices parallel to the base of the tent find the volume enclosed by the tent.



Year 12 Extension 2 Task 3 2005 Solutions

Question 1 (25)

(i) $\int \frac{1}{x \log x} dx$ $u = \log x$
 $= \int \frac{du}{u}$ $du = \frac{dx}{x}$
 $= \log u + c$
 $= \log(\log x) + c$ (3)

(ii) $\int \frac{x^2}{1+x} dx$ $\frac{x-1}{(x+1)(x^2+0x+0)}$
 $= \int \left[\frac{x-1}{x-1} + \frac{1}{1+x} \right] dx$
 $= \frac{1}{2}x^2 - x + \log(1+x) + c$ (3)

(iii) $\int \sin^4 x \cos^3 x dx$
 $= \int \sin^3 x (1 - \sin^2 x) \cos x dx$ $u = \sin x$
 $= \int (u^4 - u^6) du$ $du = \cos x dx$
 $= \frac{1}{5}u^5 - \frac{1}{7}u^7 + c$
 $= \frac{1}{5}\sin^5 x - \frac{1}{7}\sin^7 x + c$ (4)

(iv) $\int \frac{dx}{1 + \sin x \cos x}$ $t = \tan \frac{x}{2}$
 $= \int \frac{\frac{2dt}{1+t^2}}{1 + \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2}}$ $dt = \frac{2dt}{1+t^2}$
 $= \int \frac{2dt}{1+t^2+2t+1-t^2}$
 $= \int \frac{2dt}{4t^2+2t+1-t^2}$
 $= \int \frac{2dt}{(2t+1)^2}$
 $= \left[\log(t+1) \right]_0^1$
 $= \log 2$ (4)

(v) $\int x \tan^{-1} x dx$ $u = \tan^{-1} x$ $v = \frac{1}{2}x^2$
 $= \frac{1}{2}x^2 \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} dx$ $du = \frac{dx}{1+x^2}$ $dv = dx$
 $= \frac{1}{2}x^2 \tan^{-1} x - \frac{1}{2} \int \left[1 - \frac{1}{1+x^2} \right] dx$ (4)

(vi) $\int \frac{3x+1}{x^2+2x+2} dx$
 $= \frac{3}{2} \int \frac{2x+2}{x^2+2x+2} dx - 2 \int \frac{dx}{(x+1)^2+1}$
 $= \frac{3}{2} \log(x^2+2x+2) - 2 \tan^{-1}(x+1) + c$ (3)

(vii) $\int \frac{x^2-x-2}{(2x-1)(x^2+4)} dx$
 $\frac{A}{(2x-1)} + \frac{Bx+C}{(x^2+4)} = \frac{x^2-x-2}{(2x-1)(x^2+4)}$
 $A(x^2+4) + (Bx+C)(2x-1) = x^2-x-2$
 $\frac{17}{4}A = \frac{66}{4}$ $4A - C = -21$
 $A = -5$ $-2B - C = -21$
 $C = 41$
 $\frac{x=1}{5A+B+C = -21}$
 $-25+B+1 = -21$
 $B = 3$

$\int \frac{x^2-x-2}{(2x-1)(x^2+4)} dx$
 $= \int \left[\frac{-5}{2x-1} + \frac{3x+1}{x^2+4} \right] dx$
 $= \int \left[\frac{-5}{2x-1} + \frac{3}{2} \cdot \frac{2x}{x^2+4} + \frac{1}{x^2+4} \right] dx$
 $= \frac{-5}{2} \log(2x-1) + \frac{3}{2} \log(x^2+4) + \frac{1}{2} \tan^{-1} \frac{x}{2} + c$ (4)

Question 2 (13)

(i) $I_n = \int \tan^n x dx$
 $= \int \tan^{n-2} x \cdot (\sec^2 x - 1) dx$
 $= \int \tan^{n-2} x \sec^2 x dx - \int \tan^{n-2} x dx$
 $u = \tan x$
 $du = \sec^2 x dx$
 $= \int u^{n-2} du - I_{n-2}$
 $= \frac{1}{n-1} u^{n-1} - I_{n-2}$
 $= \frac{1}{n-1} \tan^{n-1} x - I_{n-2}$ (3)


(ii) $\int \tan^7 x dx$
 $= \int \left[\frac{1}{6} \tan^6 x \right] dx - I_5$
 $= \frac{1}{6} - \left[\frac{1}{4} \tan^4 x \right]_0^{\frac{\pi}{4}} + I_3$
 $= \frac{1}{6} - \frac{1}{4} + \left[\frac{1}{2} \tan^2 x \right]_0^{\frac{\pi}{4}} - I_1$
 $= \frac{1}{6} - \frac{1}{4} + \frac{1}{2} - \int \tan x dx$
 $= \frac{5}{12} + \left[\log \cos x \right]_0^{\frac{\pi}{4}}$
 $= \frac{5}{12} + \log(\sqrt{2})$ (3)

(b) (i) $\int_a^a f(x) dx$ $u = a-x$
 $du = -dx$
 $x=0, u=a$
 $x=a, u=0$
 $= - \int_a^0 f(a-u) du$
 $= \int_0^a f(a-u) du$
 $= \int_0^a f(a-x) dx$ (2)

(ii) $f(x) + f\left(\frac{\pi}{2}-x\right)$
 $= \frac{1}{1+\tan x} + \frac{1}{1+\tan\left(\frac{\pi}{2}-x\right)}$
 $= \frac{1}{1+\tan x} + \frac{1}{1+\cot x}$
 $= \frac{1}{1+\tan x} + \frac{1}{1+\frac{1}{\tan x}}$
 $= \frac{1}{1+\tan x} + \frac{\tan x}{\tan x+1}$
 $= \frac{1}{1+\tan x} + \frac{\tan x}{1+\tan x}$
 $= \frac{1+\tan x}{1+\tan x}$
 $= 1$ (2)

(iii) $\int_0^{\frac{\pi}{2}} \frac{1}{1+\tan x} dx$
 $= \int_0^{\frac{\pi}{2}} \frac{1}{1+\tan(\frac{\pi}{2}-x)} dx$
 $= \int_0^{\frac{\pi}{2}} \left[1 - \frac{1}{1+\tan x} \right] dx$
 $\therefore 2 \int_0^{\frac{\pi}{2}} \frac{1}{1+\tan x} dx = \int_0^{\frac{\pi}{2}} dx$
 $\int_0^{\frac{\pi}{2}} \frac{1}{1+\tan x} dx = \frac{1}{2} \left[x \right]_0^{\frac{\pi}{2}}$
 $= \frac{\pi}{4}$ (3)

Question 3



a)

$$A(x) = \pi (1 - \sin x)^2$$

$$\Delta V = \pi (1 - \sin x)^2 \Delta x$$

$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=0}^{\pi/2} \pi (1 - \sin x)^2 \Delta x$$



$$= \pi \int_0^{\pi/2} (1 - 2\sin x + \sin^2 x) dx$$

$$= \pi \left[\frac{3}{2}x + 2\cos x - \frac{1}{4}\sin 2x \right]_0^{\pi/2}$$

$$= \pi \left(\frac{3\pi}{4} + 0 - 0 - 2 + 0 \right)$$

$$= \pi \left(\frac{3\pi}{4} - 2 \right)$$

$$= \frac{3\pi^2 - 8\pi}{4} \text{ units}^3 \quad \textcircled{5}$$



$$A(x) = \frac{1}{2}(2y)(\sqrt{3}y)$$

$$= \frac{\sqrt{3}}{2}y^2$$

$$= \frac{\sqrt{3}}{2}(4-x^2)$$

$$\Delta V = \frac{\sqrt{3}}{2}(4-x^2)\Delta x$$


$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=1}^2 \frac{\sqrt{3}}{2}(4-x^2)\Delta x$$

$$= 2\sqrt{3} \int_1^2 (4-x^2) dx$$

$$= 2\sqrt{3} \left[4x - \frac{1}{3}x^3 \right]_1^2$$

$$= 2\sqrt{3} \left(8 - \frac{8}{3} \right)$$

$$= \frac{32\sqrt{3}}{3} \text{ units}^3 \quad \textcircled{5}$$



$$A(x) = 2\pi(4-x) \left(\frac{x^2}{x^2+1} \right)$$

$$= 2\pi(4-x) \left(1 - \frac{1}{x^2+1} \right)$$

$$\Delta V = 2\pi(4-x) \left(1 - \frac{1}{x^2+1} \right) \Delta x$$

(ii)


$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=0}^2 2\pi(4-x) \left(1 - \frac{1}{x^2+1} \right) \Delta x$$

$$= 2\pi \int_0^2 \left[4-x - \frac{4}{x^2+1} + \frac{x}{x^2+1} \right] dx$$

$$= 2\pi \left[4x - \frac{1}{2}x^2 - 4 \tan^{-1} x + \frac{1}{2} \log(x^2+1) \right]_0^2$$

$$= 2\pi (8 - 2 - 4 \tan^{-1} 2 + \frac{1}{2} \log 5 - 0)$$

$$= 2\pi (6 - 4 \tan^{-1} 2 + \frac{1}{2} \log 5) \text{ units}^3 \quad \textcircled{5}$$



$$A = 6 \times \frac{1}{2}(s)(s) \sin 60^\circ$$

$$= \frac{3\sqrt{3}}{2}s^2$$

(ii)

$$A(y) = \frac{3\sqrt{3}}{2}x^2$$

$$= \frac{3\sqrt{3}}{8}(4-y^2)$$

$$\Delta V = \frac{3\sqrt{3}}{8}(4-y^2)\Delta y$$

$$V = \lim_{\Delta y \rightarrow 0} \sum_{y=0}^2 \frac{3\sqrt{3}}{8}(4-y^2)\Delta y$$

$$= \frac{3\sqrt{3}}{8} \int_0^2 (4-y^2) dy$$

$$= \frac{3\sqrt{3}}{8} \left[4y - \frac{1}{3}y^3 \right]_0^2$$

$$= \frac{3\sqrt{3}}{8} \left(8 - \frac{8}{3} \right)$$

$$= \frac{8\sqrt{3}}{3} \text{ m}^3 \quad \textcircled{5}$$