Concavity

The second derivative measures the change in slope with respect to x, this is known as **concavity**

If f''(x) > 0, the curve is concave up If f''(x) < 0, the curve is concave down If f''(x) = 0, possible point of inflection



Turning Points

All turning points are stationary points.

If f''(x) > 0, minimum turning point If f''(x) < 0, maximum turning point

e.g. Find the turning points of $y = x^3 + x^2 - x + 1$

$$\frac{dy}{dx} = 3x^2 + 2x - 1$$
$$\frac{d^2 y}{dx^2} = 6x + 2$$

Stationary points occur when $\frac{dy}{dx} = 0$

i.e.
$$3x^{2} + 2x - 1 = 0$$

 $(3x - 1)(x + 1) = 0$
 $x = \frac{1}{3}$ or $x = -1$

when
$$x = -1$$
, $\frac{d^2 y}{dx^2} = 6(-1) + 2$
 $= -4 < 0$
 $\therefore (-1,2)$ is a maximum turning point
when $x = \frac{1}{3}$, $\frac{d^2 y}{dx^2} = 6\left(\frac{1}{3}\right) + 2$
 $= 4 > 0$
 $\therefore \left(\frac{1}{3}, \frac{22}{27}\right)$ is a minimum turning point

Inflection Points

A point of inflection is where there is a **change in concavity**, to see if there is a change, check either side of the point.

e.g. Find the inflection point(s) of $y = 4x^3 + 6x^2 + 2$

$$\frac{dy}{dx} = 12x^2 + 12x \qquad \qquad \frac{d^2y}{dx^2} = 24x + 12$$

Possible points of inflection occur when $\frac{d^2 y}{dx^2} = 0$

i.e. 24x + 12 = 0 $x = -\frac{1}{2}$

: there is a change in concavity

 $\therefore \left(-\frac{1}{2},3\right)$ is a point of inflection

$$\begin{array}{c|c} x & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ \frac{d^2 y}{dx^2} & \bigwedge^{(-12)} & \frac{0}{\sqrt{2}} \end{array}$$

 dx^3

If
$$\frac{d^3 y}{dx^3} \neq 0$$
, then it is a point of inflection
 $e.g.\frac{d^3 y}{dx^3} = 24$
when $x = -\frac{1}{2}, \frac{d^3 y}{dx^3} = 24 \neq 0$
 $\therefore \left(-\frac{1}{2},3\right)$ is a point of inflection
Horizontal Point of Inflection; $\frac{dy}{dx} = 0$ $\frac{d^2 y}{dx^2} = 0$ $\frac{d^3 y}{dx^3} \neq 0$

dx