



**SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS**

**2010
HIGHER SCHOOL CERTIFICATE
TRIAL PAPER**

Mathematics Extension 2

General Instructions

- Reading Time – 5 Minutes
- Working time – 3 Hours
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- Each Question is to be returned in a separate bundle.
- All necessary working should be shown in every question.
- Answer in simplest exact form unless otherwise instructed.

Total Marks – 120

- Attempt questions 1 – 8
- Examiner: *A.M.Gainford*
- NOTE: This is a trial paper only and does not necessarily reflect the content or format of the final Higher School Certificate examination paper for this subject.

Question 1. (15 marks) (Start a new answer sheet.)

	Marks
(a) Evaluate	2
	$\int_0^3 \frac{x}{\sqrt{x^2 + 16}} dx .$
(b) Find	1
	$\int (\cos^2 x - \sin^2 x) dx .$
(c) Use integration by parts to find	2
	$\int xe^{-x} dx .$
(d) (i) Find real numbers a and b such that	2
	$\frac{1-3x}{x^2-3x+2} = \frac{a}{x-1} + \frac{b}{x-2} .$
(ii) Hence find	2
	$\int \frac{1-3x}{x^2-3x+2} dx .$
(e) Evaluate	2
	$\int_0^{\frac{\pi}{2}} \frac{1}{\cos x + 2} dx .$
(f) (i) If $I_n = \int_0^{\frac{\pi}{4}} \tan^n x dx , n = 1, 2, 3, \dots$	2
	show that $I_n + I_{n-2} = \frac{1}{n-1}, n = 2, 3, 4, \dots$
(ii) Hence evaluate	2
	$\int_0^{\frac{\pi}{4}} \tan^5 x dx .$

Question 2. (15 marks) (Start a new answer sheet.)

- | | Marks |
|--|-------|
| (a) If $u = 3 - 4i$ and $v = 2 - 2i$ find | 4 |
| (i) $u\bar{v}$ | |
| (ii) \sqrt{u} | |
| (iii) v in modulus-argument form. | |
| (iv) v^4 using De Moivre's theorem. | |
|
 | |
| (b) On an Argand diagram shade the region that is satisfied by both the conditions | 2 |
| $3 \leq z - 4i \leq 4$ and $-\frac{\pi}{4} < \arg(z - 4i) < \frac{\pi}{4}$ | |
|
 | |
| (c) Sketch, on separate Argand diagrams, the locus of the complex number z satisfying | 4 |
| (i) $z^2 - (\bar{z})^2 = i$ | |
| (ii) $ z - 1 = \operatorname{Re}(z)$ | |
|
 | |
| (d) It is given that $z = \cos \theta + i \sin \theta$ where $0 < \arg z < \frac{\pi}{2}$. | 5 |
| (i) Show that $z + 1 = 2 \cos \frac{\theta}{2} \left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right)$ and express $z - 1$ in modulus-argument form. | |
| (ii) Hence show that $\operatorname{Re} \left(\frac{z-1}{z+1} \right) = 0$. | |


Question 3. (15 marks) (Start a new answer sheet.)

- | | Marks |
|--|--------------|
| (a) (i) Show that $z = 1+i$ is a root of $z^2 - (3-2i)z + (5-i) = 0$. | 3 |
| (ii) Find the other root of the equation. | |
| (b) If α , β and γ are roots of the equation $x^3 + qx - 2 = 0$ find, in terms of q ,
the monic cubic polynomial equation whose roots are α^2 , β^2 and γ^2 . | 3 |
| (c) (i) Use De Moivre's theorem to find $\cos 5\theta$ in terms of powers of $\cos \theta$. | 6 |
| (ii) Use the result in (i) to solve the equation
$16x^4 - 20x^2 + 5 = 0$ | |
| (d) If ω represents one of the complex roots of the equation $z^3 - 1 = 0$ | |
| (i) Show that $1 + \omega + \omega^2 = 0$. | |
| (ii) Evaluate $(1 - \omega^8)(1 - \omega^4)(1 - \omega^2)(1 - \omega)$. | |

Question 4 (15 marks) (Start a new answer sheet.)

(a) The graph of $y = f(x)$ is sketched below.

There is a stationary point at $(0, 1)$.



Use this graph to sketch the following, on separate diagrams, showing essential features.

2

(i) $y = f\left(\frac{x}{2}\right)$

2

(ii) $y = x + f(x)$

2

(iii) $y = \frac{1}{f(x)}$

2

(iv) $y = f\left(\frac{1}{x}\right)$

(b) (i) Find $\int \frac{1}{x^2 \sqrt{9-x^2}} dx$, using the substitution $x = 3\cos\theta$.

4

(ii) Evaluate $\int_1^e x^3 \log_e x dx$.

(c) Given that the quartic polynomial $p(x) = x^4 - 5x^3 - 9x^2 + 81x - 108$ has a zero of multiplicity 3, factorise $p(x)$ completely, and find all its zeroes.

3

Question 5 (15 marks) (Start a new answer sheet.)

- | | Marks |
|--|--------------|
| (a) A particle is moving under gravity in a fluid which exerts a resistance to its motion, per unit mass, k times its speed (k is constant). | |
| (i) If the particle falls vertically from rest, show that its terminal velocity
is $V_T = \frac{g}{k}$, where g is acceleration due to gravity. | 2 |
| (ii) If the particle is projected vertically upward with velocity V_T show that after time t seconds | 6 |
| (α) its speed is $V_T (2e^{-kt} - 1)$ | |
| (β) its height above the starting point is $\frac{1}{k}V_T (2 - 2e^{-kt} - kt)$ | |
| (iii) Hence find an expression for the greatest height reached in terms of V_T and k . | 2 |
| (b) A box contains 6 white balls and 2 black balls. Balls are selected at random, one at a time, and not replaced. A note is kept of the number, X , of the draw which first yields a black ball. If this experiment is repeated many times, find: | 5 |
| (i) the most probable value of X ; | |
| (ii) the probability that $X > 4$. | |


Question 6 (15 marks) (Start a new answer sheet.)

- (a) A council has 14 councillors: 6 Labor, 5 Liberal and 3 Independents. 6
Five councilors are chosen at random to form a committee.

- (i) (α) How many different committees can be formed?
(β) Find the probability that the committee will have a majority of Labor councilors.
- (ii) (α) Show that the number of different committees which can be formed with at least one councilor from each of the groups Labor, Liberal, and Independent is 1365.
(β) Given that the committee contains at least one councilor from each of the groups Labor, Liberal, and Independent, find the probability that the committee will have a majority of Labor councilors.

- (b) The circle $x^2 + y^2 = 4$ is rotated about the line $x = 5$ to form a torus. Use the method of cylindrical shells to prove that the volume of the solid is $40\pi^2$ cubic units. 4

- (c) The solid drawn at right has a circular base of radius 3 units in the horizontal plane. Vertical cross-sections perpendicular to the diameter along the x -axis are equilateral triangles.



- (i) A vertical slice of width Δa is positioned at the point where $x = a$. 3
If the volume of the slice is ΔV , show that $\Delta V = \sqrt{3}(9 - a^2)\Delta a$.
- (ii) Hence determine the volume of the solid. 2

Question 7 (15 marks) (Start a new answer sheet.)

- (a) On polling day in Rock Island City the ratio of electoral votes in the only four polling booths A, B, C, and D was 5:4:3:8 respectively. The percentages of votes for Mr Jones in these booths were 60%, 50%, 40%, and 70% respectively.

4

- (i) Find the probability that a voter chosen at random voted for Mr Jones.
(ii) If ten voters of this city were chosen at random, find the probability that Mr Jones gained
(α) at least 8 votes
(β) no more than 2 votes.


- (b) The equation $e^{2x} \log_e y = 3$ implicitly defines y as a function of x .

3

Find $\frac{dy}{dx}$ as a function of y .

(c)

8



In the diagram above, P , Q , and R are the midpoints of the sides BC , CA , and AB respectively of a triangle ABC . The circle drawn through the points P , Q , and R meets the sides BC , CA , and AB again at X , Y , and Z respectively.

Copy the diagram to your answer sheet.

- (i) Briefly explain why $RPCQ$ is a parallelogram.
(ii) Show that $\triangle XCQ$ is isosceles.
(iii) Show that $AX \perp BC$.

Question 8 (15 marks) (Start a new answer sheet.)

(a) Five women and four men are to be seated at a round table.

5

- (i) In how many ways may this be done without restrictions?
- (ii) In how many ways may this be done if no two men are to be seated together?
- (iii) If one man and one woman are a married couple, what is the probability that they are seated together, given the conditions of part (ii)?

(b) One root of the equation $x^3 + ax^2 + bx + c = 0$ is equal to the sum of the other two roots.

4

Show that $a^3 - 4ab + 8c = 0$.

(c) (i) Graph, in the same xy -plane, the curves

6

$$y = x^{-\frac{2}{3}}, x > 0 \text{ and } y = (x-1)^{-\frac{2}{3}}, x > 1$$

(ii) Hence, or otherwise, given the sum S , where

$$S = 1 + \frac{1}{\sqrt[3]{2^2}} + \frac{1}{\sqrt[3]{3^2}} + \dots + \frac{1}{\sqrt[3]{(10^9)^2}}, \text{ find the two consecutive integers between which the sum } S \text{ lies.}$$

This is the end of the paper.

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right) \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Sydney Boys Extension 2 2010

Question 1

a) $\int_0^3 \frac{x}{\sqrt{x^2+16}} dx$

$$= \left[\sqrt{x^2+16} \right]_0^3$$

$$= 5 - 4$$

$$= \underline{\underline{1}}$$

b) $\int (\cos^2 x - \sin^2 x) dx$

$$= \int \cos 2x dx$$

$$= \underline{\underline{\frac{1}{2} \sin 2x + c}}$$

c) $I = \int x e^{-x} dx$

$$u = x$$

$$v = -e^{-x}$$

$$du = dx$$

$$dv = e^{-x} dx$$

$$I = -xe^{-x} + \int e^{-x} dx$$

$$= \underline{\underline{-xe^{-x} - e^{-x} + c}}$$

d) (i) $a(x-2) + b(x-1) = 1-3x$

$$\frac{x=2}{b} = -5$$

$$\frac{x=1}{-a} = -2$$

$$a = 2$$

$$\therefore \underline{\underline{a=2, b=-5}}$$

(ii) $\int \frac{1-3x}{x^2-3x+2} dx$

$$= \int \left[\frac{2}{x-1} - \frac{5}{x-2} \right] dx$$

$$= \underline{\underline{2 \log(x-1) - 5 \log(x-2) + c}}$$

e) $\int_0^{\frac{\pi}{2}} \frac{dx}{\cos x + 2}$

$$t = \tan \frac{x}{2}$$

$$dx = \frac{2dt}{1+t^2}$$

$$x=0, t=0$$

$$x=\frac{\pi}{2}, t=1$$

$$= \int_0^1 \frac{2dt}{1-t^2+2+2t^2}$$

$$= \int_0^1 \frac{2dt}{3+t^2}$$

$$= \frac{2}{\sqrt{3}} \left[\tan^{-1} \frac{t}{\sqrt{3}} \right]_0^1$$

$$= \frac{2}{\sqrt{3}} \left(\frac{\pi}{6} - 0 \right) = \underline{\underline{\frac{\pi}{3\sqrt{3}}}}$$

f) $I_n = \int_0^{\frac{\pi}{4}} \tan^n x dx$

$$I_n + I_{n-2}$$

$$= \int_0^{\frac{\pi}{4}} (\tan^n x + \tan^{n-2} x) dx$$

$$= \int_0^{\frac{\pi}{4}} \tan^{n-2} x (\tan^2 x + 1) dx$$

$$= \int_0^{\frac{\pi}{4}} \tan^{n-2} x \sec^2 x dx$$

$$= \frac{1}{n-1} \left[\tan^{n-1} x \right]_0^{\frac{\pi}{4}}$$

$$= \frac{1}{n-1} (1-0)$$

$$= \underline{\underline{\frac{1}{n-1}}}$$

(ii) $\int_0^{\frac{\pi}{4}} \tan^5 x dx = I_5$

$$= \frac{1}{4} - I_3$$

$$= \frac{1}{4} - \frac{1}{2} + I_1$$

$$= -\frac{1}{4} + \int_0^{\frac{\pi}{4}} \tan x dx$$

$$= -\frac{1}{4} - \left[\log \cos x \right]_0^{\frac{\pi}{4}}$$

$$= -\frac{1}{4} - \log \frac{1}{\sqrt{2}} + \log 1$$

$$= \underline{\underline{-\frac{1}{4} - \log \frac{1}{\sqrt{2}}}}$$

Question 2

a) (i) $u\bar{v} = (3-4i)(2+2i)$

$$= 6+6i-8i+8$$

$$= \underline{\underline{14-2i}}$$

(ii) $\sqrt{u} = \sqrt{3-4i}$

$$a^2 - b^2 = 3 \quad 2ab = -4$$

$$a^2 - \frac{4}{a^2} = 3 \quad b = -\frac{2}{a}$$

$$a^4 - 3a^2 - 4 = 0$$

$$(a^2 - 4)(a^2 + 1) = 0$$

$$a^2 = 4 \quad \text{or} \quad a^2 = -1$$

$a = \pm 2$ or $a = \pm i$ no real solutions


$$\therefore \underline{\underline{\sqrt{u} = \pm(2-i)}}$$

(iii) $|v| = \sqrt{2^2+2^2} = 2\sqrt{2}$ $\arg v = \tan^{-1} \frac{-2}{2} = -\frac{\pi}{4}$

$$v = 2\sqrt{2} \operatorname{cis} \left(-\frac{\pi}{4} \right)$$

(iv) $v^4 = (2\sqrt{2})^4 \operatorname{cis} (-\pi)$

$$= \underline{\underline{-64}}$$



c) (i) $z^2 - (\bar{z})^2 = i$
 $4xy = 1$
 $xy = \frac{1}{4}$


(ii) $|z-1| = \operatorname{Re}(z)$

$$(x-1)^2 + y^2 = x^2$$

$$x^2 - 2x + 1 + y^2 = x^2$$

$$y^2 = 2x - 1$$

$$y^2 = 2(x - \frac{1}{2})$$



d) $z+1$
 $= \cos\theta + 1 + i\sin\theta$
 $= 2\cos^2\frac{\theta}{2} - 1 + 1 + 2i\sin\frac{\theta}{2}\cos\frac{\theta}{2}$
 $= 2\cos\frac{\theta}{2}(\cos\frac{\theta}{2} + i\sin\frac{\theta}{2})$

$$z-1$$

$$= \cos\theta - 1 + i\sin\theta$$

$$= 1 - 2\sin^2\frac{\theta}{2} - 1 + 2i\sin\frac{\theta}{2}\cos\frac{\theta}{2}$$

$$= -2\sin\frac{\theta}{2}(\sin\frac{\theta}{2} - i\cos\frac{\theta}{2})$$

$$\frac{z-1}{z+1} = \frac{-2\sin\frac{\theta}{2}(\sin\frac{\theta}{2} - i\cos\frac{\theta}{2})}{2\cos\frac{\theta}{2}(\cos\frac{\theta}{2} + i\sin\frac{\theta}{2})} \times \frac{\cos\frac{\theta}{2} - i\sin\frac{\theta}{2}}{\cos\frac{\theta}{2} + i\sin\frac{\theta}{2}}$$

$$= -\frac{\sin\frac{\theta}{2}(\sin^2\frac{\theta}{2} - i\sin\theta - i\cos^2\frac{\theta}{2} - \sin\theta\cos\frac{\theta}{2})}{\cos\frac{\theta}{2}(\cos^2\frac{\theta}{2} + \sin^2\frac{\theta}{2})}$$

$$= \frac{\sin\frac{\theta}{2}(\sin^2\frac{\theta}{2} + \cos^2\frac{\theta}{2})i}{\cos\frac{\theta}{2}}$$

$$= i\tan\frac{\theta}{2}$$

$$\therefore \operatorname{Re}\left(\frac{z-1}{z+1}\right) = 0$$

Question 3

a) $P(1+i)$

$$(1+i)^2 - (3-2i)(1+i) + 5-i$$

$$= 1+2i-1-3-3i+2i-2+5-i$$

$$= 0$$

$\therefore 1+i$ is a root

$\arg(z-4i) = \frac{\pi}{4}$ (i) $x+\beta = 3-2i$
 $1+i+\beta = 3-2i$
 $\beta = 2-3i$

b) let $y = x^2 \Rightarrow x = y^{\frac{1}{2}}$

$$y^{\frac{3}{2}} + q_1 y^{\frac{1}{2}} - 2 = 0$$

$$y^{\frac{1}{2}}(y + q_1) = 2$$

$$y(y^2 + 2q_1 y + q_1^2) = 2$$

$$y^3 + 2q_1 y^2 + q_1^2 y - 2 = 0$$

c) $\operatorname{cis} 5\theta = (\operatorname{cis}\theta)^5$

$$= \cos^5\theta + 5\cos^4\theta - 10\cos^3\theta + 10\cos^2\theta - 5\cos\theta - 1$$

equating real parts

$$\cos 5\theta = \cos^5\theta - 10\cos^3\theta \sin^2\theta + 5\cos\theta \sin^4\theta$$

$$= \cos^5\theta - 10\cos^3\theta(1-\cos^2\theta) + 5\cos\theta(1-\cos^2\theta)^2$$

$$= \cos^5\theta - 10\cos^3\theta + 10\cos^5\theta + 5\cos\theta$$

$$- 10\cos^3\theta + 5\cos^5\theta$$

$$= 16\cos^5\theta - 20\cos^3\theta + 5\cos\theta$$

(ii) $16x^4 - 20x^2 + 5 = 0$
let $x = \cos\theta$

$$\frac{\cos 5\theta}{\cos\theta} = 0$$

$$\cos 5\theta = 0, \cos\theta \neq 0$$

$$5\theta = \pm \pi k, k = 1, 2$$





$$\theta = \pm \frac{\pi}{5}, \pm \frac{2\pi}{5}$$

$$x = \cos\frac{\pi}{5}, -\cos\frac{\pi}{5}, \cos\frac{2\pi}{5}, -\cos\frac{2\pi}{5}$$

d) a) $w^3 - 1 = 0$
 $(w-1)(w^2 + w + 1) = 0$
 $w=1$ or $w^2 + w + 1 = 0$
not a solution as w is complex root

(ii) $(1-w^8)(1-w^4)(1-w^2)(1-w)$
 $= (1-w^2)(1-w)(1-w^2)(1-w)$
 $= [(1-w^2)(1-w)]^2$
 $= (1-w-w^2+w^3)^2$
 $= (2-w-w^2)^2$
 $= (3-1-w-w^2)^2$
 $= 3^2$
 $= \underline{9}$

Question 4



b) (i) $\int \frac{dx}{x^2 \sqrt{9-x^2}}$
 $\quad \quad \quad x = 3\cos\theta \quad dx = -3\sin\theta d\theta$
 $\quad \quad \quad$
 $= \int \frac{-3\sin\theta d\theta}{9\cos^2\theta \cdot 3\sin\theta}$
 $= -\frac{1}{9} \int \sec^2\theta d\theta$
 $= -\frac{1}{9} \tan\theta + C$
 $= -\frac{x}{9\sqrt{9-x^2}} + C$

(ii) $\int_1^e x^3 \log x \, dx$

$$\begin{aligned} u &= \log x & v &= \frac{1}{4}x^4 \\ du &= \frac{dx}{x} & dv &= x^3 \, dx \\ &= \left[\frac{1}{4}x^4 \log x \right]_1^e - \frac{1}{4} \int_1^e x^3 \, dx \\ &= \frac{1}{4}e^4 - \frac{1}{16} [x^4]_1^e \\ &= \frac{1}{4}e^4 - \frac{1}{16}e^4 + \frac{1}{16} \\ &= \underline{\underline{\frac{3}{16}e^4 + \frac{1}{16}}} \end{aligned}$$

c) $p(x) = x^4 - 5x^3 - 9x^2 + 81x - 108$

$p'(x) = 4x^3 - 15x^2 - 18x + 81$

$$\begin{aligned} p''(x) &= 12x^2 - 30x - 18 \\ &= 6(2x^2 - 5x - 3) \\ &= 6(2x+1)(x-3) \end{aligned}$$

$p(3) = p'(3) = p''(3) = 0$

$\therefore p(x) = \underline{\underline{(x-3)^3(x+4)}}$

zeros are 3, 3, 3 and -4

Question 5

a)

$$m\ddot{x} = mg - m\dot{x}\ddot{x}$$

$$\ddot{x} = g - kx$$

terminal velocity occurs when $\ddot{x} = 0$

$$\therefore g - kx = 0$$

$$x = \frac{g}{k}$$

(ii)

$$m\ddot{x} = -mg - m\dot{x}\ddot{x}$$

$$\ddot{x} = -g - kv$$

$$\frac{dv}{dt} = -g - kv$$

$$\int_0^t dt = - \int_{v_T}^v \frac{dv}{g + kv}$$

$$t = \left[-\frac{1}{k} \log(g + kv) \right]^{v_T}$$

$$= -\frac{1}{k} \log \left(\frac{g + kv_T}{g + kv_0} \right)$$

$$-kt = \log \left(\frac{g + kv_T}{g + kv_0} \right)$$

$$e^{-kt} = \frac{g + kv_0}{g + kv_T}$$

$$= \frac{\frac{g}{k} + v_0}{\frac{g}{k} + v_T}$$

$$= \frac{v_T + v_0}{2v_T}$$

$$2v_T e^{-kt} = v_T + v$$

$$v = 2v_T e^{-kt} - v_T$$

$$= \underline{\underline{v_T (2e^{-kt} - 1)}}$$

(B) $\frac{dx}{dt} = v_T (2e^{-kt} - 1)$

$$\int_0^x dx = v_T \int_0^t (2e^{-kt} - 1) dt$$

$$x = v_T \left[-\frac{2}{k} e^{-kt} - t \right]_0^t$$

$$= v_T \left(-\frac{2}{k} e^{-kt} - t + \frac{2}{k} \right)$$

$$= \underline{\underline{\frac{1}{k} v_T (2 - 2e^{-kt} - kt)}}$$

(iii) greatest height occurs when $v=0$

$$\therefore v_T (2e^{-kt} - 1) = 0$$


$$e^{-kt} = \frac{1}{2}$$

$$-kt = \log \frac{1}{2}$$

$$t = -\frac{1}{k} \log \frac{1}{2} = \frac{1}{k} \log 2$$

$$x = v_T \left(2 - 2 \left(\frac{1}{2} \right) + \log \frac{1}{2} \right)$$

$$= \underline{\underline{v_T \left(1 + \log \frac{1}{2} \right)}}$$



(i) The most probable value of X is 1

(ii) $P(X > 4) = P(\text{first 4 balls are } W)$

$$= \frac{6}{8} \times \frac{5}{7} \times \frac{4}{6} \times \frac{3}{5}$$

$$= \underline{\underline{\frac{3}{14}}}$$

Question 6

a)(i) Committees = $\frac{14C_5}{2002}$

(B) $P(\text{majority Labor})$

$$= P(3 \text{ Labor}) + P(4 \text{ Labor}) + P(5 \text{ Labor})$$

$$= \frac{6C_3 \times 8C_2 + 6C_4 \times 8C_1 + 6C_5}{14C_5}$$

$$= \underline{\underline{\frac{49}{143}}}$$

(ii) Committees

$$= 6C_3 \times 5C_1 \times 3C_1 + \frac{6}{2} (5C_2 \times 3C_1 + 5C_1 \times 3C_2)$$

$$+ 6C_1 (5C_3 \times 3C_1 + 5C_2 \times 3C_2 + 5C_1 \times 3C_3)$$

$$= 300 + 675 + 390$$


$$= \underline{\underline{1365}}$$

(B) $P(\text{majority Labor})$

$$= \frac{6C_3 \times 5C_1 \times 3C_1}{1365}$$

$$= \frac{20}{91}$$

b)



$$\Delta A(x) = 2\pi(5-x) \cdot 2\sqrt{4-x^2}$$

$$\Delta V = 4\pi(5-x)\sqrt{4-x^2} \Delta x$$

$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=-2}^{2} 4\pi(5-x)\sqrt{4-x^2} \Delta x$$

$$= 4\pi \int_{-2}^{2} (5-x)\sqrt{4-x^2} dx$$


$$= 20\pi \int_{-2}^{2} \sqrt{4-x^2} dx - 4\pi \int_{-2}^{2} x\sqrt{4-x^2} dx$$

$$= 20\pi \times \frac{1}{2}\pi(2)^2 - 0$$

$$= 20\pi \times 2\pi$$

$$= \underline{\underline{40\pi^2 \text{ units}^3}}$$

c)



$$\begin{aligned} \frac{h}{y} &= \tan 60^\circ \\ h &= y \tan 60^\circ \\ &= \sqrt{3}y \end{aligned}$$

$$\begin{aligned} \Delta A(a) &= \frac{1}{2} \times 2y \times \sqrt{3}y \\ &= \sqrt{3}y^2 \\ &= \sqrt{3}(\sqrt{9-a^2})^2 \end{aligned}$$

$$\underline{\underline{\Delta V = \sqrt{3}(\sqrt{9-a^2}) \Delta a}}$$

$$\begin{aligned} \text{(iii)} V &= \lim_{\Delta x \rightarrow 0} \sum_{x=-3}^{3} \sqrt{3}(\sqrt{9-x^2}) \Delta x \\ &= \sqrt{3} \int_{-3}^{3} \sqrt{9-x^2} dx \\ &= \sqrt{3} \times \frac{1}{2}\pi(3)^2 \\ &= \frac{9\sqrt{3}}{2}\pi \text{ units}^3 \end{aligned}$$

Question 7

$$\begin{aligned} \text{(a)(i)} P(\text{Jones}) &= \frac{0.6 \times 5 + 0.5 \times 4 + 0.4 \times 3 + 0.7 \times 8}{20} \\ &= \underline{\underline{\frac{59}{100}}} \end{aligned}$$

(ii) Let $X = \# \text{ votes for Jones}$

$$\begin{aligned} P(X \geq 8) &= P(X=8) + P(X=9) + P(X=10) \\ &= \binom{10}{8} \left(\frac{41}{100}\right)^2 \left(\frac{59}{100}\right)^8 + \binom{10}{9} \left(\frac{41}{100}\right) \left(\frac{59}{100}\right)^9 \\ &\quad + \left(\frac{59}{100}\right)^{10} \end{aligned}$$

$$\begin{aligned} &= 0.1516993349 \\ &= \underline{\underline{0.1517}} \text{ (to 4dp)} \end{aligned}$$

$$\begin{aligned} P(X \leq 2) &= P(X=0) + P(X=1) + P(X=2) \\ &= \left(\frac{41}{100}\right)^0 + \binom{10}{1} \left(\frac{41}{100}\right)^1 \left(\frac{59}{100}\right)^9 + \binom{10}{2} \left(\frac{41}{100}\right)^2 \left(\frac{59}{100}\right)^8 \\ &= 0.01457376613 \\ &= \underline{\underline{0.0146}} \text{ (to 4dp)} \end{aligned}$$

b) $e^{2x} \log y = 3$

$$(e^{2x}) \left(\frac{1}{y} \cdot \frac{dy}{dx}\right) + (\log y) (2e^{2x}) = 0$$

$$\frac{e^{2x}}{y} \frac{dy}{dx} = -2e^{2x} \log y$$

$$\underline{\underline{\frac{dy}{dx} = -2y \log y}}$$

c) $\frac{AR}{RB} = \frac{AQ}{QC} = \frac{1}{1}$ (given)

$\therefore RQ \parallel BC$ (ratio of transversals are $\frac{1}{1}$)

$PQ \parallel AB$ (by similar method)

$\therefore RPQCQ$ is \parallel gram (opposite sides are \parallel)

(ii) $\angle QRP = \angle QCP$ (opposite \angle 's in \parallel gram =)

$\angle QRP = \angle QXC$ (\angle 's in same segment)

$\therefore \angle QCP = \angle QXC$

Thus $\triangle QXC$ is isosceles ($2 = \angle$'s)

$$(iii) QY = QX \quad (\text{given})$$

$$QY = QX \quad (= \text{sides in isosceles})$$

\therefore AC is a diameter of the circumcircle of $\triangle AXC$, with Q being the centre.

$$\angle AXC = 90^\circ \quad (\angle \text{ in semicircle})$$

$$\therefore \underline{AX \perp BC}$$

Question 8

a) (i) Ways = $8!$
 $= \underline{\underline{40320}}$

(ii) No two M together

$$\begin{matrix} W & \times & \times & W \\ & \times & & \times \\ & & W & \times & W \end{matrix}$$

$$\text{Ways} = 4! \times {}^5P_4$$

$$= \underline{\underline{2880}}$$

$$(iii) P(M_A \text{ WA together})$$

$$= \frac{4! \times 2 \times {}^4P_3}{2880}$$

$$= \underline{\underline{\frac{2}{5}}}$$

b) $x^3 + ax^2 + bx + c = 0$
roots $\alpha, \beta, \alpha + \beta$

$$2\alpha + 2\beta = -a$$

$$\alpha + \beta = -\frac{a}{2}$$

$$\alpha\beta + \alpha^2 + a\beta + a\alpha + \beta^2 = b$$

$$\alpha^2 + 2\alpha\beta + \beta^2 = b$$

$$(\alpha + \beta)^2 + \alpha\beta = b$$

$$\alpha\beta(\alpha + \beta) = -c$$

$$\alpha\beta(-\frac{a}{2}) = -c$$



$$\alpha\beta = \frac{2c}{a}$$

$$(\alpha + \beta)^2 + \alpha\beta = b$$

$$\frac{a^2}{4} + \frac{2c}{a} = b$$

$$a^3 + 8c = 4ab$$

$$\underline{\underline{a^3 - 4ab + 8c = 0}}$$



$$\int_{1}^{10^{9+1}} x^{-\frac{2}{3}} dx < S < 1 + \int_{1}^{10^{9+1}} (x-1)^{-\frac{2}{3}} dx$$

$$3 \left[x^{\frac{1}{3}} \right]_{1}^{10^{9+1}} < S < 1 + 3 \left[(x-1)^{\frac{1}{3}} \right]_{1}^{10^{9+1}}$$

$$3 \left[(10^{9+1})^{\frac{1}{3}} - 1 \right] < S < 1 + 3 \left[10^3 - 1 \right]$$

$$2997.000001 < S < 2998$$

$\therefore S$ lies between 2997
and 2998.