



SYDNEY GRAMMAR SCHOOL  
MATHEMATICS DEPARTMENT  
TRIAL EXAMINATIONS 2005

# FORM VI

# MATHEMATICS EXTENSION 1

## Examination date

Wednesday 10th August 2005

## Time allowed

2 hours

## Instructions

- All seven questions may be attempted.
- All seven questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

## Collection

- Write your candidate number clearly on each booklet.
- Hand in the seven questions in a single well-ordered pile.
- Hand in a booklet for each question, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Keep the printed examination paper and bring it to your next Mathematics lesson.

## Checklist

- SGS booklets: 7 per boy. A total of 1000 booklets should be sufficient.
- Candidature: 117 boys.

## Examiner

KWM

**QUESTION ONE** (12 marks) Use a separate writing booklet.

Marks

(a) Evaluate  $\sum_{n=1}^4 n!$ . 1


(b) Differentiate the following with respect to  $x$ .  
 (i)  $y = \log_e(\sin x)$  1

(ii)  $y = \cos^{-1} 3x$  1

(c) State the domain and range of the function  $f(x) = 2 \cos^{-1} \frac{x}{3}$ . 2

(d) Given the points  $A(5, 1)$  and  $B(-3, 6)$ , find the co-ordinates of the point  $P$  that divides the interval  $AB$  externally in the ratio 3 : 4. 2

(e) 2



The diagram above shows the tangents  $AC$  and  $AE$  drawn to a circle.  $BF$  and  $DF$  are chords drawn from the points of contact at  $B$  and  $D$  respectively. Given that  $\angle BFD = 66^\circ$ , find  $\angle BAD$  giving reasons for your answer.

(f) Use the substitution  $u = 1 - x^2$  to evaluate the definite integral 3

$$\int_0^{\frac{\sqrt{3}}{2}} x\sqrt{1-x^2} dx.$$

**QUESTION TWO** (12 marks) Use a separate writing booklet.

**Marks**

(a) Simplify  $\frac{{}^nC_{r+1}}{{}^nC_r}$ . 2

(b) Find the term independent of  $x$  in the expansion of  $\left(3x^2 + \frac{2}{x}\right)^{12}$ . 3

(c) A couple, purchasing a house, negotiates a \$300 000 mortgage to be repaid in equal monthly instalments over a period of 25 years. The interest on the loan is 7.2% per annum, compounded monthly. Let  $\$A_n$  be the amount owing on the loan after  $n$  months, and  $\$M$  the monthly repayment.

(i) Write down an expression for  $A_1$ . 1

(ii) Hence show that  $A_2 = 300\,000(1.006)^2 - 1.006M - M$ . 1

(iii) Show that  $A_n = 300\,000(1.006)^n - \frac{M(1.006^n - 1)}{0.006}$ . 1

(iv) Find, to the nearest dollar, the monthly repayment  $M$  required to repay the loan over 25 years under the agreed terms. 1

(d) Find  $\int_0^{\frac{\pi}{3}} \tan^2 x \, dx$ . 3


**QUESTION THREE** (12 marks) Use a separate writing booklet.

Marks

(a) Prove that  $\frac{1 - \cos 2A}{\sin 2A} = \tan A$ .

2

(b)



In the diagram above the curve  $y = \frac{1}{\sqrt{4-x^2}}$  is sketched showing vertical asymptotes at  $x = -2$  and  $x = 2$ . Find the exact area of the shaded region bounded by the curve, the line  $x = 1$  and the co-ordinate axes.

2

(c) Let the equation  $x^3 - 3x^2 - 4x + 12 = 0$  have roots  $\alpha$ ,  $\beta$  and  $\gamma$ .

(i) Find the value of  $\alpha + \beta + \gamma$ .

1


(ii) Find the value of  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$ .

2

(iii) Given that two of its roots sum to zero, solve the equation.

2

(d)



The diagram above shows the curve  $y = e^x$  and the line  $y = 2x + 1$  intersecting at point  $P(0, 1)$  and at another point  $Q$ . Use Newton's Method once, with initial approximation  $x = 1$ , to find a better approximation to the  $x$  co-ordinate of the point  $Q$ . Write your approximation correct to one decimal place.

3

**QUESTION FOUR** (12 marks) Use a separate writing booklet.

Marks

(a) Find  $\cos^{-1}(-\frac{1}{2}) + \sin^{-1}(-\frac{\sqrt{3}}{2})$  in radians. 1

(b) Find the values of  $a$  and  $b$  that make the polynomial  $P(x) = 2x^3 + ax^2 - 13x + b$  exactly divisible by  $x^2 - x - 6$ . 3

(c) (i) Express  $\cos x - \sqrt{3}\sin x$  in the form  $R \cos(x + \alpha)$ , where  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ . 2

(ii) Hence, or otherwise, find the general solution of the equation 2

$$\cos x - \sqrt{3}\sin x = 1.$$

(d) If the surrounding air temperature is  $20^\circ\text{C}$ , it takes 15 minutes for a cup of tea at a temperature of  $80^\circ\text{C}$  to cool to a temperature of  $40^\circ\text{C}$ . Given that  $T$  is the temperature in degrees Celsius of the tea after  $t$  minutes, then Newton's Law of cooling states that  $T$  satisfies the differential equation  $\frac{dT}{dt} = k(T - 20)$ .

(i) Show that  $T = 20 + Ae^{kt}$  is a solution of the differential equation. 1


(ii) Find the value of  $A$ , and show that  $k = -\frac{\ln 3}{15}$ . 2

(iii) Find the temperature of the tea after 30 minutes. 1

**QUESTION FIVE** (12 marks) Use a separate writing booklet.

Marks

(a)



In the diagram above a focal chord  $PQ$  intersects the parabola  $x^2 = 4ay$  at points  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$ . The tangents to the parabola at point  $P$  and point  $Q$  intersect at  $T$ .


- (i) Show that the equation of the tangent to the parabola at the point  $P$  is given by  $y = px - ap^2$ . 2
- (ii) Show that  $pq = -1$ . 2
- (iii) Show that the acute angle between the focal chord  $QP$  and the tangent  $TP$  to the parabola at  $P$  is given by  $\tan^{-1} |q|$ . 2

(b) A particle is moving in simple harmonic motion about the origin.

- (i) Assuming that  $\ddot{x} = -n^2x$ , show that  $v^2 = n^2(a^2 - x^2)$ , where  $a$  is the amplitude. 2
- (ii) When the particle is 3 metres from the origin, its speed is 8 m/s, and when it is 4 metres from the origin its speed is 6 m/s. Find the period and amplitude of the motion. 3
- (iii) Find the greatest acceleration of the particle. 1

**QUESTION SIX** (12 marks) Use a separate writing booklet.

Marks




The diagram above shows a plane inclined at  $30^\circ$  to the horizontal, meeting level ground at  $B$ . A ball is projected from a point  $A$  on the plane, 10 metres above the horizontal. The angle of projection is  $60^\circ$  to the horizontal and the initial speed of the ball is 40 m/s.

- (i) Take  $g = 10 \text{ m/s}^2$ , and show that the displacement equations of motion of the ball are given by 3

$$y = 20\sqrt{3}t - 5t^2 + 10 \quad \text{and}$$

$$x = 20t.$$

- (ii) Show that the ball hits the inclined plane at the point  $P$  after  $t = \frac{16\sqrt{3}}{3}$  seconds. 3



In the diagram above,  $TA$  is a tangent and  $TBC$  is a secant drawn to a circle of centre  $O$ . Let the midpoint of the chord  $BC$  be  $D$ . Prove that  $\angle AOT = \angle ADT$ . 3


Exam continues overleaf ...

- (c) A rectangle is expanding in such a way that at all times it is twice as long as it is wide. If its area is increasing at a rate of  $18 \text{ cm}^2/\text{s}$ , find the rate at which its perimeter is increasing at the instant its width is 1 metre. 3

**QUESTION SEVEN** (12 marks) Use a separate writing booklet.

Marks

(a)



The diagram above shows two vertical towers  $BD$  and  $CE$  of heights  $h$  and  $2h$  respectively, on a horizontal plane  $ABC$ . Point  $A$  is due south of point  $B$ , and the angles of elevation of the tops of the towers from  $A$  are both  $30^\circ$ . Given that the angle of elevation from  $D$  to  $E$  is  $15^\circ$ , find the bearing of the taller tower from point  $A$  correct to the nearest degree. 4

- (b) By considering the expansion of  $(1 + x)^{n-1}$ , prove that: 4

$$\frac{7}{1} \binom{n-1}{0} + \frac{7^2}{2} \binom{n-1}{1} + \frac{7^3}{3} \binom{n-1}{2} + \dots + \frac{7^n}{n} \binom{n-1}{n-1} = \frac{1}{n} (2^{3n} - 1).$$

- (c) Use induction, or otherwise, to prove that the sum of the products of all the pairs of different integers that can be formed from the first  $n$  positive integers is 4

$$\frac{n}{24} (n-1)(n+1)(3n+2).$$

**END OF EXAMINATION**



The following list of standard integrals may be used:

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE:  $\ln x = \log_e x, \quad x > 0$

QUESTION 1

(a)  $\sum_{n=1}^4 n! = 1! + 2! + 3! + 4!$   
 $= 1 + 2 + 6 + 24$   
 $= 33 \checkmark$

(b) (i)  $y = \ln(\sin x)$   
 $y' = \frac{1}{\sin x} \times \cos x$   
 $y' = \cot x \checkmark$

(ii)  $y = \cos^{-1} 3x$   
 $y' = \frac{1}{\sqrt{1-9x^2}} \times 3$   
 $y' = \frac{3}{\sqrt{1-9x^2}} \checkmark$

(c)  $f(x) = 2 \cos^{-1} \frac{x}{3}$

Domain:  $-1 \leq \frac{x}{3} \leq 1$   
 so  $-3 \leq x \leq 3 \checkmark$

Range:  $0 \leq y \leq 2\pi \checkmark$

(d) A(5, 1), B(-3, 6)  
 $K: P = -3:4$

$P \left( \frac{1x_1 + Kx_2}{1+K}, \frac{1y_1 + Ky_2}{1+K} \right) \checkmark$

$P \left( \frac{20+9}{1}, \frac{4-18}{1} \right)$

$P(29, -14) \checkmark$

(e)  $\angle DBA = 66^\circ$  (The angle between a chord and tangent at the point of contact equals the angle drawn in the alternate segment.)  $\checkmark$

Tangents drawn from an external point are equal, so  $AB = AD$  and  $\triangle ABD$  is isosceles.

$\angle BAD = \angle DBA = 66^\circ$

(base angles of an isosceles triangle are equal.)

so  $\angle BAD = 180^\circ - 2 \times 66^\circ$   
 $= 48^\circ \checkmark$

(angle sum of a triangle.)

(f)  $\int_0^{\frac{\sqrt{3}}{2}} x\sqrt{1-x^2} dx$   $u = 1-x^2$   
 $du = -2x dx$   
 $-\frac{1}{2} du = x dx$

when  $x=0$ ,  $u=1$   $\checkmark$

when  $x = \frac{\sqrt{3}}{2}$ ,  $u = \frac{1}{4}$  :

$-\frac{1}{2} \int_1^{\frac{1}{4}} u^{\frac{1}{2}} du = \frac{1}{3} \left[ u^{\frac{3}{2}} \right]_{\frac{1}{4}}^1$

$= \frac{1}{3} \left( 1 - \frac{1}{8} \right)$

$= \frac{1}{3} \times \frac{7}{8}$

$= \frac{7}{24} \checkmark$

QUESTION 2

$$(a) \frac{{}^n C_{r+1}}{{}^n C_r} = \frac{n!}{(n-r-1)!(r+1)!} \times \frac{(n-r)! r!}{n!}$$

$$= \frac{n-r}{r+1} \checkmark$$

$$(b) \left(3x^2 + \frac{2}{x}\right)^{12}$$

$$\text{General term } T_r = {}^{12} C_r (3x^2)^{12-r} \left(\frac{2}{x}\right)^r$$

using rules of indices:

$$24 - 3r = 0$$

$$3r = 24$$

$$r = 8 \checkmark$$

$${}^{12} C_8 \times 3^4 \times 2^8 \checkmark$$

(e)

$$(i) A_1 = 300000 \times 1.006 - M \checkmark$$

$$(ii) A_2 = A_1 \times 1.006 - M$$

$$= 300000 (1.006)^2 - M (1.006) - M \checkmark$$

$$(iii) A_n = 300000 (1.006)^n$$

$$- M (1 + 1.006 + 1.006^2 + \dots + 1.006^{n-1})$$

a geometric progression

$$a = 1 \text{ and } r = 1.006.$$

$$S_n = \frac{a(r^n - 1)}{r - 1} \checkmark$$

$$S_n = \frac{(1.006)^n - 1}{0.006}$$

$$A_n = 300000 (1.006)^n - \frac{M (1.006^n - 1)}{0.006}$$

(iv) after the loan is repaid

$$A_n = 0, \text{ and } n = 300$$

$$M = \frac{300000 (1.006)^{300} (0.006)}{(1.006)^{300} - 1}$$

$$M \doteq \$2159 \checkmark$$

(d)

$$\int_0^{\frac{\pi}{3}} \tan^2 x \, dx$$

$$= \int_0^{\frac{\pi}{3}} \sec^2 x - 1 \, dx \checkmark$$

$$= \left[ \tan x - x \right]_0^{\frac{\pi}{3}} \checkmark$$

$$= \left( \sqrt{3} - \frac{\pi}{3} \right) - (0)$$

$$= \sqrt{3} - \frac{\pi}{3} \checkmark$$

(12)

QUESTION 3

$$\begin{aligned}
 \text{(a) LHS} &= \frac{1 - \cos 2A}{\sin 2A} \\
 &= \frac{1 - (1 - 2\sin^2 A)}{2\sin A \cos A} \\
 &= \frac{2\sin^2 A}{2\sin A \cos A} \\
 &= \frac{\sin A}{\cos A} \\
 &= \tan A \\
 &= \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{(b) Area} &= \int_0^1 \frac{dx}{\sqrt{4-x^2}} \\
 &= \left[ \sin^{-1} \frac{x}{2} \right]_0^1 \\
 &= \frac{\pi}{6} - 0 \\
 &= \frac{\pi}{6} \text{ sq. units.}
 \end{aligned}$$

$$\text{(c) } x^3 - 3x^2 - 4x + 12 = 0$$

$$\begin{aligned}
 \text{(i) } \alpha + \beta + \gamma &= -\frac{b}{a} \\
 &= 3
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} &= \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma} \\
 &= \frac{c}{a} \\
 &= -\frac{d}{a} \\
 &= -\frac{c}{d} \\
 &= \frac{1}{3}
 \end{aligned}$$

(iii) Let the roots be  $\alpha, -\alpha, \beta$ .

$$\text{then } \alpha - \alpha + \beta = 3$$

$$\text{and } \beta = 3. \checkmark$$

$$\text{now } \alpha(-\alpha)\beta = -12$$

$$-\alpha^2 = -4$$

$$\alpha = \pm 2$$

the roots are  $-2, 2$  and  $3. \checkmark$

$$\begin{aligned}
 \text{(d) } y &= e^x \\
 y &= 2x + 1
 \end{aligned}$$

The points of intersection correspond to the roots of the equation

$$e^x - 2x - 1 = 0$$

$$f(x) = e^x - 2x - 1. \checkmark$$

$$f'(x) = e^x - 2$$

$$\text{put } x_0 = 1 : f(1) = -0.282$$

$$f'(1) = 0.718$$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} \checkmark$$

$$= 1 + \frac{0.282}{0.718}$$

$$0.718$$

$$\approx 1.4 \text{ (1 dec. place.)} \checkmark$$

(12)

QUESTION 4.

(a)  $\cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{\sqrt{3}}{2}\right)$

$= \frac{2\pi}{3} - \frac{\pi}{3}$   
 $= \frac{\pi}{3}$  ✓

(b)

$P(x) = 2x^3 + ax^2 - 13x + b$

$x^2 - x - 6 = (x-3)(x+2)$

$(x+2)$  is a factor:  $P(-2) = 0$

$(x-3)$  is a factor:  $P(3) = 0$ .

$P(-2): -16 + 4a + 26 + b = 0$  ①

$P(3): 54 + 9a - 39 + b = 0$  ②

$4a + b = -10$  — ①

$9a + b = -15$  — ②

② - ①:  $5a = -5$

$a = -1$  ✓

$b = -6$  ✓

(c) (i)

Let  $\cos x - \sqrt{3} \sin x \equiv R \cos(x+\alpha)$

$\cos x - \sqrt{3} \sin x \equiv R \cos x \cos \alpha - R \sin x \sin \alpha$

equating co-efficients:

$\cos x: R \cos \alpha = 1$  — ①

$\sin x: R \sin \alpha = \sqrt{3}$  — ②

② / ①:  $\tan \alpha = \sqrt{3}$

①:  $\alpha = \frac{\pi}{3}$  ✓

①<sup>2</sup> + ②<sup>2</sup>:  $R^2 = 4$

$R = 2$  ✓

$\cos x - \sqrt{3} \sin x = 2 \cos\left(x + \frac{\pi}{3}\right)$

(ii)  $\cos x - \sqrt{3} \sin x = 1$

$2 \cos\left(x + \frac{\pi}{3}\right) = 1$

$\cos\left(x + \frac{\pi}{3}\right) = \frac{1}{2}$

$x + \frac{\pi}{3} = 2\pi n + \frac{\pi}{3}$  or  $2\pi n - \frac{\pi}{3}$

$x = 2\pi n$  or  $x = 2\pi n - \frac{2\pi}{3}$  ✓

(d) (i)  $T = 20 + Ae^{Kt}$

$\frac{dT}{dt} = KAe^{Kt}$

$\frac{dT}{dt} = K(T-20)$

It satisfies the DE. ✓

(ii) when  $t=0, T=80^\circ\text{C}$

$80 = 20 + A$

$A = 60$  ✓

$T = 20 + 60e^{Kt}$

when  $t=15, T=40^\circ\text{C}$

$40 = 20 + 60e^{15K}$

$e^{15K} = \frac{1}{3}$

$15K = \ln \frac{1}{3}$  ✓

$15K = -\ln 3$

$K = -\frac{1}{15} \ln 3$  as required.

(iii)  $T = 20 + 60e^{-\frac{20 \ln 3}{15} t}$

$= 20 + 60e^{-\frac{2 \ln 3}{3} t}$

$= 20 + 60e^{\ln \frac{1}{9}}$

$= 20 + 60 \times \frac{1}{9}$

$= 26\frac{2}{3}^\circ\text{C}$  ✓

(12)

QUESTION 5

(i)  $x = 2ap$   $y = ap^2$   
 $\frac{dx}{dp} = 2a$   $\frac{dy}{dp} = 2ap$

$\frac{dy}{dx} = \frac{dy}{dp} \times \frac{dp}{dx}$

$= 2ap \times \frac{1}{2a}$   
 $= p$  ✓

gradient at  $P = p$ ,  $(2ap, ap^2)$

$y - y_1 = m(x - x_1)$

$y - ap^2 = p(x - 2ap)$

$y - ap^2 = px - 2ap^2$

$y = px - ap^2$  is ✓

the equation of the tangent at P.

(ii) Gradient of PS = Gradient of QP

$\frac{ap^2 - a}{2ap} = \frac{ap^2 - aq^2}{2ap - 2aq}$  ✓

$\frac{p^2 - 1}{2p} = \frac{a(p - q)(p + q)}{2a(p - q)}$

$p^2 - 1 = p(p + q)$

$p^2 - 1 = p^2 + pq$  ✓

∴  $pq = -1$  as required.

(iii)

Let  $m_1$  be the gradient of PQ and  $m_2$  be the gradient of PT.

$m_1 = \frac{p+q}{2}$   $m_2 = p$

$\tan \angle TPQ = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

$= \left| \frac{\frac{p+q}{2} - p}{1 + p \frac{p+q}{2}} \right|$  ✓

$= \left| \frac{p+q - 2p}{2 + p^2 + pq} \right|$  ( $pq = -1$ )

$= \left| \frac{q-p}{p^2+1} \right|$  ( $p = -\frac{1}{q}$ )

$= \left| \frac{q + \frac{1}{q}}{\frac{1}{q^2} + 1} \right|$  ✓

$= \left| \frac{q^3 + q}{1 + q^2} \right|$

$= \left| \frac{q(q^2+1)}{(q^2+1)} \right|$

$= |q|$

(b) (i)

$\ddot{x} = -n^2 x$

$\frac{d}{dx} \left( \frac{1}{2} v^2 \right) = -n^2 x$  ✓

$\frac{1}{2} v^2 = -\frac{n^2 x^2}{2} + c$

when  $x = a$ ,  $v = 0$  and  $c = \frac{n^2 a^2}{2}$

so  $v^2 = -n^2 x^2 + n^2 a^2$  ✓

$v = n^2 (a^2 - x^2)$

(ii)  $v^2 = n^2 (a^2 - x^2)$

when  $x = 3$ ,  $v = 8$

$64 = n^2 (a^2 - 9)$  — (1)

when  $x = 4$ ,  $v = 6$  ✓

$36 = n^2 (a^2 - 16)$  — (2)

(1) :  $\frac{a^2 - 9}{a^2 - 16} = \frac{64}{36}$

Q5 continued.

$$9(a^2 - 9) = 16(a^2 - 16)$$

$$9a^2 - 81 = 16a^2 - 256$$

$$7a^2 = 175$$

$$a^2 = 25$$

$$a = 5 \text{ (amplitude } > 0)$$

$$\textcircled{2} \quad 6 = n\sqrt{25-16}$$

$$3n = 6$$

$$n = 2.$$

$$T = \frac{2\pi}{n}$$

$$= \frac{2\pi}{2}$$

$$\underline{T = \pi \text{ s}}$$

(ii) The maximum acceleration occurs when  $x = a$ .

$$\ddot{x} = -n^2 x$$

$$\ddot{x} = -4 \times 5$$

(12)


maximum acceleration is

$$20 \text{ m/s}^2$$

(negative implies direction.)

QUESTION 6.

(a)(i)



when  $t=0$ .

$$\begin{aligned} \dot{x} &= 40 \cos 60^\circ & \dot{y} &= 40 \sin 60^\circ \\ \dot{x} &= 20 \text{ m/s} & \dot{y} &= 20\sqrt{3} \text{ m/s} \end{aligned}$$

$$\begin{aligned} \ddot{y} &= -10 & \ddot{x} &= 0 \\ \text{integrate w.r.t } t. & & & \end{aligned}$$

$$y = -10t + c_1 \quad x = c_2$$

when  $t=0$ ,  $y = 20\sqrt{3}$  and  $x = 20$   
thus  $c_1 = 20\sqrt{3}$  and  $c_2 = 20$ .

$$y = 20\sqrt{3} - 10t \quad x = 20 \quad \checkmark$$


integrate w.r.t  $t$ .

$$y = 20t\sqrt{3} - 5t^2 + c_3 \quad x = 20t + c_4$$

when  $t=0$ ,  $y = 10$ ,  $x = 0$   
thus  $c_3 = 10$  and  $c_4 = 0$

The equations of motion  $\checkmark$   
are  $y = 20t\sqrt{3} - 5t^2 + 10$   
 $x = 20t$

(ii) When the ball hits the plane at P:



$$\frac{10-y}{x} = \tan 30^\circ \quad \checkmark$$

$$\frac{5t^2 - 20t\sqrt{3}}{20t} = \frac{1}{\sqrt{3}} \quad \checkmark$$

$$\frac{t - 4\sqrt{3}}{4} = \frac{1}{\sqrt{3}}$$


$$t - 4\sqrt{3} = \frac{4}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$$

$$t = 4\sqrt{3} + \frac{4\sqrt{3}}{3}$$

$$t = \frac{16\sqrt{3}}{3} \text{ s. } \checkmark$$

(b)  $\angle TAO = 90^\circ$  (TA is a tangent)  
 $\angle TDO = 90^\circ$  (a line drawn from the centre to the mid-point of the chord is perpendicular to the chord.)

TAOD is a cyclic quadrilateral since opposite angles are supplementary. Hence  $\angle AOT = \angle ADT$  (angles standing on the same arc - drawn to the circumference are equal.)  $\checkmark$



$$A = 2x^2$$

$$\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt} \quad \text{by chain rule}$$

$$\frac{dA}{dt} = 4x \times \frac{dx}{dt}$$

at  $x = 100 \text{ cm}$ ,  $18 = 400 \times \frac{dx}{dt}$

$$\frac{dx}{dt} = \frac{9}{200} \quad \checkmark$$

Now  $P = 6x \quad \checkmark$

$$\frac{dP}{dt} = 6 \times \frac{dx}{dt}$$

$$= 6 \times \frac{9}{200}$$

$$= \frac{27}{100} \text{ cm/s. } \checkmark$$

(12)



QUESTION 7


(a)  $\triangle DEF: \tan 15^\circ = \frac{h}{DF}$

$DF = h \cot 15^\circ$

and  $BC = h \cot 15^\circ$ . ✓

$\triangle ABD: AB = h \cot 30^\circ$

$\triangle AEC: AC = 2h \cot 30^\circ$  ✓



$\triangle ABC$ , using the cosine rule:

$$\begin{aligned} \cos \theta &= \frac{h^2 \cot^2 30^\circ + 4h^2 \cot^2 30^\circ - h^2 \cot^2 15^\circ}{4h^2 \cot^2 30^\circ} \checkmark \\ &= \frac{5 \cot^2 30^\circ - \cot^2 15^\circ}{4 \cot^2 30^\circ} \\ &= \frac{15 - \cot^2 15^\circ}{12} \end{aligned}$$

$\cos \theta = 0.089316$

$\theta = 84^\circ 53'$

The bearing of the taller tower from A is  $N 84^\circ 53' E$ . ✓

QUESTION 7 Continued.

$$(b) (1+x)^{n-1} = \binom{n-1}{0} + \binom{n-1}{1}x + \binom{n-1}{2}x^2 + \dots + \binom{n-1}{n-1}x^{n-1} \quad \checkmark$$

integrating both sides w.r.t  $x$  :

$$\frac{(1+x)^n}{n} = \binom{n-1}{0}x + \frac{1}{2}\binom{n-1}{1}x^2 + \frac{1}{3}\binom{n-1}{2}x^3 + \dots + \frac{1}{n}\binom{n-1}{n-1}x^n + C_1$$

put  $x=0$ .

$$\frac{1}{n} = C_1$$

$$\frac{(1+x)^n}{n} = \binom{n-1}{0}x + \frac{1}{2}\binom{n-1}{1}x^2 + \frac{1}{3}\binom{n-1}{2}x^3 + \dots + \frac{1}{n}\binom{n-1}{n-1}x^n + \frac{1}{n} \quad \checkmark$$

put  $x=7$ .

$$\frac{8^n}{n} = 7\binom{n-1}{0} + \frac{7^2}{2}\binom{n-1}{1} + \frac{7^3}{3}\binom{n-1}{2} + \dots + \frac{7^n}{n}\binom{n-1}{n-1} + \frac{1}{n} \quad \checkmark$$

$$\frac{1}{n}(8^n - 1) = 7\binom{n-1}{0} + \frac{7^2}{2}\binom{n-1}{1} + \frac{7^3}{3}\binom{n-1}{2} + \dots + \frac{7^n}{n}\binom{n-1}{n-1}$$

$$\therefore 7\binom{n-1}{0} + \frac{7^2}{2}\binom{n-1}{1} + \frac{7^3}{3}\binom{n-1}{2} + \dots + \frac{7^n}{n}\binom{n-1}{n-1} = \frac{1}{n}(2^{3n} - 1) \quad \checkmark$$

(c) Show the statement is true for  $n=2$ .

A 1, 2. Sum of the products =  $1 \times 2 = 2$ .

$$\begin{aligned} \frac{n}{24}(n-1)(n+1)(3n+2) &= \frac{2}{24}(2-1)(2+1)(6+2) \\ &= \frac{1}{24} \times 1 \times 3 \times 8 \\ &= 2 \quad \checkmark \end{aligned}$$

Its true for  $n=2$ .

B Assume that the result is true for  $n=k$ .

ie/ the sum of the products of all the pairs of integers that can be formed from the first  $k$  positive integers is  $\frac{k}{24}(k-1)(k+1)(3k+2)$ .

Prove that it is true for  $n=k+1$ .

The addition of the integer  $(k+1)$  will add another  $(1+2+3+\dots+k)(k+1)$  products in pairs.

Q7 (c) continued.

10.

When  $n = k+1$  the sum is :

$$\frac{k}{24} (k-1)(k+1)(3k+2) + (k+1)(1+2+3+\dots+k) \checkmark$$

$(1+2+3+\dots+k)$  is an arithmetic series.  $\text{Sum} = \frac{k}{2}(1+k)$

$$= \frac{k}{24} (k-1)(k+1)(3k+2) + (k+1) \frac{k}{2} (k+1) \checkmark$$

$$= \frac{k}{24} (k+1) \{ (k-1)(3k+2) + 12(k+1) \}$$

$$= \frac{k}{24} (k+1) \{ 3k^2 + 11k + 10 \}$$

$$= \frac{k}{24} (k+1)(k+2)(3k+5) \text{ as required. } \checkmark$$

c// It follows from parts A and B by mathematical induction that the statement is true for all positive integers  $n \geq 2$ .

12

