

Using Symmetry

(1) Even

$$f(-x) = f(x)$$

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

NOTE: horizontal shift

$$\int_{c-a}^{c+a} f(x-c) dx = 2 \int_c^{c+a} f(x-c) dx$$

(2) Odd

$$f(-x) = -f(x)$$

$$\int_{-a}^a f(x) dx = 0$$

NOTE: horizontal shift

$$\int_{c-a}^{c+a} f(x-c) dx = 0$$

(3)

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

Proof:

$$\begin{aligned} & \int_0^a f(a-x)dx && u = a-x \\ &= -\int_0^a f(u)du && du = -dx \\ &= \int_0^a f(u)du && x=0, u=a \\ &= \int_0^a f(x)dx && x=a, u=0 \end{aligned}$$

odd \times odd = even

odd \times even = odd

even \times even = even

e.g. (i) $\int_{-1}^1 \sin^3 x dx = 0$ (odd function)³ = odd function

$$\begin{aligned}(ii) \int_0^1 x^2 \sqrt{1-x} dx &= \int_0^1 (1-x)^2 \sqrt{x} dx \\&= \int_0^1 \left(x^{\frac{1}{2}} - 2x^{\frac{3}{2}} + x^{\frac{5}{2}} \right) dx \\&= \left[\frac{2}{3}x^{\frac{3}{2}} - \frac{4}{5}x^{\frac{5}{2}} + \frac{2}{7}x^{\frac{7}{2}} \right]_0^1 \\&= \frac{2}{3} - \frac{4}{5} + \frac{2}{7} - 0 \\&= \underline{\underline{\frac{16}{105}}}\end{aligned}$$

Improper Integrals

An **improper** integral is a definite integral where the integrand is undefined at some point in the interval or unbounded.

We must use limits to solve, if a solution exists.

$$\begin{aligned}\text{e.g. } (i) \int_0^1 \frac{dx}{x} &= \lim_{N \rightarrow 0} \int_N^1 \frac{dx}{x} \\&= \lim_{N \rightarrow 0} [\log x]_N^1 \\&= \lim_{N \rightarrow 0} (-\log N) \\&\underline{\text{Undefined}}\end{aligned}$$

\therefore integral is undefined


$$\begin{aligned}(ii) \int_1^\infty xe^{x^2} dx &= \lim_{N \rightarrow \infty} \int_1^N xe^{-x^2} dx \\&= \lim_{N \rightarrow \infty} -\frac{1}{2} [e^{-x^2}]_1^N \\&= \lim_{N \rightarrow \infty} -\frac{1}{2} (e^{-N^2} - e^{-1}) \\&= \underline{\frac{1}{2e}}$$

Area and Inequalities

If $g(x) \leq f(x) \leq h(x)$ for $a \leq x \leq b$

$$\text{then } \int_a^b g(x)dx \leq \int_a^b f(x)dx \leq \int_a^b h(x)dx$$

e.g. (2009 Question 8 b)



Let n be a positive integer greater than 1.

The area of the region under the curve $y = \frac{1}{x}$ from $x = n - 1$ to $x = n$ is between the area of two rectangles, as shown in the diagram.

$$\text{Show that } e^{-\frac{n}{n-1}} < \left(1 - \frac{1}{n}\right)^n < e^{-1}$$

Area of inner rectangle < Area under the curve < Area of outer rectangle

$$(1) \left(\frac{1}{n} \right) < \int_{n-1}^n \frac{dx}{x} < \left(1 \right) \left(\frac{1}{n-1} \right)$$

$$\frac{1}{n} < [\log x]_{n-1}^n < \frac{1}{n-1}$$

$$\frac{1}{n} < \log \left(\frac{n}{n-1} \right) < \frac{1}{n-1}$$

$$1 < n \log \left(\frac{n}{n-1} \right) < \frac{n}{n-1}$$

$$1 < \log \left(\frac{n}{n-1} \right)^n < \frac{n}{n-1}$$

as $y = e^x$ is continually increasing

$$e < \left(\frac{n}{n-1} \right)^n < e^{\frac{n}{n-1}}$$

$$\frac{1}{e} > \left(\frac{n-1}{n} \right)^n > \frac{1}{e^{\frac{n}{n-1}}}$$

i.e. $\underline{e^{-\frac{n}{n-1}} < \left(1 - \frac{1}{n} \right)^n < e^{-1}}$

Patel Exercise 2I; 1 bdf, 2 ace, 3
Exercise 2J; 42, 44

Cambridge Exercise 2I; 1, 2ac, 3, 6, 8, 9c, 10, 11ad, 12c, 13, 18

Note: $(2a - x)$ instead of $(a - x)$

Exercise 2H; 1, 2defgh, 3b to h, 4, 6, 7b, 8, 12, 15

The 100 (*not* 78)