



Sydney Girls High School 2013

TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Extension 1 Mathematics

General Instructions

- Reading Time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
Black pen is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11 – 14, show relevant mathematical reasoning and/or calculations

Total marks – 70

Section I Pages 3 – 6

10 Marks

- Attempt Questions 1 – 10
- Answer on the Multiple Choice answer sheet provided
- Allow about 15 minutes for this section

Section II Pages 7 – 13

60 Marks

- Attempt Questions 11 – 14
- Answer on the blank paper provided
- Begin a new page for each question
- Allow about 1 hour and 45 minutes for this section

Name:

Teacher:

THIS IS A TRIAL PAPER ONLY

It does not necessarily reflect the format or the content of the 2013 HSC Examination Paper in this subject.

BLANK PAGE

Section I

10 marks

Attempt Questions 1 – 10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1 – 10

(1) The acute angle between the straight lines $y = \sqrt{3}x + 2$ and $y = 2$ is :

(A) 30°

(B) 60°

(C) 47°

(D) 68°

(2) The value of $\lim_{n \rightarrow \infty} \frac{5(10^n) + 3}{2(10^n) + 5}$ is:

(A) $\frac{3}{5}$

(B) 0

(C) 1

(D) $\frac{5}{2}$

(3) The exact value of k given $\int_0^1 \frac{dx}{x^2 + 3} = k\pi$ is:

(A) $\sqrt{3}$

(B) $\frac{\sqrt{3}}{9}$

(C) $\frac{\sqrt{3}}{18}$

(D) $6\sqrt{3}$

(4) Which of the following is the derivative of $x^2 \cos^{-1} 3x$?

(A) $2x \sin^{-1} 3x$

(B) $2x \cos^{-1} 3x + x^2 \sin^{-1} 3x$

(C) $2x \cos^{-1} 3x - \frac{x^2}{\sqrt{1-9x^2}}$

(D) $2x \cos^{-1} 3x - \frac{3x^2}{\sqrt{1-9x^2}}$

(5) The solution to $\ln(x^3 + 19) = 3 \ln(x + 1)$ is:

(A) $x = -3$ or $x = 2$

(B) $x = 3$

(C) $x = -2$

(D) $x = 2$

(6) The exact value of $\int_0^{\frac{\pi}{4}} \cos^2 \frac{1}{2}x \, dx$ is :

(A) $\frac{1+\pi}{\sqrt{2}}$

(B) $\frac{2\sqrt{2}+\pi}{8}$

(C) $\frac{2\sqrt{2}+\pi}{4}$

(D) $\frac{\sqrt{2}+\pi}{8}$

(7) The domain of $y = \cos^{-1} \sqrt{\frac{1}{4} - x^2}$ is :

(A) $0 \leq x \leq \frac{1}{2}$

(B) $-\frac{1}{4} \leq x \leq \frac{1}{2}$

(C) $-\frac{1}{2} \leq x \leq \frac{1}{2}$

(D) $\frac{1}{4} \leq x \leq \frac{1}{2}$

(8) A metal disc , 5 cm radius , expands when heated. If the radius is increasing at the rate of

0.01 cm / sec , the rate at which the area of one of the faces is increasing is given by:

(A) $\frac{\pi}{10} cm^2 / sec$

(B) $\frac{\pi}{5} cm^2 / sec$

(C) $\frac{2\pi}{5} cm^2 / sec$

(D) $\frac{5\pi}{2} cm^2 / sec$

(9) Two roots of the equation $x^3 - 2x^2 + kx + 18 = 0$ are opposites. The value of k is :

(A) - 9

(B) 9

(C) - 6

(D) 6

(10) A point moving with simple harmonic motion starts from a point 5cm from the centre of the

motion with a speed of 1cm / s . The period is 8 seconds. The maximum acceleration is:

(A) $4.9ms^{-2}$

(B) $5.2ms^{-2}$

(C) $24.4ms^{-2}$

(D) $25.6ms^{-2}$

Section II

60 marks

Attempt Questions 11 – 14

Allow about 1 hours and 45 minutes for this section

Answer on the blank paper provided. Begin a new page for each question

Your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (Begin a New Page)

(15 Marks)

(a) By making the substitution $u^2 = x + 1$, find $\int \frac{x+2}{\sqrt{x+1}} dx$ [2]

(b) Solve: $x + 2 < \frac{4}{x-1}$ ($x \neq 1$) [3]

(c) Find the general solution (in radian form) of the equation $\cos 2x = \cos x$ [3]

(d) i) Sketch the graph of the curve $y = 3 \sin^{-1}(x/2)$, clearly indicating the domain and range. [1]

ii) Find the area enclosed between the curve $y = 3 \sin^{-1}(x/2)$, the line $x=1$ and the positive x axis. [3]

(e) Consider the series $\tan x + \tan^3 x + \tan^5 x + \dots$, where $0 \leq x \leq \frac{\pi}{4}$

i) Explain why this series has a limiting sum [1]

ii) Show that $S_{\infty} = \frac{1}{2} \tan 2x$ [2]

End of Question 11

Question 12 (Begin a New Page)

(15 Marks)

(a) Use mathematical induction to show that $5^n + 2(11^n)$ is a multiple of 3 for all positive integers n . [3]

(b) At time t minutes the number of individuals in each of population

A and B is given by $N_A = 15 + 20e^{-0.5t}$ and $N_B = 5 + 40e^{-0.5t}$ respectively.

i) Find the initial size of population A [1]

ii) Find the initial rate of change of population B [1]

iii) Find the time at which the two population sizes are equal. [2]

(c) A particle moves along the x axis according to the equation

$$x = 6 \sin 2t - 2\sqrt{3} \cos 2t .$$

i) Express x in the form $R \sin(2t - \alpha)$ where $R > 0$ and $0 \leq \alpha \leq \pi/2$. [2]

ii) Prove that the particle moves in simple harmonic motion. [2]

iii) Find when the particle is 2m to the right of the origin. [2]
(correct to 2 decimal places)

Question 12 continues on the next page

Question 13 (Begin a New Page)

(15 Marks)

(a) The function $f(x)$ is given by $f(x) = \sqrt{x+6}$ for $x \geq -6$


- i) Find the inverse function $f^{-1}(x)$ and find its domain. [2]
- ii) On the same diagram, sketch the graphs of $y = f(x)$ and $y = f^{-1}(x)$. Showing Clearly all the intercepts on the coordinates axes. [2]
- iii) Show that the x coordinates of any points of intersection of the graphs of $y = f(x)$ and $y = f^{-1}(x)$ satisfy the equation $x^2 - x - 6 = 0$. [1]
- iv) Hence find the point of the intersection of the two graphs. [1]

(b) A vertical flagpole CD of height h metres stands with its base C on horizontal ground. A is a point on the ground due west of C and B is a point on the ground 40 metres due south of A . From A and B the angles of elevation of the top D of the flagpole are 20° and 10° respectively.

- i) Draw a diagram for the information given [1]
- ii) Find the height of the flagpole to the nearest metre. [3]

Question 13 continues on the next page

- (c) Two concentric circles with centre O have radii 2 cm and 4 cm . The points P and Q lie on the larger Circle and $\angle POQ = x$, where $0 \leq x \leq \frac{\pi}{2}$



- i) If the area $A\text{ cm}^2$ of the shaded region is $\frac{1}{16}$ the area of the larger circle, show that x satisfies the equation $8 \sin x - 2x - \pi = 0$. [1]
- ii) Show that this equation has a solution $x = \alpha$, where $0.5 \leq \alpha \leq 0.6$ [2]
- iii) Taking 0.6 as a first approximation for α , use one application of Newton's method to find a second approximation, giving the answer correct to 2 decimal places. [2]

End of Question 13

Question 14 (Begin a New Page)

(15 Marks)

(a) A particle moves in a straight line. At time t seconds its displacement is x metres from a fixed point O on the line, its acceleration is $a \text{ ms}^{-2}$, and its velocity is $v \text{ ms}^{-1}$, where v is given by $v = \frac{32}{x} - \frac{x}{2}$. Initially the particle is at $x = 2$.


i) Find an expression for a in terms x . [2]

ii) Show that $t = \int \frac{2x}{64-x^2} dx$, and hence show that $x^2 = 64 - 60e^{-t}$. [3]

iii) Sketch the graph of x^2 against t and describe the limiting behaviour of the particle. [1]

(b) $P(2t, t^2)$ is a point on the parabola $x^2 = 4y$ with focus F . The point M divides the interval FP externally in the ratio $3:1$. Show that as P moves on the parabola $x^2 = 4y$, then the locus of M is given by $x^2 = 6y + 3$. [3]

Question 14 continues on the next page



- (c) The diagram above shows two particles A and B projected from the origin. Particle A is projected with initial velocity V m/s at an angle α and Particle B is projected T seconds later with the same initial velocity V m/s but an angle of β . The particles collide at the point R .

- i) You may assume that the equation of the path of A is given by

$$y = -\frac{gx^2}{2V^2} \sec^2 \alpha + x \tan \alpha$$

Write down the equation of the path of B .

[1]

- ii) Show that the x -coordinate of the collision point R is given by

$$x = \frac{2V^2 \cos \alpha \cos \beta}{g \sin(\alpha + \beta)}$$

[2]

- iii) You may assume that the horizontal displacement of A after t seconds is given by

$$x = Vt \cos \alpha$$

Write down the equation for the horizontal displacement of B .

[1]

- iv) Show that, for the collision to take place, the value of T is given by

$$T = \frac{2V(\cos \beta - \cos \alpha)}{g \sin(\alpha + \beta)}$$

[2]

End of paper

BLANK PAGE

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$





Sydney Girls High School

Mathematics Faculty

Years 12 HSC Mathematics Extension 1 2013 Trial

Question	Marker's Comment
11	<p>This question overall was completed very average. Some students answered it very well and others very poorly, there was a large range.</p> <ul style="list-style-type: none">a) Substitution and integration was done well but substituting back in terms of x was very poor.b) Most students realised to multiply both sides by $(x-1)^2$ but then most couldn't complete and solve.c) Most students didn't know correct general solution and some even failed to recognise to substitute $\cos 2x = 2\cos^2 x - 1$ to be able to factorise and solve.d) Overall was done well, most knew domain and range. Shaded area was also done well.e) i) To obtain the mark both : $r = \tan^2 x$ and if $0 < x < \frac{\pi}{4}$ then $0 < \tan^2 x < 1$ had to be stated. ii) Was done very well.
12	<ul style="list-style-type: none">b) iii) Many students successfully found the derivative but forgot to substitute for t and thus did not answer the question.c) ii) All calculus answers involving trigonometric functions must be in terms of radians. Radians are a linear measure, degrees are a measure of turning.d) There were many solutions to this question because of the number of equal angles in the diagram. Deductive solutions should be a logical progression of statements to a conclusion. Students who simply wrote down everything they could see from the diagram without a logical progression failed to gain full marks.
13	<p>This question was done well. Only some students had difficulties with inverse functions and sketching these functions. A number of students couldn't do the three dimensional trigonometry.</p>
14	<p>This question was done very poorly. Many students didn't know the formula $a = \frac{d}{dx} \frac{1}{2} v^2$ or didn't know how to use it properly. Many students have no idea how to show a given formula in all parts of this question.</p>



Sydney Girls High School Mathematics Faculty

Multiple Choice Answer Sheet – Trial HSC 2013
Extension 1



Student Number: Answers

Completely fill the response oval representing the most correct answer.

1. A B C D
2. A B C D
3. A B C D
4. A B C D
5. A B C D
6. A B C D
7. A B C D
8. A B C D
9. A B C D
10. A B C D *NO ANSWER*

Section II

Question 11.

a) $u^2 = x+1$

$u = (x+1)^{1/2}$

$$\frac{du}{dx} = \frac{1}{2}(x+1)^{-1/2}$$

$$\frac{du}{dx} = \frac{1}{2\sqrt{x+1}}$$

$$2du = \frac{dx}{\sqrt{x+1}}$$

$$\therefore \int \frac{2x+2}{\sqrt{x+1}} dx = 2 \int (u^2+1) du$$


$$= 2 \left[\frac{u^3}{3} + u \right] + C$$

$$= 2 \left[\frac{(x+1)^{3/2}}{3} + (x+1)^{1/2} \right] + C$$

$$= 2\sqrt{x+1} \left[\frac{(x+1)^3}{3} + 1 \right] + C$$

b) $(x-1)^2 (x+2) < \frac{4}{(x-1)^2}$

$$(x-1)(x+3)(x-2) < 0$$



$$\therefore x < -3$$

$$1 < x < 2$$

$$(x-1)^2 (x+2) < 4(x-1)$$

$$(x-1)^2 (x+2) - 4(x-1) < 0$$

$$(x-1) [(x-1)(x+2) - 4] < 0$$

$$(x-1) [x^2 - x + 2x - 2 - 4] < 0$$

$$(x-1) [x^2 + x - 6] < 0$$

c) General solution

$$\cos 2x = \cos x$$

$$\cos 2x - \cos x = 0$$

$$2\cos^2 x - 1 - \cos x = 0$$

$$2\cos^2 x - \cos x - 1 = 0$$

$$(2\cos x + 1)(\cos x - 1) = 0$$

$$2\cos x = -1 \quad \cos x = 1$$

$$\cos x = -\frac{1}{2} \quad x = \cos^{-1}(1)$$

$$x = \cos^{-1}\left(-\frac{1}{2}\right) \quad x = 0$$

$$x = \frac{2\pi}{3} \quad (120^\circ)$$

$$\therefore x = 2n\pi \pm \frac{2\pi}{3} \quad x = 2n\pi$$

d) i) $y = 3 \sin^{-1}\left(\frac{x}{2}\right)$

Domain: $y = \sin^{-1} x \quad -1 \leq x \leq 1$


Domain: $y = 3 \sin^{-1} \frac{x}{2} \quad -1 \leq \frac{x}{2} \leq 1$


$$-2 \leq x \leq 2$$

Range: $y = \sin^{-1} x \quad -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}$

Range: $y = 3 \sin^{-1} \frac{x}{2} \quad -\frac{3\pi}{2} \leq 3 \sin^{-1} \frac{x}{2} \leq \frac{3\pi}{2}$

$$-\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$$





d) ii)

$$y = 3 \sin^{-1} \frac{x}{2}$$

at $x=1$, $y = 3 \sin^{-1} \frac{1}{2}$

$$y = 3 \sin^{-1} \frac{1}{2}$$

$$y = \frac{\pi}{2}$$

$$\sin\left(\frac{y}{3}\right) = \frac{x}{2}$$

$$2 \sin\left(\frac{y}{3}\right) = x$$

Shaded

Area = Area of rectangle $- \int_0^{\frac{\pi}{2}} 2 \sin \frac{y}{3} dy$

$$= \left(1 \times \frac{\pi}{2}\right) - 2 \times 3 \left[-\cos \frac{y}{3}\right]_0^{\pi/2}$$

$$= \frac{\pi}{2} - 6 \left[-\cos \frac{\pi}{6} - (-\cos 0)\right]$$

$$= \frac{\pi}{2} - 6 \left[-\frac{\sqrt{3}}{2} + 1\right]$$

$$= \frac{\pi}{2} + 6\frac{\sqrt{3}}{2} - 6$$

$$= \frac{\pi}{2} + 3\sqrt{3} - 6 \text{ units}^2$$

$$\approx 0.77 \text{ units}^2$$

e) i) $\tan x + \tan^3 x + \tan^5 x + \dots$ $0 \leq x \leq \frac{\pi}{4}$

Common ratio $\frac{\tan^3 x}{\tan x} = \frac{\tan^5 x}{\tan^3 x}$

$$\therefore r = \tan^2 x$$

limiting sum exists $-1 < |r| < 1$

if $0 < x < \frac{\pi}{4}$ then

$$0 < \tan^2 x < 1$$

\therefore limiting sum exists

$$ii) S_{\infty} = \frac{a}{1-r}$$

$$= \frac{\tan x}{1 - \tan^2 x}$$

$$= \frac{1}{2} \left(\frac{2 \tan x}{1 - \tan^2 x} \right)$$

$$= \frac{1}{2} \tan 2x$$

Question 12

a) When $n=1$
 $\therefore 5^1 + 2(11^1) = 27$ which is a multiple of 3

Assume true for $n=k$
 $5^k + 2(11^k) = 3m$ (an integer)

Prove true for $n=k+1$
 $5^{k+1} + 2(11^{k+1}) = 5 \times 5^k + 22 \times 11^k$
 $= 5(3m - 2(11^k)) + 22 \times 11^k$
 $= 15m - 10(11^k) + 22 \times 11^k$
 $= 15m + 12(11^k)$
 $= 3(5m + 4(11^k))$ which is a multiple of 3

Hence it is true for $n=k$, true for $n=k+1$
 True for $n=1$, hence true for $n \in \mathbb{N}$

b) i) Initially $t=0$
 then $N_A = 15 + 20e^0 = 35$

ii) $\frac{dN_A}{dt} = -20e^{-0.5t}$
 when $t=0$
 $\frac{dN_A}{dt} = -20$

iii) $N_A = N_0 e^{-0.5t} = 15 + 20e^{-0.5t} = 5 + 40e^{-0.5t}$
 $10 = 20e^{-0.5t}$
 $-0.5t = \log_e \left(\frac{1}{2}\right)$
 $t = 2 \log_e(2) = 1.39 \text{ min}$

c) $1) R = \sqrt{6^2 + (2\sqrt{3})^2}$
 $= \sqrt{48}$
 $= 4\sqrt{3}$

$R \sin(2t - \alpha) = 4\sqrt{3} \sin 2t \cos \alpha - 4\sqrt{3} \cos 2t \sin \alpha$
 $= 6 \sin 2t - 2\sqrt{3} \cos 2t$
 $\therefore 4\sqrt{3} \cos \alpha = 6$
 $\cos \alpha = \frac{6}{4\sqrt{3}} = \frac{\sqrt{3}}{2}$
 $\alpha = \frac{\pi}{6}$

ii) $\hat{r} = 4\sqrt{3} \sin(2t - \frac{\pi}{6})$
 $\hat{r} = 8\sqrt{3} \cos(2t - \frac{\pi}{6})$
 $\hat{r} = -16\sqrt{3} \sin(2t - \frac{\pi}{6})$
 $= -16r$ which is in the form $\ddot{r} = -\omega^2 r$

iii) when $r=2$
 $4\sqrt{3} \sin(2t - \frac{\pi}{6}) = 2$
 $\sin(2t - \frac{\pi}{6}) = \frac{1}{2\sqrt{3}}$
 $2t - \frac{\pi}{6} = \sin^{-1}\left(\frac{1}{2\sqrt{3}}\right) + \frac{\pi}{6}$
 $t = \frac{\sin^{-1}\left(\frac{1}{2\sqrt{3}}\right) + \frac{\pi}{6}}{2}$
 $= 0.408 \text{ s}$
 $= 0.41 \text{ seconds}$ (must be in radians)

d) Let $\angle DAC = \pi$
 then $\angle ADC = \pi$ Base \angle 's in ΔADC
 then $\angle CAB = \angle ADC$ (\angle in alt segment) $= \pi$
 $\angle CAB = \angle ACD$ Alt \angle 's $AB \parallel EC$
 $= \pi$


Also $\angle DEB = \pi$ ($= \angle$'s on chord DC)
 $\therefore ED \parallel AC$ (equal alt \angle 's)
 There are many variations on this proof

13) $y = \sqrt{x+6}$

i) $x = \sqrt{y+6}$

$y = x^2 - 6$

D: $x \geq 0$



iii) $x^2 - 6 = x$

$x^2 - x - 6 = 0$


iv) $x^2 - x - 6 = 0$

$(x-3)(x+2) = 0$

$x = 3 \quad x \neq -2$

$y = 3 \quad P(3,3)$

b)



$\tan 20 = \frac{h}{AC}$

$\tan 10 = \frac{h}{BC}$

$AC^2 + AB^2 = BC^2$

$\left(\frac{h}{\tan 20}\right)^2 + 40^2 = \left(\frac{h}{\tan 10}\right)^2$

$40^2 = h^2 \left[\frac{1}{(\tan 10)^2} - \frac{1}{(\tan 20)^2} \right]$

$5 h^2 (\tan 20)^2 - h^2 (\tan 10)^2$

$h^2 [(\tan 20)^2 - (\tan 10)^2] = 5 \cdot 40^2 (\tan 10)^2$

$h = 8 \text{ m}$

c) i)

$$\frac{1}{2} 4 \times 4 \sin x - \frac{1}{2} \times 2^2 x = \frac{1}{16} \pi \times 16$$

$$8 \sin x - 2x = \pi$$

$$8 \sin x - 2x - \pi = 0$$

ii) $P(0.5) = 8 \sin(0.5) - 1 - \pi$

$$= -0.306$$

$$P(0.6) = 8(\sin(0.6)) - 1 - \pi$$

$$= 0.1755$$

$$P(0.5) < 0, P(0.6) > 0$$

$\therefore x = \alpha$ is a

solution

iii) $x = 0.6 - \frac{P(0.6)}{P'(0.6)}$

$$= 0.6 - \frac{0.1755}{4.603}$$

$$= 0.516$$

14) a)

$$a = \frac{d}{dx} \frac{1}{2} \sqrt{x}$$

$$\frac{1}{2} \sqrt{x} = \frac{1}{2} \left(\frac{32}{x} - \frac{x}{2} \right)^2$$

$$= \frac{1}{2} \left(\frac{1024}{x^2} - 32 + \frac{x^2}{4} \right)$$

$$= \frac{1024}{2x^2} - 16 + \frac{x^2}{8}$$

$$= \frac{512}{x^2} - 16 + \frac{x^2}{8}$$

$$a = -1024x^{-3} + \frac{x}{4}$$

$$= \frac{1024}{x^3} + \frac{x}{4}$$

$$v = \frac{dx}{dt} \quad \int dt = \int \frac{-2x}{64-x^2}$$

$$v = \frac{32-x}{64-x^2} \quad \int \frac{32-x}{64-x^2} = \ln(64-x^2) + C$$

$$= \frac{2x}{64-x^2} \quad \text{at } t=0 \quad x=2$$

$$0 = \ln(60) + C$$

$$C = \ln 60$$

$$t = \ln(64-x^2) + \ln 60$$

$$-t = \ln(64-x^2) - \ln 60$$


$$t = \ln\left(\frac{64-x^2}{60}\right)$$

$$e^{-t} = \frac{64-x^2}{60}$$

$$60e^{-t} = 64-x^2$$

$$x^2 = 64 - 60e^{-t}$$

iii)



$$b) M = \frac{-3 \times 2t + 1 \times 0}{-3+1} = \frac{-3t^2+1}{-2}$$

$$(x, y) = \frac{-6t}{-2}, \frac{-3t^2+1}{-2}$$

$$x = 3t \rightarrow t = \frac{x}{3}$$

$$y = \frac{3t^2+1}{2}$$

$$\text{at } t = \frac{x}{3}$$

$$y = \frac{3\left(\frac{x}{3}\right)^2+1}{2} = \frac{3x^2+1}{6}$$

$$6y = x^2+3$$

$$y = \frac{3x^2+1}{6} \quad x^2 = 6y+3$$

$$18y = 3x^2+9$$

c) i)

$$y = \frac{-gx^2}{2v^2} \sec^2 B + x \tan B$$

$$ii) \frac{-gx^2}{2v^2} \sec^2 B + x \tan B = \frac{-gx^2}{2v^2} \sec^2 \alpha + x \tan \alpha$$

$$\frac{-gx^2}{2v^2} \sec^2 B + \frac{gx^2}{2v^2} \sec^2 \alpha = x \tan \alpha - x \tan B$$

$$\frac{gx^2}{2v^2} (\sec^2 \alpha - \sec^2 B) = x (\tan \alpha - \tan B)$$

$$\frac{gx}{2v^2} (1 + \tan^2 \alpha - 1 - \tan^2 B) = \tan \alpha - \tan B$$

$$\frac{gx}{2v^2} (\tan \alpha - \tan B) (\tan \alpha + \tan B) = \tan \alpha - \tan B$$

$$\frac{gx}{2v^2} = \frac{1}{\tan \alpha + \tan B}$$

$$x = \frac{2v^2}{g} \cdot \frac{1}{\frac{\sin \alpha + \sin B}{\cos \alpha \cos B}}$$

$$= \frac{2v^2}{g} \cdot \frac{1}{\frac{\sin \alpha \cos B + \sin B \cos \alpha}{\cos \alpha \cos B}}$$

$$= \frac{2v^2 \cos \alpha \cos B}{g \sin(\alpha + B)}$$

$$iii) x_B = V(t-T) \cos B$$

$$iv) \sqrt{t \cos \alpha} = \frac{2\sqrt{v \cos \alpha \cos B}}{g \sin(\alpha + B)}$$

$$V(t-T) \cos B = \frac{2v^2 \cos \alpha \cos B}{g \sin(\alpha + B)}$$

$$g \sin(\alpha + B)$$

From ①

$$t = \frac{2v \cos B}{g \sin(\alpha + B)}$$

$$g \sin(\alpha + B)$$

From ②

$$t-T = \frac{2v \cos \alpha}{g \sin(\alpha + B)}$$

$$T = \frac{2v \cos B}{g \sin(\alpha + B)} - \frac{2v \cos \alpha}{g \sin(\alpha + B)}$$

$$= \frac{2v(\cos B - \cos \alpha)}{g \sin(\alpha + B)}$$

