



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2015**

**HIGHER SCHOOL CERTIFICATE  
TRIAL PAPER**

# Mathematics

# Extension 1

## *General Instructions*

- Reading time – 5 minutes.
- Working time – 2 hours.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for messy or badly arranged work.
- Leave your answers in the simplest exact form, unless otherwise stated.
- Start each **NEW** question in a separate answer booklet.

## **Total Marks – 70**

### **Section I**

Pages 2–4

#### **10 Marks**

- Attempt Questions 1–10
- Allow about 15 minutes for this section.

### **Section II**

Pages 6–11

#### **60 marks**

- Attempt Questions 11–14
- Allow about 1 hour and 45 minutes for this section

Examiners: *R. Elliot & J. Chen*

**This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.**

## Section I

10 marks

Attempt Questions 1–10


Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10.

---

- 1 The roots of  $3x^3 - 2x^2 + x - 1 = 0$  are  $\alpha$ ,  $\beta$  and  $\gamma$ .  
What is the value of  $\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2$ ?
- (A)  $-\frac{1}{9}$   
(B)  $-\frac{2}{9}$   
(C) 1  
(D)  $\frac{2}{9}$
- 2 What is the minimum value of  $\sqrt{7}\sin x - 3\cos x$ ?
- (A) -2  
(B) -4  
(B) -16  
(D)  $\sqrt{7} - 3$
- 3 What is the domain and range of  $y = \sin^{-1}\left(\frac{2x}{5}\right)$ ?
- (A) Domain:  $-1 \leq x \leq 1$ ; Range:  $-\pi \leq y \leq \pi$   
(B) Domain:  $-1 \leq x \leq 1$ ; Range:  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$   
(C) Domain:  $-\frac{5}{2} \leq x \leq \frac{5}{2}$ ; Range:  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$   
(D) Domain:  $-\frac{5}{2} \leq x \leq \frac{5}{2}$ ; Range:  $-\pi \leq y \leq \pi$
- 4 Evaluate  $\lim_{x \rightarrow 0} \frac{\sin 3x}{2x}$ .
- (A) 0  
(B)  $\frac{2}{3}$   
(C) 1  
(D)  $\frac{3}{2}$

- 5 In the diagram below,  $AB$  is a tangent to the circle  $BCD$ .  
 Also,  $CD$  is a tangent to the circle  $ABD$ .  
 $\angle BAD = \theta$  and  $\angle BCD = \phi$ .



Which of the following is a true statement?

- (A)  $\triangle ABD \equiv \triangle BDC$
- (B)  $ABCD$  is a cyclic quadrilateral
- (C)  $\triangle ABD \parallel \triangle BDC$
- (D)  $AB \parallel CD$
- 6 A particle moves in simple harmonic motion so that its velocity,  $v$ , is given by

$$v^2 = 6 - x - x^2.$$

Between which two points does it oscillate?

- (A)  $x = 6$  and  $x = 3$
- (B)  $x = -2$  and  $x = 3$
- (C)  $x = 1$  and  $x = 2$
- (D)  $x = 2$  and  $x = -3$
- 7 Which of the following is an expression for  $\int \cos^3 x \sin x \, dx$ ?
- (A)  $-\cos^4 x + c$
- (B)  $-\frac{1}{4}\cos^4 x + c$
- (C)  $\cos^4 x + c$
- (D)  $\frac{1}{4}\cos^4 x + c$

- 8 Which of the following is the correct expression for the inverse of  $f(x) = e^{1-2x}$ ?
- (A)  $f^{-1}(x) = -2e^{1-2x}$
- (B)  $f^{-1}(x) = -\frac{1}{2}e^{1-2x}$
- (C)  $f^{-1}(x) = -\frac{1}{2}\log_e(1-2x)$
- (D)  $f^{-1}(x) = \frac{1}{2}(1 - \log_e x)$
- 9 Three Mathematics study guides, four Mathematics textbooks and five exercise books are randomly placed along a bookshelf. What is the probability that the Mathematics textbooks are all next to each other?
- (A)  $\frac{4!}{12!}$
- (B)  $\frac{9!}{12!}$
- (C)  $\frac{4!3!5!}{12!}$
- (D)  $\frac{4!9!}{12!}$
- 10 A particle moves on the  $x$ -axis with velocity  $v$  m/s, such that  $v^2 = 16x - x^2$ . Which of the following is the particle's maximum speed and the position of where this maximum speed occurs?
- (A) Maximum speed = 16 m/s at  $x = 0$
- (B) Maximum speed = 8 m/s at  $x = -8$
- (C) Maximum speed =  $-8$  m/s at  $x = 8$
- (D) Maximum speed = 8 m/s at  $x = 8$

## Section II

60 marks

Attempt Questions 11–14

Allow about 1 hour and 45 minutes for this section

Answer each question in a NEW writing booklet. Extra pages are available

In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.

---

**Question 11** (15 Marks)      Start a NEW Writing Booklet

- (a) Differentiate  $\sin^{-1}(\log_e x)$ . 1
- (b) Find  $\int \frac{1}{\sqrt{4-9x^2}} dx$ . 1
- (c) (i) Simplify  $\sin(A+B) + \sin(A-B)$ . 1
- (ii) Hence, evaluate  $\int_0^{\frac{\pi}{6}} \sin 3x \cos x dx$ . 2
- (d) The point  $P(6p, 3p^2)$  is a point on the parabola  $x^2 = 12y$ .
- (i) Find the equation of the tangent at  $P$ . 2
- (ii) The tangent at  $P$  cuts the  $y$ -axis at  $B$ . 3  
The point  $A$  divides  $PB$  internally in the ratio 1 : 2.  
Find the locus of the point  $A$  as  $P$  varies.
- (e) A piece of meat at temperature  $T^\circ \text{C}$  is placed in an oven, which has a constant temperature of  $H^\circ \text{C}$ .  
The rate at which the temperature of the meat warms is given by
- $$\frac{dT}{dt} = -K(T - H),$$
- where  $t$  is in minutes and for some positive constant  $K$ .
- (i) Show that  $T = H + Be^{-Kt}$ , where  $B$  is a constant, is a solution of the differential equation above. 1
- (ii) If the meat warms from  $10^\circ \text{C}$  to  $50^\circ \text{C}$  in the oven, which has a constant temperature of  $180^\circ \text{C}$ , in 30 minutes, find the value of  $K$ . 2
- (iii) How long will it take the meat to get to a temperature of  $150^\circ \text{C}$ ? 2  
Express your answer correct to the nearest minute.

**Question 12** (15 Marks)      Start a NEW Writing Booklet

(a) (i) Solve  $\cos x - \sqrt{3} \sin x = 1$  for  $0 \leq x \leq 2\pi$ . **2**

(ii) Hence, or otherwise, find a general solution to  $\cos x - \sqrt{3} \sin x = 1$ . **1**

(b) (i) On the same set of axes sketch the graphs of  $y = \cos 2x$  and  $y = \frac{x+1}{3}$  **2**

(ii) Use the graph to determine the number of solutions to the equation **1**

$$3\cos 2x = x + 1$$

(iii) One solution of the equation  $3\cos 2x = x + 1$  is close to 0.5. **3**  
Use one application of Newton's Method to find another approximation, correct to 3 decimal places.

(c) Evaluate  $\int_0^{\frac{\pi}{4}} \sin^2 2x \, dx$  **3**

(d) When  $x$  cm from the origin, the acceleration of a particle moving in a straight line is given by: **3**


$$\frac{d^2x}{dt^2} = -\frac{5}{(x+2)^3}$$

It has an initial velocity of 2 cm/s at  $x = 0$ . If the velocity is  $V$  cm/s, find  $V$  in terms of  $x$ .

**Question 13** (15 Marks)      Start a NEW Writing Booklet

- (a) In the diagram below,  $DC$  is a diameter of the larger circle centred at  $A$ . 4  
 $AC$  is a diameter of the smaller circle centred at  $B$ .  
 $DE$  is tangent to the smaller circle at  $F$  and  $DC = 12$ .

Copy the diagram to your answer booklet.  
 Determine the length of  $DE$ .



- (b) (i) Simplify  $k! + k \times k!$  1  
 (ii) Prove, by mathematical induction, that 3

$$1 \times 1! + 2 \times 2! + 3 \times 3! + \dots + n \times n! = (n+1)! - 1$$


for all positive integers  $n$ .

- (c) (i) Using the substitution  $x = 3 + 3\sin\theta$  find  $\int \sqrt{x(6-x)} \, dx$  4  
 (ii) Let  $R$  be the region bounded by the curve  $y = \sqrt{x(6-x)}$  and the  $x$ -axis. 3  
 Find the volume of the solid of revolution generated by revolving  $R$  about the  $x$ -axis.

**End of Question 13**

**Question 14** (15 Marks)      Start a NEW Writing Booklet

(a)



The figure above shows an inverted conical cup with base radius 8 cm and height 10 cm.

Some water is poured into the cup at a constant rate of  $\frac{2\pi}{5}$  cm<sup>3</sup> per minute.

Let the depth of the water be  $h$  cm at time  $t$  minutes.

Find the rate of change in the area of the water surface when  $h = 4$

**3**

- (b) A particle is projected horizontally at  $30 \text{ ms}^{-1}$  from the top of a 100 m high wall. Assume that acceleration due to gravity is  $10 \text{ ms}^{-2}$  and that there is no air resistance.

The flight path of the particle is given by:

$$x = 30t, y = 100 - 5t^2 \text{ (Do NOT prove this)}$$

where  $t$  is the time in seconds after take-off.

- (i) Find the time taken for the particle to reach the ground.

**1**

- (ii) Find the angle and speed at which the particle strikes the ground.


**2**

**Question 14 continues on page 11**



Question 14 (continued)

- (c) The diagram below shows a tetrahedron such that  $VA = VB = AB = 2a$ ,  
 $CA = CB = 3a$  and  $VC = \sqrt{5}a$ .  
 $O$  is the foot of the perpendicular from  $V$  to the base  $ABC$ .  
 $M$  is the midpoint of  $AB$ .  
 $P$  is a point on  $BC$  such that  $BP = ra$  where  $0 \leq r \leq 3$ .  
 $\angle VMC = \theta$  and  $\angle VPO = \varphi$ .



- (i) By considering  $\triangle VMC$ , show that  $\cos \theta = \frac{\sqrt{6}}{4}$ . 3
- (ii) Hence find the exact value of  $VO$ . 1
- (iii) Show that  $VP^2 = \frac{1}{3}(3r^2 - 8r + 12)a^2$  2
- (iv) Hence show that  $\sin \varphi = \sqrt{\frac{45}{8(3r^2 - 8r + 12)}}$  1
- (v) Hence, or otherwise, find the maximum value of  $\varphi$  as  $r$  varies. 2

**End of paper**



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2015**

**HIGHER SCHOOL CERTIFICATE  
TRIAL PAPER**

# Mathematics Extension 1

## Sample Solutions

Question	Teacher
Q11	<b>RB</b>
Q12	<b>BK</b>
Q13	<b>BD</b>
Q14	<b>PB</b>

### MC Answers

Q1	D
Q2	B
Q3	C
Q4	D
Q5	C
Q6	D
Q7	B
Q8	D
Q9	D
Q10	D

Section I 10 marks

---

- 1 The roots of  $3x^3 - 2x^2 + x - 1 = 0$  are  $\alpha$ ,  $\beta$  and  $\gamma$ .  
What is the value of  $\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2$ ?

- (A)  $-\frac{1}{9}$   
(B)  $-\frac{2}{9}$   
(C) 1  
(D)  $\frac{2}{9}$

**ANSWER: D**

$$3x^3 - 2x^2 + x - 1 = 0$$

$$\alpha\beta\gamma = -\frac{d}{a} \quad \alpha + \beta + \gamma = -\frac{b}{a}$$
$$= \frac{1}{3} \quad = \frac{2}{3}$$

$$\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2 = \alpha\beta\gamma(\alpha + \beta + \gamma)$$
$$= \frac{1}{3} \times \frac{2}{3}$$
$$= \frac{2}{9}$$

- 2 What is the minimum value of  $\sqrt{7}\sin x - 3\cos x$ ?

- (A) -2  
(B) -4  
(B) -16  
(D)  $\sqrt{7} - 3$

**ANSWER: B**

$$\sqrt{7}\sin x - 3\cos x$$

$$r = \sqrt{(\sqrt{7})^2 + 3^2}$$

$$= \sqrt{7+9}$$

$$= 4$$

$$\text{Let } \sqrt{7}\sin x - 3\cos x = r\sin(\theta - \alpha)$$
$$= 4\sin(\theta - \alpha)$$

No matter what the value of  $\alpha$

$$-1 \leq \sin(x - \alpha) \leq 1$$

$$-4 \leq 4\sin(x - \alpha) \leq 4$$

Therefore the minimum value is  $x = -4$ .

3 What is the domain and range of  $y = \sin^{-1}\left(\frac{2x}{5}\right)$ ?

(A) Domain:  $-1 \leq x \leq 1$ ; Range:  $-\pi \leq y \leq \pi$

(B) Domain:  $-1 \leq x \leq 1$ ; Range:  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

(C) Domain:  $-\frac{5}{2} \leq x \leq \frac{5}{2}$ ; Range:  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

(D) Domain:  $-\frac{5}{2} \leq x \leq \frac{5}{2}$ ; Range:  $-\pi \leq y \leq \pi$

**ANSWER: C**

$$y = \sin^{-1}\left(\frac{2x}{5}\right) \Rightarrow \sin y = \left(\frac{2x}{5}\right)$$

Domain:  $-1 \leq \sin y \leq 1$

$$-1 \leq \frac{2x}{5} \leq 1$$

$$-\frac{5}{2} \leq x \leq \frac{5}{2}$$

Range:  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$  as this is the range of

$y = \sin^{-1} x$

4 Evaluate  $\lim_{x \rightarrow 0} \frac{\sin 3x}{2x}$ .

(A) 0

(B)  $\frac{2}{3}$

(C) 1

(D)  $\frac{3}{2}$

**ANSWER: D**


$$\lim_{x \rightarrow 0} \frac{\sin 3x}{2x} = \frac{3}{2} \lim_{x \rightarrow 0} \frac{\sin 3x}{2x} \times \frac{2}{3}$$

$$= \frac{3}{2} \lim_{x \rightarrow 0} \frac{2 \sin 3x}{2 \times 3x}$$

$$= \frac{3}{2} \times 1$$

$$= \frac{3}{2}$$

- 5 In the diagram below,  $AB$  is a tangent to the circle  $BCD$ .  
Also,  $CD$  is a tangent to the circle  $ABD$ .  
 $\angle BAD = \theta$  and  $\angle BCD = \phi$ .



Which of the following is a true statement?

- (A)  $\triangle ABD \cong \triangle BDC$   
 (B)  $ABCD$  is a cyclic quadrilateral  
 (C)  $\triangle ABD \parallel \triangle BDC$   
 (D)  $AB \parallel CD$

**ANSWER: C**

In  $\triangle ABD$  and  $\triangle DCB$  :  
 $\angle DCB = \angle DBA$  (angle in the alternate segment)  
 ie  $\angle DCB = \angle ABD$   
 $\angle BAD = \angle BDC$  (angle in the alternate segment)  
 ie  $\angle BAD = \angle CDB$   
 $BD$  is but not respective to angles.  
 Therefore  $\triangle ABD \not\cong \triangle DCB$

Hence, triangles are equiangular they are similar and  $\triangle ABD \parallel \triangle DCB$

- 6 A particle moves in simple harmonic motion so that its velocity,  $v$ , is given by

$$v^2 = 6 - x - x^2.$$

Between which two points does it oscillate?

- (A)  $x = 6$  and  $x = 3$   
 (B)  $x = -2$  and  $x = 3$   
 (C)  $x = 1$  and  $x = 2$   
 (D)  $x = 2$  and  $x = -3$

**ANSWER: D**

$v^2 = 6 - x - x^2$   
 For the particle to reach its oscillation points  $v = 0$ .  
 $v^2 = 6 - x - x^2$   
 $0 = 6 - x - x^2$   
 $0 = (3 + x)(2 - x)$   
 $\therefore x = -3$  and  $2$

7 Which of the following is an expression for  $\int \cos^3 x \sin x \, dx$ ?

(A)  $-\cos^4 x + c$

(B)  $-\frac{1}{4}\cos^4 x + c$

(C)  $\cos^4 x + c$

(D)  $\frac{1}{4}\cos^4 x + c$

**ANSWER: B**

$\int \cos^3 x \sin x \, dx$ , testing solutions:

$$\frac{d}{dx}(\cos^4 x) = 4\cos^3 x \times -\sin x$$

$$\frac{d}{dx} -\frac{1}{4}(\cos^4 x) = \cos^3 x \sin x$$

$$-\frac{1}{4}\cos^4 x = \int \cos^3 x \sin x \, dx$$

8 Which of the following is the correct expression for the inverse of  $f(x) = e^{1-2x}$ ?

(A)  $f^{-1}(x) = -2e^{1-2x}$

(B)  $f^{-1}(x) = -\frac{1}{2}e^{1-2x}$

(C)  $f^{-1}(x) = -\frac{1}{2}\log_e(1-2x)$

(D)  $f^{-1}(x) = \frac{1}{2}(1 - \log_e x)$

**ANSWER: D**

Let  $y = e^{1-2x}$

$$y = e^{1-2x}$$

$$\ln y = 1 - 2x$$

$$2x = 1 - \ln y$$

$$x = \frac{1}{2}(1 - \ln y)$$

$$\therefore f^{-1}(x) = \frac{1}{2}(1 - \ln y)$$

- 9 Three Mathematics study guides, four Mathematics textbooks and five exercise books are randomly placed along a bookshelf. What is the probability that the Mathematics textbooks are all next to each other?

(A)  $\frac{4!}{12!}$

(B)  $\frac{9!}{12!}$

(C)  $\frac{4!3!5!}{12!}$

(D)  $\frac{4!9!}{12!}$

**ANSWER: D**

Since there are 9 elements counting the textbooks as 1 element, hence these can be arranged in  $9!$  ways. Also the textbooks can be arranged in  $4!$  ways.

As there are 12 separate elements, the divisor for population can be counted in  $12!$  ways.

Therefore, the probability is  $\frac{9!4!}{12!}$

- 10 A particle moves on the  $x$ -axis with velocity  $v$  m/s, such that  $v^2 = 16x - x^2$ . Which of the following is the particle's maximum speed and the position of where this maximum speed occurs?

(A) Maximum speed = 16 m/s at  $x = 0$

(B) Maximum speed = 8 m/s at  $x = -8$

(C) Maximum speed =  $-8$  m/s at  $x = 8$

(D) Maximum speed = 8 m/s at  $x = 8$

**ANSWER: D**

$$v^2 = 16x - x^2$$

$$\frac{1}{2}v^2 = 8x - \frac{x^2}{2}$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = 8 - \frac{2x}{2}$$

$$= 8 - x$$

$$\ddot{x} = 8 - x$$

When  $\ddot{x} = 0$  the speed is the greatest so,  
 $\ddot{x} = 8 - x$

$$0 = 8 - x$$

$$x = 8$$

At  $x = 8$ ,

$$v^2 = 16x - x^2$$

$$= 16(8) - (8)^2$$

$$= 64$$

$$v = \pm 8$$

As  $v$ , velocity can take positive and negative values, but the speed can only be positive, the maximum speed is 8 m/s.

(a) let  $y = \sin^{-1}(\ln x)$

$$y' = \frac{1}{\sqrt{1-(\ln x)^2}} \times \frac{1}{x}$$

$$= \frac{1}{x\sqrt{1-(\ln x)^2}}$$

generally well answered but some students forgot the  $\frac{1}{x}$ .

15

① Others thought  $(\ln x)^2 = 2\ln x$  or  $\ln x^2$  No!

(b)  $\int \frac{1 dx}{\sqrt{4-9x^2}} = \frac{1}{3} \sin^{-1}\left(\frac{3x}{2}\right) + C$  (1)

generally well done but some students left off the  $\frac{1}{3}$ .

(c) (i)  $\sin(A+B) + \sin(A-B)$   
 $\sin A \cos B + \cos A \sin B + \sin A \cos B - \cos A \sin B$

$= 2 \sin A \cos B$  (1) *this part, very well answered by 95% of students.*

(ii)  $\int_0^{\frac{\pi}{6}} \sin 3x \cos x dx$

$= \frac{1}{2} \int_0^{\frac{\pi}{6}} 2 \sin 3x \cos x dx$

$= \frac{1}{2} \int_0^{\frac{\pi}{6}} [\sin 4x + \sin 2x] dx$

$= -\frac{1}{2} \left[ \frac{1}{4} \cos 4x \right]_0^{\frac{\pi}{6}} + -\frac{1}{2} \left[ \frac{1}{2} \cos 2x \right]_0^{\frac{\pi}{6}}$

$= -\frac{1}{8} \cos 4x \Big|_0^{\frac{\pi}{6}} - \frac{1}{4} \cos 2x \Big|_0^{\frac{\pi}{6}}$



(ii) (d) (ii)  $\left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right)$  *ratio formula often messed up!*

$P(6p, 3p^2)$   
 $B(0, -3p^2)$

$m:n = 1:2$

$A = \left( \frac{1 \times 0 + 2 \times 6p}{3}, \frac{-3p^2 + 6p^2}{3} \right)$

$A(4p, p^2)$  (1)

*Many students found 'A' but could not find locus equation.*

So  
 $x = 4p \Rightarrow p = \frac{x}{4}$   
 $y = p^2$

So  $y = \frac{x^2}{16}$

$\Rightarrow x^2 = 16y$  is the locus. (1)

(e) (i)  $T = H + Be^{-kt} \Rightarrow Be^{-kt} = T - H$

$\frac{dT}{dt} = 0 - Bke^{-kt}$   
 $= -k(T - H)$  (1)

(ii)  $H = 180^\circ$

When  $t = 0, T = 10^\circ$

$10 = 180 + Be^{-0}$

$10 - 180 = B$

$B = -170$

So  $T = 180 - 170e^{-kt}$

data  
 $t = 30$   
 $T = 50$

*Well answered. Using the method shown. Some boys wanted to integrate the  $\frac{dT}{dt}$  and got into a lot of bother.*

$$\text{So } 50 = 180 - 170e^{-30k}$$

$$-130 = -170e^{-30k}$$

$$\left(\frac{13}{17}\right) = e^{-30k}$$

$$\ln\left(\frac{13}{17}\right) = \ln e^{-30k}$$
$$= -30k$$

$$k = \frac{\ln\left(\frac{13}{17}\right)}{-30} = 0.008942$$

*Lucky k either as exact or approx will give the correct exact or approx. Final answer.*

$$\text{So } T = 180 - 170e^{-0.008942t}$$

②

$$\text{(iii) } 150 = 180 - 170e^{-0.008942t}$$

$$-30 = -170e^{-0.008942t}$$

$$\left(\frac{3}{17}\right) = e^{-0.008942t}$$

$$\ln\left(\frac{3}{17}\right) = \ln e^{-0.008942t}$$

$$= -0.008942t$$

$$t = \frac{\ln\left(\frac{3}{17}\right)}{-0.008942} \doteq 194 \text{ mins}$$

3 hrs 14 mins

*Well answered.*

②

$$\begin{aligned}
& \left( -\frac{1}{8} \cos \frac{4\pi}{6} + \frac{1}{8} \cos 0 \right) - \left( \frac{1}{4} \cos \frac{2\pi}{6} - \frac{1}{4} \cos 0 \right) \\
&= -\frac{1}{8} \cos \frac{2\pi}{3} + \frac{1}{8} - \left( \frac{1}{4} \cos \frac{\pi}{3} - \frac{1}{4} \right) \\
&= -\frac{1}{8} \cos 120^\circ + \frac{1}{8} - \frac{1}{4} \cos 60^\circ + \frac{1}{4} \\
&= -\frac{1}{8} \times -\frac{1}{2} + \frac{1}{8} - \frac{1}{4} \cdot \frac{1}{2} + \frac{1}{4} \\
&= \frac{1}{16} + \frac{1}{8} - \frac{1}{8} + \frac{1}{4} = \frac{1}{16} + \frac{1}{4} = \frac{5}{16} //
\end{aligned}$$

Many students did not use hint from (c)(i) or used it badly. The  $\frac{1}{2}$  out the front was there in most answers. Those that made the correction that (i) can help in (ii), most students worked out the answer. (2)

(d)  $P(6p, 3p^2)$

$$x^2 = 12y$$

$$\begin{aligned}
y - 3p^2 &= px - 6p^2 \\
y &= px - 3p^2 \quad (2)
\end{aligned}$$

(i)  $x^2 = 12y$

$$y = \frac{x^2}{12}$$

$$y' = \frac{2x}{12} = \frac{x}{6}$$

at  $x = 6p$   $m = \frac{6p}{6} = p$

$$(y - 3p^2) = p(x - 6p)$$

well answered.

(ii) Cuts y axis at B.  
 $x = 0$   $y = -3p^2$

$B(0, -3p^2)$  (1)

$P(6p, 3p^2)$ .

easy to find

$$12. (a) (i) \cos x - \sqrt{3} \sin x = 1 \quad 0 \leq x \leq 2\pi$$

$$\cos x - \sqrt{3} \sin x = R \cos(x + \alpha)$$

$$\cos x - \sqrt{3} \sin x = R \cos x \cos \alpha - R \sin x \sin \alpha$$

$$\Rightarrow R = \sqrt{1+3} = 2$$

$$\text{Then } 2 \cos \alpha = 1 \quad \text{and} \quad 2 \sin \alpha = \sqrt{3}$$

$$\cos \alpha = \frac{1}{2} \quad \sin \alpha = \frac{\sqrt{3}}{2}$$

$$\therefore \alpha = \frac{\pi}{3}$$

$$\text{Then } 2 \cos\left(x + \frac{\pi}{3}\right) = 1$$

$$\cos\left(x + \frac{\pi}{3}\right) = \frac{1}{2}$$

$$x + \frac{\pi}{3} = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \dots$$

$$x = 0, \frac{4\pi}{3}, 2\pi, \dots$$

$$(ii) \text{ General Soln: } x = 2n\pi \text{ or } 2n\pi \pm \frac{4\pi}{3}$$


$$\text{or } x = -\frac{\pi}{3} + 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$$

$$\text{or } x = \frac{-2\pi}{3} + 2k\pi \text{ or } 2k\pi, k \in \mathbb{Z}$$

(i) Most students realised they needed the auxiliary angle method. Common errors included:

- evaluating tan incorrectly and having  $1/\sqrt{3}$
- not finding all the solutions in the given domain.

(ii) Some had the incorrect formula.



(ii) 3 solutions ✓

(1)

x	0	2
y	1/3	1

(iii)  $f(x) = 3\cos 2x - x - 1$

Let  $f(0.5) = 1.6209 - 0.5 - 1$   
 $f(0.5) = 0.1209069$

Also  $f'(x) = -6\sin 2x - 1$  ✓  
 $f'(x) = -6.0488$

Then  $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$   
 $= 0.5 - \frac{0.1209069}{-6.0488259}$  ✓

$= 0.5199884907$   
 $x_{n+1} = 0.520$  to 3dp ✓

An inaccurate graph resulted in the wrong number of solutions. Using Newton's Method done well on the whole. Some students did not use the given starting value and so were incorrect.

$$(c) \int_0^{\pi/4} \sin^2 2x \, dx$$

$$\sin^2 \theta = \frac{1}{2}(1 - \cos 2\theta)$$

$$\sin^2 2x = \frac{1}{2}(1 - \cos 4x)$$

$$= \frac{1}{2} \int_0^{\pi/4} (1 - \cos 4x) \, dx$$

$$= \frac{1}{2} \left[ x - \frac{\sin 4x}{4} \right]_0^{\pi/4}$$

$$= \frac{1}{2} \left[ \left( \frac{\pi}{4} - 0 \right) - \left( 0 - 0 \right) \right]$$

$$= \frac{\pi}{8}$$

The most common mistake was not using the double angle formula correctly.

(3)

$$(d) \quad x = \frac{-5}{(x+2)^3}$$

When  $t=0$ ,  $x = x \text{ cm}$ .  
 $t=0$ ,  $v=2$ ,  $x=0$ .

$$\frac{d(\frac{1}{2}v^2)}{dx} = \frac{-5}{(x+2)^3}$$

$$\frac{1}{2}v^2 = -5 \int \frac{1}{(x+2)^3} \, dx$$

$$v^2 = \frac{-10}{-2} (x+2)^{-2} + C$$

$$v^2 = \frac{5}{(x+2)^2} + C$$

When  $x=0$ ,  $v=2 \Rightarrow 4 = \frac{5}{4} + C$   
 $C = 2\frac{3}{4} = \frac{11}{4}$

(d) (cont)

$$v^2 = \frac{5}{(x+2)^2} + \frac{11}{4}$$

$v > 0$  since  $v$  can never be 0

Also when  $t=0$ ,  $v=2$  and

$\frac{d^2x}{dt^2} < 0 \Rightarrow$  decreasing speed.


$$\therefore v = \sqrt{\frac{5}{(x+2)^2} + \frac{11}{4}}$$

The most common error was to differentiate the given function in terms of  $t$ .

Half a mark was deducted if no statement about sign of  $v$  was included.

✓ (3)

(a)



$FB \perp DF$  (radius  $\perp$  tangent at point of contact)

$DB^2 = FB^2 + DF^2$  (Pythagoras' Theorem)

$\therefore 9^2 = 3^2 + DF^2$

$\therefore DF^2 = 72$

$\therefore DF = 6\sqrt{2}$

$EC \perp DE$  (DC is a diameter  
 $\therefore \angle DEC = 90^\circ$ , angle in a semicircle)

$\therefore \triangle DBF \parallel \triangle DCE$  (equiangular)

as  $\angle D$  is common

$\angle DFB = \angle DEC = 90^\circ$   
 (as indicated above)

$\therefore \frac{DE}{DF} = \frac{DC}{CB}$  (corresponding sides in similar triangles)

$\therefore \frac{DE}{6\sqrt{2}} = \frac{12}{9}$

$\therefore DE = \frac{12 \times 6\sqrt{2}}{9}$

$= 8\sqrt{2}$  (4)

There seemed to be a reluctance to give reasons for geometrical conclusions

(b) (i)  $K! + K \times K!$   
 $= K!(1+K)$   
 $= (K+1)!$  (1)

Done well although some stopped at  $K!(1+K)$

0	0.5	1	Mean
6	18	138	0.91

(ii)  $S(n) \equiv 1 \times 1! + 2 \times 2! + \dots + n \times n! = (n+1)! - 1$

Show  $S(1)$  is true

i.e.  $1 \times 1! = 2! - 1$

LHS = 1

RHS =  $2 - 1$

= 1

$\therefore S(1)$  is true

Assume  $S(k)$  is true

i.e.  $1 \times 1! + 2 \times 2! + \dots + k \times k! = (k+1)! - 1$

Show  $S(k+1)$  is true

i.e.  $1 \times 1! + 2 \times 2! + \dots + k \times k! + (k+1) \times (k+1)!$   
 $= (k+2)! - 1$

LHS =  $(k+1)! - 1 + (k+1) \times (k+1)!$

=  $(k+1)! (1 + k+1) - 1$

=  $(k+1)! (k+2) - 1$

=  $(k+2)! - 1$

= RHS

$\therefore$  If  $S(k)$  is true,  $S(k+1)$  is true

$S(1)$  is true and, if  $S(k)$  is true,

$S(k+1)$  is true

$\therefore$  By the process of Mathematical Induction,  $S(n)$  is true for all integral  $n \geq 1$ . (3)

0	0.5	1	1.5	2	2.5	3	3.5	4	Mean
7	6	11	14	25	8	20	22	49	2.70



Most students demonstrated an understanding of the process of Mathematical Induction. However, many statements were sloppy. For example, "Assume  $n=k$ " rather than "Assume the statement is true if  $n=k$ " or, having defined the statement as  $S(n)$  as above, "Assume that  $S(k)$  is true". Many concluding statements were also sloppy.

$$\begin{aligned}
 &= \frac{9}{2} \left[ \frac{\sin 2\theta}{2} + \theta \right] + C \\
 &= \frac{9}{2} \left[ \sin \theta \cos \theta + \theta \right] + C \\
 &= \frac{9}{2} \left[ \frac{x-3}{3} \sqrt{1 - \frac{(x-3)^2}{9}} + \sin^{-1} \frac{x-3}{3} \right] + C \\
 &= \frac{9}{2} \left[ \frac{x-3}{3} \frac{\sqrt{6x-x^2}}{3} + \sin^{-1} \frac{x-3}{3} \right] + C \\
 &= \frac{1}{2} \left[ (x-3)\sqrt{x(6-x)} + 9 \sin^{-1} \frac{x-3}{3} \right] + C
 \end{aligned}$$

(4)

Many students found their integration challenging. Some left the integral at the form  $\frac{9}{2} [\sin \theta \cos \theta + \theta] + C$ , or equivalent rather than returning to an expression in terms of  $x$ .

0	0.5	1	1.5	2	2.5	3	3.5	4	Mean
3	5	11	4	20	15	25	24	55	2.93

0	0.5	1	1.5	2	2.5	3	Mean
2	8	2	3	0	4	143	2.77

(c) (i) If  $x = 3 + 3 \sin \theta$ ,  
 $dx = 3 \cos \theta$   
 and  $\sin \theta = \frac{x-3}{3}$ .

$$\begin{aligned}
 &\int \sqrt{x(6-x)} \, dx \\
 &= \int \sqrt{(3+3 \sin \theta)(6-3-3 \sin \theta)} \cdot 3 \cos \theta \, d\theta \\
 &= \int 3 \sqrt{(1 - \sin^2 \theta)} \cdot 3 \cos \theta \, d\theta \\
 &= 9 \int \cos^2 \theta \, d\theta
 \end{aligned}$$

NOTE: ORIGINAL INTEGRAL IS POSITIVE

$$= 9 \int \frac{\cos 2\theta + 1}{2} \, d\theta$$

$$\begin{aligned}
 \text{(ii)} \quad V &= \pi \int_0^6 \left( \sqrt{x(6-x)} \right)^2 \, dx \\
 &= \pi \int_0^6 x(6-x) \, dx \\
 &= \pi \times \frac{1}{2} \left[ (x-3)\sqrt{x(6-x)} + 9 \sin^{-1} \left( \frac{x-3}{3} \right) \right]_0^6 \\
 &= \frac{\pi}{2} \left\{ [9 \sin^{-1} 1] - [9 \sin^{-1} (-1)] \right\} \\
 &= \frac{\pi}{2} \left\{ 9 \cdot \frac{\pi}{2} - 9 \left( -\frac{\pi}{2} \right) \right\} \\
 &= \frac{9\pi^2}{2}
 \end{aligned}$$

(3)

Most who progressed through (c) (i) found the appropriate volume.

0	0.5	1	1.5	2	2.5	3	Mean
15	10	21	4	24	23	65	2.05

14(a)

$$\left| \frac{dV}{dt} = \frac{2\pi}{5} \right| \text{ (given)}$$

$$S = \pi r^2$$

$$V = \frac{1}{3} \pi r^2 h.$$

$$\left| \frac{dS}{dr} = 2\pi r \right|$$

$$d \frac{h}{10} = \frac{r}{8} \text{ (SIMILARITY)}$$

$$\therefore V = \frac{5}{12} \pi r^3$$

$$\left| \frac{dV}{dr} = \frac{5}{4} \pi r^2 \right|$$

$$\text{Now } \frac{dS}{dt} = \frac{dS}{dr} \times \frac{dr}{dt} \times \frac{dV}{dt}$$

$$= 2\pi r \times \frac{4}{5\pi r^2} \times \frac{2\pi}{5}$$

$$= \frac{16\pi}{25r}$$

$$\text{When } h=4$$

$$r = \frac{16}{5}$$

$$= \frac{16}{25} \times \pi \times \frac{5}{16}$$

$$= \frac{\pi}{5} \text{ cm/min}$$

COMMENT Not particularly well done.

many students treated  $h$  as a constant in the differentiation

of  $\frac{dV}{dr}$ .

(b) (i) let  $y = 0$ .


$$100 - 5t^2 = 0$$

$$5t^2 = 100$$

$$t^2 = 20$$

$$t = 2\sqrt{5} \text{ sec.}$$

(ii)  $\dot{x} = 30$        $\dot{y} = -10t$   
 $= -20\sqrt{5}$



$$V^2 = 900 + 2000$$

$$= \sqrt{2900}$$

$$V = 10\sqrt{29} \text{ m/s.}$$


$$\theta = \tan^{-1} \frac{2\sqrt{5}}{3}$$

COMMENT


Common error was

to let  $y = -100$ . Generally will done.

(C).



$$\begin{aligned}VM &= 2a \sin 60^\circ \\ &= 2a \frac{\sqrt{3}}{2} \\ \underline{VM} &= \underline{a\sqrt{3}}.\end{aligned}$$



$$\begin{aligned}CM^2 &= (3a)^2 - a^2 \\ &= 8a^2 \\ \therefore \underline{CM} &= \underline{\sqrt{8a^2}}.\end{aligned}$$

$$\begin{aligned}\cos \theta &= \frac{CM^2 + VM^2 - VC^2}{2 \times a\sqrt{3} \times 2\sqrt{2}a} \\ &= \frac{8a^2 + 3a^2 - 5a^2}{4\sqrt{6}a^2} \\ &= \frac{6a^2}{4\sqrt{6}a^2}\end{aligned}$$

$$\therefore \boxed{\cos \theta = \frac{\sqrt{6}}{4}}.$$

COMMENT:

Quite well done.

$$\begin{aligned}(ii) \quad VO &= VM \sin \theta \\ &= a\sqrt{3} \times \frac{\sqrt{10}}{4} \\ &= \frac{a\sqrt{30}}{4}\end{aligned}$$

$$\begin{aligned}\sin^2 \theta &= 1 - \frac{6}{16} \\ &= \frac{10}{16} \\ \therefore \sin \theta &= \frac{\sqrt{10}}{4}\end{aligned}$$

COMMENT:

many students unable to find  $\sin \theta$ .

$$\begin{aligned}
 \text{(III)} \quad \cos \angle B^1 C &= \frac{(2a)^2 + (3a)^2 - (\sqrt{r}a)^2}{2 \times 2a \times 3a} \\
 &= \frac{4a^2 + 9a^2 - 5a^2}{12a^2} \\
 &= \frac{2}{3}
 \end{aligned}$$

$$\begin{aligned}
 \therefore VP^2 &= VB^2 + (ra)^2 - 2 \times 2a \times ra \times \frac{2}{3} \\
 &= 4a^2 + r^2 a^2 - \frac{8ra^2}{3} \\
 &= \frac{a^2}{3} [12 + 3r^2 - 8r]
 \end{aligned}$$

COMMENT Very few students were able to obtain this answer.

The common error was to assume

$\triangle COP \parallel \triangle CMB$ . Hence finding an expression for OP then using Pythagoras to obtain  $VP^2$ . (this was not given marks)

$$\text{(IV)} \quad VP = a \sqrt{\frac{12 + 3r^2 - 8r}{3}}$$

$$\begin{aligned}
 \therefore \sin \theta &= \frac{\frac{a\sqrt{30}}{4}}{a \sqrt{\frac{12 + 3r^2 - 8r}{3}}} \\
 &= \frac{\sqrt{90}}{4 \sqrt{12 + 3r^2 - 8r}} \\
 &= \frac{\sqrt{45}}{\sqrt{8(12 + 3r^2 - 8r)}}
 \end{aligned}$$

COMMENT-

most obtained a mark, allowing for previous error in VO.

(V). Best done by recognizing

that  $\sin \phi$  is maximised  
by  $3r^2 - 8r + 12$  being a minimum  
this occurs when  $6r - 8 = 0$   
 $r = \frac{4}{3}$ .

$$\begin{aligned} \text{ii. } \sin \phi &= \sqrt{\frac{45}{8(3r^2 - 8r + 12)}} \quad \text{when } r = \frac{4}{3} \\ &= \sqrt{\frac{45 \times 9}{8 \times 60}} \\ &= \frac{3\sqrt{6}}{8} \end{aligned}$$

$$\therefore \phi = \sin^{-1} \left( \frac{3\sqrt{6}}{8} \right)$$

COMMENT:

many students were  
able to obtain a mark or two.

many maximised  $\sin \phi$  by  
calculus. Few saw the easier  
approach.