

--	--	--	--	--	--	--	--

CANDIDATE NUMBER

SYDNEY GRAMMAR SCHOOL



2015 Trial Examination

# FORM VI

## MATHEMATICS EXTENSION II

Friday 31st July 2015

### General Instructions

- Reading time — 5 minutes
- Writing time — 3 hour
- Write using black or blue pen.
- Board-approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

### Total — 100 Marks

- All questions may be attempted.

### Section I – 10 Marks

- Questions 1–10 are of equal value.
- Record your solutions to the multiple choice on the sheet provided.

### Section II – 90 Marks

- Questions 11–16 are of equal value.
- All necessary working should be shown.
- Start each question in a new booklet.

### Collection

- Write your candidate number on each answer booklet and on your multiple choice answer sheet.
- Hand in the booklets in a single well-ordered pile.
- Hand in a booklet for each question in Section II, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Write your candidate number on this question paper and hand it in with your answers.
- Place everything inside the answer booklet for Question Eleven.

### Checklist

- SGS booklets — 6 per boy
- Multiple choice answer sheet
- Candidature — 73 boys

Examiner

BDD

**SECTION I - Multiple Choice**

Answers for this section should be recorded on the separate answer sheet handed out with this examination paper.

---

**QUESTION ONE**

The roots of the quadratic equation  $x^2 - 8ix - 20 = 0$  are:

- (A)  $4i \pm 2$  (B)  $4 \pm 2i$   
(C)  $-4i \pm 2$  (D)  $-4 \pm 2i$

**QUESTION TWO**

The value of  $\int_0^{\frac{\pi}{2}} \sin x \cos x dx$  is:

- (A)  $-\frac{1}{2}$  (B)  $\frac{1}{4}$   
(C)  $\frac{1}{2}$  (D) 1

**QUESTION THREE**

The gradient of the tangent to the parametric curve  $x = 2 \sec \theta$ ,  $y = 3 \tan \theta$  at  $\theta$  is:


- (A)  $\frac{2}{3} \sin \theta$  (B)  $\frac{2}{3} \operatorname{cosec} \theta$   
(C)  $\frac{3}{2} \sin \theta$  (D)  $\frac{3}{2} \operatorname{cosec} \theta$

**QUESTION FOUR**

Which of the following functions is odd?

- (A)  $y = x \sin x$  (B)  $y = \sin(\sin(x))$   
(C)  $y = \ln |x|$  (D)  $y = \sin^2(x)$


**QUESTION FIVE**



The size of angle  $\theta$  in the diagram above is:

- (A)  $50^\circ$
- (B)  $55^\circ$
- (C)  $60^\circ$
- (D)  $65^\circ$

**QUESTION SIX**



The point  $P$  in quadrant one represents complex number  $z$ . The points  $O, P, Q, R$  are the vertices of a square, as in the diagram.

Which statement is NOT true about the square:

- (A) side  $OR$  is represented by  $iz$
- (B) the centre of the square is represented by  $\frac{1}{2}(1 - i)z$
- (C) diagonal  $RP$  is represented by  $(1 - i)z$
- (D) vertex  $Q$  is represented by  $(1 + i)z$

**QUESTION SEVEN**


A pupil makes the following claims about the roots of the equation  $z^6 = 1$ :

- (I) The roots lie on the vertices of a hexagon in the complex plane
- (II) The roots lie on the unit circle in the complex plane
- (III) If  $\omega$  is a root, then so is  $\frac{1}{\omega}$
- (IV) If  $\omega$  is a root, then so is  $\bar{\omega}$

Which of these statements are TRUE?

- (A) I and IV
- (B) II and III
- (C) I, II and III
- (D) I, II, III and IV

**QUESTION EIGHT**



The base and top of the solid depicted are right angled isosceles triangles. A pupil is required to determine the volume by slicing parallel to the base. A typical slice parallel to the base at height  $z$  from the base is marked.

The cross-sectional area of the slice is:

- (A)  $\frac{1}{2}(6 - \frac{3}{4}z)^2$
- (B)  $\frac{1}{4}(6 - \frac{1}{4}z)^2$
- (C)  $\frac{1}{2}(7 - z)^2$
- (D)  $\frac{1}{2}(36 - \frac{27}{4}z)$

**QUESTION NINE**

The point defined by the complex number  $z$  moves in the complex plane subject to the constraint  $|z - 3i| + |z + 3i| = 12$ .

The locus of  $z$  is a conic with eccentricity:

(A)  $\frac{1}{4}$ .

(B)  $\frac{1}{2}$ .

(C) 1.

(D) 2.

**QUESTION TEN**

A polynomial  $P(x)$  of fourth degree with real coefficients has the following properties:

$$P(1) = 0, P'(1) \neq 0$$

$$P(2) \neq 0, P'(2) = P''(2) = 0$$

What is the greatest number of complex non-real roots the polynomial could have?

(A) 0

(B) 1

(C) 2

(D) 3

————— End of Section I —————

**SECTION II - Written Response**

Answers for this section should be recorded in the booklets provided.

Show all necessary working.

Start a new booklet for each question.

---

**QUESTION ELEVEN** (15 marks) Use a separate writing booklet. Marks

(a) Given  $w = 1 - 2i$ ,  $z = 3 + 4i$ , express the following in the form  $a + ib$  for real  $a, b$ :

(i)  $w^2$  1

(ii)  $\frac{z}{w}$  1

(b) Given  $t = 1 + i\sqrt{3}$  find:

(i)  $t$  in modulus–argument form, 2

(ii)  $t^8$  in Cartesian form. 1

(c) Find:

(i)  $\int \frac{1}{x^2 + 6x + 13} dx$  2

(ii)  $\int x \sin x dx$  2

(d) Evaluate the following integral, expressing your answer in simplest exact form: 2

$$\int_{10}^{17} \frac{dx}{\sqrt{x^2 - 64}}.$$

(e) (i) Find constants  $A, B$  and  $C$  such that 2

$$\frac{-4x^2 + 5x + 1}{(x - 1)^2} = \frac{A}{(x - 1)^2} + \frac{B}{x - 1} + C.$$

(ii) Hence find 2

$$\int \frac{-4x^2 + 5x + 1}{(x - 1)^2} dx.$$

**QUESTION TWELVE** (15 marks) Use a separate writing booklet.

**Marks**

(a) Consider the hyperbola  $9x^2 - 16y^2 = 144$ .

(i) Find the eccentricity, foci, directrices and asymptotes of the hyperbola. 3

(ii) Sketch the curve, locating the foci, directrices,  $y$ -intercepts and asymptotes. 2


(iii) Use calculus to find the gradient of the tangent at  $x = 5$  in quadrant one. 2

(iv) Consider a tangent with point of contact in quadrant one. Explain geometrically why its gradient will always be greater than 0.75. 1

(b) (i) Solve the equation  $z^5 = -1$ , leaving your answers in modulus-argument form. 1

(ii) Hence factorise  $z^5 + 1$  as a product of real linear and quadratic factors. 2

(c)



A certain solid has a base which is the ellipse  $\frac{x^2}{16} + \frac{y^2}{4} = 1$ . Slices perpendicular to the base and parallel to the  $y$ -axis are right-angled triangles of height 3 units.

(i) Show that the cross-sectional area of a slice parallel to the  $y$ -axis is 2


$$A(x) = \frac{3}{2}\sqrt{16 - x^2}$$

(ii) Hence find the volume of the solid. 2

**QUESTION THIRTEEN** (15 marks) Use a separate writing booklet. **Marks**

- (a) The cubic polynomial  $P(x) = 2x^3 + 15x^2 + 24x + d$  is known to have a repeated real root and a distinct real root. The distinct root and repeated roots have opposite sign. Find the constant  $d$ . **3**

(b)



The curve  $y = e^{-x^2}$  is shown above.

- (i) Use the method of cylindrical shells to find the volume obtained when the region bounded by the axes, the curve and the line  $x = 2$  is rotated about the  $y$ -axis. **3**

- (ii) What is the limiting value as  $N \rightarrow \infty$  of the volume obtained when the region bounded by the axes, the curve and the line  $x = N$  is rotated about the  $y$ -axis? **1**

- (c) A landing aeroplane of mass  $m$  kg is brought to rest by the action of two retarding forces: a force of  $4m$  Newtons due to the reverse thrust of the engines; and a force due to the brakes of  $\frac{mv^2}{40\,000}$  Newtons.

- (i) Show that the aeroplane's equation of motion for its speed  $v$  at time  $t$  seconds after landing is **1**

$$\dot{v} = -\frac{v^2 + 400^2}{40\,000}.$$

- (ii) Assuming the aeroplane lands at a speed of  $U$  m/s, find an expression for the time it takes to come to rest. **3**

- (iii) Show that, given a sufficiently long runway, then no matter how fast its landing speed, it will always come to rest within approximately 2.6 minutes of landing. **1**

- (d) Consider the locus of  $z$  such that  $|z - \sqrt{2} - i| = 1$ .

- (i) Sketch the locus of  $z$  in the complex plane. **1**

- (ii) Find the minimum value of  $|z|$ . **1**


- (iii) Find the maximum value of  $\arg(z)$ , for  $0^\circ < \arg(z) < 90^\circ$ , correct to the nearest degree. **1**



**QUESTION FOURTEEN** (15 marks) Use a separate writing booklet.

**Marks**

(a)



The curve  $y = f(x)$ , sketched above, has asymptotes  $y = 0$  and  $y = -x$ .

Copy or trace the above graph onto three separate number planes. Use your diagrams to show sketches of the following graphs, showing all essential features clearly.

(i)  $y = (f(x))^2$

**2**

(ii)  $|y| = f(x)$

**2**

(iii)  $y = \ln f(x)$

**2**

(b) (i) Let  $z = \text{cis } \theta$ . Use de Moivre's Theorem to prove that for any integer  $n$ ,

**1**

$$z^n - \frac{1}{z^n} = 2i \sin n\theta.$$

(ii) By considering  $\left(z + \frac{1}{z}\right)^5$ , show that  $\sin^5 \theta = \frac{1}{16}(\sin 5\theta - 5 \sin 3\theta + 10 \sin \theta)$ .

**2**

(iii) Solve the following equation for  $0 \leq \theta \leq 2\pi$ :

**2**

$$\sin 5\theta - 5 \sin 3\theta + 9 \sin \theta = 0.$$

(c) You may assume the equation for the chord of contact for the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

from  $(x_0, y_0)$  is  $\frac{x_0 x}{a^2} - \frac{y_0 y}{b^2} = 1$ .

Show that that chord of contact from a point on a directrix is a focal chord.

**2**

(d) Use the substitution  $x = \frac{\pi}{2} - u$  to evaluate


**2**

$$\int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\cos^3 x + \sin^3 x} dx.$$

**QUESTION FIFTEEN** (15 marks) Use a separate writing booklet.

Marks

(a)



Two intersecting circles  $C_1$  and  $C_2$  share a common chord  $AB$ . Points  $P$  and  $H$  lie on circle  $C_1$  and points  $Q$  and  $K$  lie on circle  $C_2$ , such that  $PAQ$  and  $HAK$  are straight. Lines  $HP$  and  $QK$  intersect at  $X$ .

Let  $\angle BKQ = \theta$ .

Copy or trace the diagram into your answer book.

- (i) Find  $\angle BAQ$  in terms of  $\theta$ , giving a reason for your answer. 1
  - (ii) Show that  $BKXH$  is cyclic. 2
  - (iii) Assuming that  $XAB$  is straight, show that  $XAB$  bisects angle  $PBK$ . 2
- (b) (i) List all 10 ways that 3 non-negative integers can add to 3. 1
- (ii) Use the identity  $(1 + x)^{3n} = ((1 + x)^n)^3$  to prove that 2
- $${}^{3n}C_3 = n^3 + 6n \times {}^nC_2 + 3 \times {}^nC_3$$

The question continues over the page

**QUESTION FIFTEEN** (Continued)

- (c) As part of a test of a new capsule delivery system, a capsule of mass  $m$  is fired straight up at speed  $u$  m/s. Air resistance is negligible and the magnitude of the acceleration due to gravity is  $g$ .


The capsule subsequently deploys a parachute and falls back to earth, subject to gravity and to a resistive force of magnitude  $mkv^2$ .

- (i) Use calculus to show that the maximum height attained by the capsule is  $H = \frac{u^2}{2g}$ . 2
- (ii) For the return trip, take the origin at the point it begins falling and assume down is positive. Show that the motion is determined by the equation  $\ddot{x} = k(\alpha^2 - v^2)$ , where  $\alpha^2 = \frac{g}{k}$ . 1
- (iii) Let  $U$  be the impact speed of the package. Find an expression for the square of the speed  $U$  in terms of  $H$ ,  $k$  and  $\alpha$ . 3
- (iv) Assume that the package is launched at speed  $u = \alpha$ . Find the impact speed as a percentage of the launch speed. 1

**QUESTION SIXTEEN** (15 marks) Use a separate writing booklet.

Marks

(a)



An infinite sequence of complex numbers is defined by

$$z_1 = 1, z_{n+1} = \frac{3}{4}iz_n.$$

The path  $z_1z_2z_3 \dots$  defines a piecewise linear spiral in the Argand plane.

- (i) Show that the  $n^{\text{th}}$  edge satisfies the relationship 1

$$z_{n+1} - z_n = \left(\frac{3}{4}i - 1\right) z_n.$$

- (ii) Hence find a simplified expression for the length of the  $n^{\text{th}}$  edge. 1
- (iii) Find the length of the spiral, by considering the limiting sum of the lengths of its edges as  $n \rightarrow \infty$ . 1


- (b) (i) Use algebra to prove, for any integer  $k \geq 0$ , that 1

$$\frac{2k+1}{2k+2} \leq \frac{\sqrt{2k+1}}{\sqrt{2k+3}}.$$

- (ii) Prove, by induction on  $n \geq 0$ , that the central binomial coefficient  $\binom{2n}{n}$  satisfies 3

$$\binom{2n}{n} \leq \frac{4^n}{\sqrt{2n+1}}.$$

(c)



Consider a cubic polynomial  $y = P(x)$  with real coefficients and roots  $0, g \pm hi$  where  $g$  and  $h$  are real and  $h > 0$ . In the diagram above, the roots form the vertices of an isosceles triangle  $OAB$  in the complex plane. The roots of  $P'(x) = 0$  are the foci of the sketched ellipse which touches the triangle at  $g + 0i$ . We have sketched the case  $g > 0$  and with major axis lying on the real axis. The centre of the ellipse is NOT the origin.

- (i) Show that the cubic polynomial has equation  $y = x^3 - 2gx^2 + (g^2 + h^2)x$ . 1  
 (ii) Show that the turning points of  $y = P(x)$ , and hence the foci of the ellipse, occur at 1

$$x = \frac{2}{3}g \pm \frac{1}{3}\sqrt{g^2 - 3h^2}.$$

- (iii) Find the condition on  $g$  and  $h$  and hence on  $\angle AOB$  which ensures that the major axis of this ellipse lies on the real axis. You may assume this condition holds in parts (iv) and (v). 1  
 (iv) Find the equation of the ellipse. 2  
 (v) Show that the ellipse is tangential to the triangle at the midpoints of  $OA$  and  $OB$ . 2  
 (vi) If the triangle is equilateral, describe the behaviour of the polynomial  $P(x)$  at the centre of the ellipse, for real  $x$ . 1

————— End of Section II —————

**END OF EXAMINATION**

**SECTION I - Multiple Choice****QUESTION ONE**

The discriminant  $\Delta = (-8i)^2 - 4 \times 1 \times (-20) = 16 = (4)^2$ . Hence the roots are

$$\frac{8i \pm 4}{2} = 4i \pm 2$$

Hence **A**.

**QUESTION TWO**

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \sin x \cos x \, dx &= \frac{1}{2} \int_0^{\frac{\pi}{2}} \sin 2x \, dx \\ &= \left[ -\frac{1}{4} \cos 2x \right]_0^{\frac{\pi}{2}} \\ &= \frac{1}{4} ((-\cos \pi) + \cos 0) \\ &= \frac{1}{2} \end{aligned}$$

Hence **C**.

**QUESTION THREE**

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy/d\theta}{dx/d\theta} \\ &= 3 \sec^2 \theta \div 2 \sec \theta \tan \theta \\ &= \frac{3}{2} \sec \theta \div \tan \theta \\ &= \frac{3}{2} \frac{1}{\cos \theta} \frac{\cos \theta}{\sin \theta} \\ &= \frac{3}{2} \operatorname{cosec} \theta \end{aligned}$$

Hence **D**.

**QUESTION FOUR**

Options A, C and D are even. Only option B is odd. Hence **B**.

**QUESTION FIVE**

Angle  $XDY = \theta + 20$  (Exterior opposite angle in  $\triangle AXD$ )

Angle  $BCD = \theta + 20 + 30$  (Exterior opposite angle in  $\triangle DCY$ )

Thus  $\theta + (\theta + 50) = 180$  opposite angles of cyclic quad  $ABCD$ )

So  $\theta = 65$ .

Hence **D**.

**QUESTION SIX**

Option **C** is incorrect.

**QUESTION SEVEN**

All statements are TRUE. Hence **D**.

**QUESTION EIGHT**

The linear equation  $y = (6 - \frac{3}{4}z)$  satisfies the conditions  $y = 6$  when  $z = 0$  and  $y = 3$  when  $z = 4$ . The area of the triangle is  $\frac{1}{2}y^2 = \frac{1}{2}(6 - \frac{3}{4})^2$ , hence the correct answer is **A**.

**QUESTION NINE**

The foci are  $\pm 3i$ , hence the distance between the two foci is  $2ae = 6$ . But the sum of the distances from a point on the ellipse to the foci is  $2a = 12$ . Combining these two equations,  $e = \frac{1}{2}$ .

Hence the correct answer is **B**.

**QUESTION TEN**

The correct answer is 2. Note that there must be an even number of complex non-real roots, because of the real coefficients, and two is a possible answer. This is easily seen by drawing a polynomial with zero when  $x = 1$ , stationary point of inflexion when  $x = 2$  and a turning point at some larger  $x$  value.

**SECTION II - Written Response**

**QUESTION ELEVEN**

$$\begin{aligned} \text{(a) (i) } w^2 &= (1 - 2i)^2 \\ &= 1 - 4 - 4i \\ &= -3 - 4i \end{aligned}$$

$$\begin{aligned} \text{(ii) } \frac{z}{w} &= \frac{3 + 4i}{1 - 2i} \\ &= \frac{(3 + 4i)(1 + 2i)}{1 + 4} \\ &= \frac{3 - 8 + 4i + 6i}{5} \\ &= -1 + 2i \end{aligned}$$

$$\text{(b) (i) We have } |t| = 2, \arg(t) = \tan^{-1}(\sqrt{3}) = \frac{\pi}{3}.$$

$$\text{Hence } t = 2 \operatorname{cis} \frac{\pi}{3}.$$

$$\begin{aligned} \text{(ii) } t^8 &= 2^8 \operatorname{cis} \frac{8\pi}{3} \\ &= 256 \operatorname{cis} \frac{2\pi}{3} \\ &= 256\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) \\ &= 128(-1 + i\sqrt{3}) \end{aligned}$$

$$\begin{aligned} \text{(c) (i) } \int \frac{1}{x^2 + 6x + 13} dx &= \int \frac{1}{(x + 3)^2 + 2^2} dx \\ &= \frac{1}{2} \tan^{-1} \frac{x + 3}{2} + C \end{aligned}$$

$$\begin{aligned} \text{(ii) } \int x \sin x dx &= x(-\cos x) - \int 1 \times (-\cos x) dx \\ &= -x \cos x + \sin x + C \end{aligned}$$

$$\begin{aligned} \text{(d) } \int_{10}^{17} \frac{dx}{\sqrt{x^2 - 64}} &= \left[ \ln(x + \sqrt{x^2 - 64}) \right]_{10}^{17} \\ &= \ln(17 + \sqrt{17^2 - 64}) - \ln(10 + \sqrt{10^2 - 64}) \\ &= \ln 32 - \ln 16 \\ &= \ln 2 \end{aligned}$$

(e)

$$\begin{aligned} \text{(i) } \frac{-4x^2 + 5x + 1}{(x - 1)^2} &= \frac{A}{(x - 1)^2} + \frac{B}{x - 1} + C \\ -4x^2 + 5x + 1 &= A + B(x - 1) + C(x - 1)^2 \end{aligned}$$

Equating coefficients of  $x^2$  tells us  $C = -4$ .  
Substituting  $x = 1$  tells us  $A = 2$ .

Substituting  $x = 0$ , tells us:

$$A - B + C = 1$$

$$2 - B + 4 = 1$$

$$B = -3$$

$$\begin{aligned} \text{(ii) Hence } \int \frac{-4x^2 + 5x + 1}{(x-1)^2} dx &= \int \frac{2}{(x-1)^2} dx + \int \frac{-3}{(x-1)} dx - \int 4 dx \\ &= \frac{-2}{(x-1)} - 3 \ln |x-1| - 4x + C \end{aligned}$$

### QUESTION TWELVE

(a)

$$\text{(i) } \frac{x^2}{16} - \frac{y^2}{9} = 1$$

Hence  $a = 4$  and  $b = 3$ .

$$\text{Now } b^2 = a^2(e^2 - 1)$$

$$e^2 = \frac{9}{16} + 1$$


$$e^2 = \frac{25}{16}$$

$$e = \frac{5}{4}$$

The foci are  $(\pm ae, 0) = (\pm 5, 0)$ . The directrices are  $x = \frac{a}{e}$  and  $x = -\frac{a}{e}$ . Thus  $x = \frac{16}{5}$  and  $x = -\frac{16}{5}$ .

The asymptotes are  $y = \frac{b}{a}x$  and  $y = -\frac{b}{a}x$ . Thus  $y = \frac{3}{4}x$  and  $y = -\frac{3}{4}x$ .

(ii)



(iii) By substituting in the equation for the hyperbola, when  $x = 5$ , we find  $y = \frac{9}{4}$  in quadrant one. Differentiating with respect to  $x$ :



$$\begin{aligned}
 18x - 32y \frac{dy}{dx} &= 0 \\
 \frac{dy}{dx} &= \frac{9x}{16y} \\
 &= \frac{45}{36} \quad \text{when } x = 5 \\
 &= 1.25
 \end{aligned}$$

(iv) The tangent in Quadrant One will be steeper than the asymptote, thus its gradient will never be less than that of the asymptote.

(b) (i) Let  $z = \text{cis } \theta$ . Then

$$z^5 = -1$$

$$\text{cis } 5\theta = \text{cis}(\pi + 2k\pi) \quad \text{for any integer } k$$

Equating arguments gives:

$$5\theta = (2k + 1)\pi$$

$$\theta = \frac{\pi}{5}, \frac{3\pi}{5}, \pi, -\frac{\pi}{5}, -\frac{3\pi}{5}$$

Hence (in conjugate pairs) the roots are:

$$z = -1, \quad z = \text{cis}\left(\pm\frac{1}{5}\pi\right), \quad z = \text{cis}\left(\pm\frac{3}{5}\pi\right)$$

(ii) Grouping the conjugate roots, we get:

$$\begin{aligned}
 z^5 + 1 &= (z + 1) \times \left(z - \text{cis}\frac{1}{5}\pi\right) \left(z - \text{cis}\left(-\frac{1}{5}\pi\right)\right) \times \left(z - \text{cis}\frac{3}{5}\pi\right) \left(z - \text{cis}\left(-\frac{3}{5}\pi\right)\right) \\
 &= (z + 1) \left(z^2 - 2\cos\left(\frac{1}{5}\pi\right)z + 1\right) \left(z^2 - 2\cos\left(\frac{3}{5}\pi\right)z + 1\right)
 \end{aligned}$$

(c) (i) The area of the triangle is:

$$\begin{aligned}
 \frac{1}{2}bh &= \frac{1}{2}(2y) \times 3 \\
 &= 3y \\
 &= 6 \times \sqrt{1 - \frac{x^2}{16}} \\
 &= \frac{6}{4} \times \sqrt{16 - x^2} \\
 &= \frac{3}{2} \sqrt{16 - x^2}
 \end{aligned}$$

(ii) The volume is

$$\begin{aligned}
 V &= 3 \times \int_0^4 \sqrt{16 - x^2} \, dx \\
 &= 3 \times \frac{1}{4} \pi 4^2
 \end{aligned}$$

since the integral is the area of a quarter circle of radius 4.

Thus the volume is  $V = 12\pi$ .

(This integral can also be evaluated using the trig substitution  $x = 4 \sin u$ .)

**QUESTION THIRTEEN**

(a) At a double root we have a zero of the derivative.

$$P'(x) = 6x^2 + 30x + 24$$

$$= 6(x + 4)(x + 1)$$

Hence the possibilities are  $x = -1$  or  $x = -4$ . The other root must be positive and the product of roots must be negative, i.e.  $d < 0$ .

If  $x = -1$  then  $P(-1) = 2(-1)^3 + 15(-1)^2 + 24(-1) + d$

$$0 = -2 + 15 - 24 + d$$

$$d = 11$$

If  $x = -4$  then  $P(-4) = 2(-4)^3 + 15(-4)^2 + 24(-4) + d$


$$0 = -128 + 240 - 96 + d$$

$$d = -16$$

Hence  $d = -16$ .

This question can also be solved using sum and product of roots methods.

(b) (i)



The volume of the cylindrical shell  $dV = 2\pi xy dx$ . Total volume is

$$V = \int_0^2 2\pi xy dx$$

$$= \pi \int_0^2 2xe^{-x^2} dx$$

$$= \pi \times [e^{-x^2}]_0^2$$

$$= \pi \times (1 - e^{-4})$$

(ii)  $V = \pi \times \lim_{N \rightarrow \infty} (1 - e^{-N^2})$

$$= \pi$$

(c) (i)  $m\dot{v} = \frac{-mv^2}{40\,000} - 4m$

$$\dot{v} = -\frac{v^2}{40\,000} - 4$$


$$\dot{v} = -\frac{v^2 + 160\,000}{40\,000}$$

$$= -\frac{v^2 + 400^2}{40\,000}$$

$$\begin{aligned}
 \text{(ii)} \quad \frac{dv}{dt} &= -\frac{v^2 + 400^2}{40\,000} \\
 dt &= \frac{-40\,000\,dv}{v^2 + 400^2} \\
 \int_0^T dt &= -40\,000 \times \int_U^0 \frac{dv}{v^2 + 400^2} \\
 T &= 40\,000 \times \frac{1}{400} \tan^{-1} \frac{U}{400} \\
 T &= 100 \tan^{-1} \frac{U}{400}
 \end{aligned}$$

(iii) As  $U \rightarrow \infty$ ,  $T \rightarrow 100 \times \frac{\pi}{2}$  seconds, which is about 2.6 minutes.

(d) (i)




(ii) The point  $z$  of minimum modulus is the point on the circle closest to the origin. This distance is:

$$(\text{distance from origin to centre of circle}) - (\text{radius}) = \sqrt{3} - 1$$


(iii) The point with maximum  $\arg(z)$  on the circle is defined by the tangent to the circle. The argument is  $2 \times \tan^{-1} \frac{1}{\sqrt{2}} \doteq 71^\circ$ .

**QUESTION FOURTEEN**


(a) (i)



(ii)



(iii)



(b) (i) Let  $z = \text{cis } \theta$ . Then by de Moivre's Theorem,

$$\begin{aligned} z^n - \frac{1}{z^n} &= (\text{cis } \theta)^n - (\text{cis } \theta)^{-n} \\ &= \text{cis}(n\theta) - \text{cis}(-n\theta) \\ &= \cos(n\theta) + i \sin(n\theta) - \cos(-n\theta) - i \sin(-n\theta) \\ &= \cos(n\theta) + i \sin(n\theta) - \cos(n\theta) + i \sin(n\theta) \\ &= 2i \sin(n\theta) \end{aligned}$$

Where we have used the evenness of the cosine function and the oddness of the sine function.

$$\begin{aligned} \text{(ii)} \quad \left(z - \frac{1}{z}\right)^5 &= z^5 - 5z^4 \frac{1}{z} + 10z^3 \frac{1}{z^2} - 10z^2 \frac{1}{z^3} + 5z \frac{1}{z^4} - \frac{1}{z^5} \\ &= \left(z^5 - \frac{1}{z^5}\right) - 5\left(z^3 - \frac{1}{z^3}\right) + 10\left(z - \frac{1}{z}\right) \\ &= 2i \sin 5\theta - 5 \times 2i \sin 3\theta + 10 \times 2i \sin \theta \end{aligned}$$

Now the LHS of this expression is  $(2i \sin \theta)^5$ , hence

$$32i \sin^5 \theta = 2i \sin 5\theta - 5 \times 2i \sin 3\theta + 10 \times 2i \sin \theta$$

$$16 \sin^5 \theta = \sin 5\theta - 5 \sin 3\theta + 10 \sin \theta$$

$$\sin^5 \theta = \frac{1}{16} \left( \sin 5\theta - 5 \sin 3\theta + 10 \sin \theta \right)$$

(iii) From this equation  $16 \sin^5 \theta - 10 \sin \theta = \sin 5\theta - 5 \sin 3\theta$ .

$$\text{Hence} \quad \sin 5\theta - 5 \sin 3\theta + 9 \sin \theta = 0$$

$$\text{Becomes } 16 \sin^5 \theta - 10 \sin \theta + 9 \sin \theta = 0$$

$$16 \sin^5 \theta - \sin \theta = 0$$

$$\sin \theta (16 \sin^4 \theta - 1) = 0.$$

$$\text{So } \sin \theta = 0 \text{ or } \sin \theta = \pm \frac{1}{2}.$$

The solutions of these equations in the given domain are:

$$\theta = 0, \pi, 2\pi, \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$$

(c) Let the point be  $(x_0, y_0) = \left(\frac{a}{e}, y_0\right)$ . The chord of contact is  $\frac{x}{a^2} \left(\frac{a}{e}\right) - \frac{y_0 y}{b^2} = 1$ .

Is  $(ae, 0)$  on this chord?

$$LHS = \frac{x}{a^2} \left(\frac{a}{e}\right) - \frac{y_0 y}{b^2}$$

$$= 1 - 0$$

$$= RHS$$

So yes, the chord passes through the focus.

$$\begin{aligned}
 \text{(d)} \quad \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\cos^3 x + \sin^3 x} dx &= \int_{\frac{\pi}{2}}^0 \frac{\cos^3(\frac{\pi}{2} - u)}{\cos^3(\frac{\pi}{2} - u) + \sin^3(\frac{\pi}{2} - u)} (-dx) \\
 &= \int_0^{\frac{\pi}{2}} \frac{\sin^3 u}{\sin^3 u + \cos^3 u} du \\
 &= \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx \quad (\text{relabelling } u \text{ as } x)
 \end{aligned}$$

$$\begin{aligned}
 \text{Hence } 2 \times \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\cos^3 x + \sin^3 x} dx &= \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\cos^3 x + \sin^3 x} dx + \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx \\
 &= \int_0^{\frac{\pi}{2}} \frac{\cos^3 x + \sin^3 x}{\cos^3 x + \sin^3 x} dx \\
 &= \int_0^{\frac{\pi}{2}} 1 dx \\
 &= \frac{\pi}{2}
 \end{aligned}$$

$$\text{Thus } \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\cos^3 x + \sin^3 x} dx = \frac{\pi}{4}.$$

### QUESTION FIFTEEN

- (a) (i)  $\angle BAQ = \theta$  (angles at the circumference on arc  $BQ$ )  
 (ii) Hence  $\angle BHP = \theta$  (interior opposite angle in cyclic quadrilateral  $BHPA$ )  
 Since  $\angle BHP = \angle BHX = \theta$  and the exterior opposite angle  $\angle BKQ = \theta$ , we have that quadrilateral  $BKXH$  is cyclic.  
 (iii)  $\angle PBA = \angle PHA$  (angles at the circumference on arc  $PA$  in circle  $\mathcal{C}_1$ )  
 $= \angle XHK$  (same angle)  
 $= \angle XBK$  (angles at the circumference on arc  $XK$  in circle  $BKXH$ )  
 $= \angle ABK$  (same angle)

- (b) (i) In any order, the ways three integers can add to 3 are:

$$\begin{aligned}
 &3 + 0 + 0, 0 + 3 + 0, 0 + 0 + 3 \\
 &1 + 2 + 0, 2 + 1 + 0, 0 + 2 + 1, 0 + 1 + 2, 1 + 0 + 2, 2 + 0 + 1 \\
 &1 + 1 + 1
 \end{aligned}$$

- (ii) We look to be equating coefficients of  $x^3$ .

$$\begin{aligned}
 LHS &= (1 + x)^{3n} \\
 &= {}^{3n}C_0 x^0 + {}^{3n}C_1 x^1 + \dots
 \end{aligned}$$

... and the coefficient of  $x^3$  is  ${}^{3n}C_3$ .

The  $RHS$  is  $(1 + x)^n (1 + x)^n (1 + x)^n$ .

We need to consider how we can get an  $x^3$  term when we expand the brackets. Part (i) gives us a hint here, since the sum of the three indices (one from each bracket) must be 3.

Method 1 (list them all): The coefficient of  $x^3$  is:

$$\begin{aligned} & {}^n C_3 {}^n C_0 {}^n C_0 + {}^n C_0 {}^n C_3 {}^n C_0 + {}^n C_0 {}^n C_0 {}^n C_3 \\ & + {}^n C_1 {}^n C_2 {}^n C_0 + {}^n C_2 {}^n C_1 {}^n C_0 + {}^n C_0 {}^n C_2 {}^n C_1 + {}^n C_0 {}^n C_1 {}^n C_2 + {}^n C_1 {}^n C_0 {}^n C_2 \\ & + {}^n C_2 {}^n C_0 {}^n C_1 \\ & + {}^n C_1 {}^n C_1 {}^n C_1 \\ & = 3 \times {}^n C_3 + 6 \times {}^n C_1 {}^n C_2 + ({}^n C_1)^3 \end{aligned}$$

Method 2 (Avoid listing them all):

- There are 3 ways to get  $x^3$  from one bracket,  $x^0$  from each of the others.  
This gives a contribution  $3 \times {}^n C_3 x^3 \times x^0 \times x^0$
- There are 6 ways to get  $x^1$  from one bracket,  $x^2$  from a second bracket and  $x^0$  from a third.  
This gives a contribution  $6 \times {}^n C_1 x^1 \times {}^n C_2 x^2 \times x^0$
- There is 1 way to get  $x^1$  from all of the brackets  
This gives a contribution  $1 \times {}^n C_1 x \times {}^n C_1 x \times {}^n C_1 x$

Thus from the *RHS* the coefficient of  $x^3$  is:

$$3 \times {}^n C_3 + 6 \times {}^n C_1 {}^n C_2 + ({}^n C_1)^3$$

Thus either method gives us

$$\begin{aligned} & 3 \times {}^n C_3 + 6 \times {}^n C_1 {}^n C_2 + ({}^n C_1)^3 \\ & = 3 \times {}^n C_3 + 6n \times {}^n C_2 + n^3 \end{aligned}$$

Equating coefficients of  $x^3$  yields the required result.

- (c) (i) Starting from the equation of motion  $\ddot{x} = -g$ , we get

$$\begin{aligned} \ddot{x} &= -g \\ v \frac{dv}{dx} &= -g \\ \int_u^0 v \, dv &= - \int_0^H g \, dx \\ \left[ \frac{1}{2} v^2 \right]_u^0 &= \left[ -gx \right]_0^H \\ -gH &= -\frac{1}{2} u^2 \\ H &= \frac{u^2}{2g} \end{aligned}$$

- (ii) Working the equations with down as positive and including the resistive term;

$$\begin{aligned} m\ddot{x} &= mg - kv^2 \\ \ddot{x} &= g - kv^2 \\ &= k \left( \frac{g}{k} - v^2 \right) \\ &= k(\alpha^2 - v^2) \end{aligned}$$

where  $\alpha^2 = \frac{g}{k}$ .

(iii) We need to integrate the equation of motion:

$$v \frac{dv}{dx} = k(\alpha^2 - v^2)$$

$$-\frac{1}{2} \times \int_0^U \frac{-2v dv}{\alpha^2 - v^2} = k \times \int_0^H dx$$

$$\left[-\frac{1}{2} \ln(\alpha^2 - v^2)\right]_0^U = [kx]_0^H$$

$$-\frac{1}{2} \ln \frac{\alpha^2 - U^2}{\alpha^2} = kH$$

Hence  $kH = -\frac{1}{2} \ln \frac{\alpha^2 - U^2}{\alpha^2}$ . (Note that  $U < \alpha$ , the terminal velocity.)

Rearranging we get:

$$U^2 = \alpha^2(1 - e^{-2kH})$$

(iv) We have:

$$\frac{(\text{impact speed})^2}{(\text{launch speed})^2} = \frac{\alpha^2(1 - e^{-2kH})}{\alpha^2}$$

$$= 1 - e^{-2kH}$$

But  $2kH = 2 \times \frac{g}{\alpha^2} \times \frac{\alpha^2}{2g} = 1$ . Hence

$$\frac{(\text{impact speed})^2}{(\text{launch speed})^2} = 1 - e^{-1}$$

$$= \frac{e - 1}{e}$$

$$\doteq 0.632$$

So  $\frac{\text{impact speed}}{\text{launch speed}} = 79.5\%$

That is, the impact speed = 79.5% of the launch speed.

## QUESTION SIXTEEN

(a) (i)  $z_{n+1} - z_n = \frac{3}{4}iz_n - z_n$

$$= \left(\frac{3}{4}i - 1\right)z_n$$

(ii)  $|z_{n+1} - z_n| = \left|\frac{3}{4}i - 1\right| \times |z_n|$

$$= \frac{5}{4}|z_n|$$

But  $|z_n| = \left(\frac{3}{4}\right)^{n-1}$ , so

(iii)  $|z_{n+1} - z_n| = \frac{5}{4} \times \left(\frac{3}{4}\right)^{n-1}$

$$\text{Total length} = \frac{5}{4} \left(1 + \frac{3}{4} + \frac{9}{16} + \dots\right)$$

$$= \frac{5}{4} \times \frac{1}{1 - \frac{3}{4}}$$

$$= \frac{5}{4} \times 4$$

$$= 5$$



(b) Consider  $RHS/LHS$ . We want to show that this ratio is greater than 1.

$$\begin{aligned} \left(\frac{RHS}{LHS}\right)^2 &= \frac{(2k+1)(2k+2)^2}{(2k+3)(2k+1)^2} \\ &= \frac{(2k+2)^2}{(2k+3)(2k+1)} \\ &= \frac{4k^2+8k+4}{4k^2+8k+3} \\ &> 1 \end{aligned}$$

(Since the numerator is greater than the denominator.)

(i)

Step A: Let us check the result for  $n = 0$ . When  $n = 0$ :

$$\begin{aligned} LHS &= \binom{0}{0} & RHS &= \frac{4^n}{\sqrt{2n+1}} \\ &= 1 & &= 1 \end{aligned}$$

Step B: Assume the results holds for  $n = k$ , that is assume

$$\binom{2k}{k} \leq \frac{4^k}{\sqrt{2k+1}}$$

We need to show that the result holds for  $n = k + 1$ , that is to show that

$$\binom{2k+2}{k+1} \leq \frac{4^{k+1}}{\sqrt{2k+3}}$$

$$\begin{aligned} LHS &= \frac{(2k+2)!}{(k+1)!(k+1)!} \\ &= \frac{(2k)!}{k!k!} \times \frac{(2k+1)(2k+2)}{(k+1)(k+1)} \\ &\leq \frac{4^k}{\sqrt{2k+1}} \times \frac{4(2k+1)}{(2k+2)} \\ &\leq \frac{4^{k+1}}{\sqrt{2k+1}} \times \frac{\sqrt{2k+1}}{\sqrt{2k+3}} \\ &\leq \frac{4^{k+1}}{\sqrt{2k+3}} \end{aligned}$$

as required.

Step C: Hence the result holds for all  $n$  by the Principle of Mathematical Induction.

(c) (i)  $y = x(x - (g + hi))(x - (g - hi))$   
 $= x(x^2 - 2gx + (g^2 + h^2))$   
 $= x^3 - 2gx^2 + x(g^2 + h^2)$

(ii)  $y' = 3x^2 - 4gx + (g^2 + h^2)$

So  $y' = 0$  when

$$\begin{aligned} x &= \frac{4g \pm \sqrt{16g^2 - 4 \times 3 \times (g^2 + h^2)}}{6} \\ &= \frac{2}{3}g \pm \frac{1}{3}\sqrt{g^2 - 3h^2} \end{aligned}$$

(iii) We need  $g^2 - 3h^2 \geq 0$ , so that the foci are real.

$$\text{Thus } \frac{h^2}{g^2} \leq \frac{1}{3}$$

$$\frac{h}{g} \leq \frac{1}{\sqrt{3}}$$

But  $\tan \frac{1}{2} \angle AOB = \frac{h}{g}$ , so  $\frac{1}{2} \angle AOB \leq 30^\circ$ . Thus  $\angle AOB \leq 60^\circ$ .

(iv) The centre of the ellipse occurs at the midpoint of the foci, which are the stationary points of  $P(x)$ . Thus the centre is at  $x = \frac{2}{3}g$ .

The equation of the ellipse is

$$\frac{(x - \frac{2}{3}g)^2}{a^2} + \frac{y^2}{b^2} = 1$$

The endpoint of the ellipse is given to be  $(g, 0)$ , so that  $a = g - \frac{2}{3}g = \frac{1}{3}g$ . The distance between the foci is  $2ae$ , so using part (i)

$$2ae = \frac{2}{3}\sqrt{g^2 - 3h^2}$$

$$\begin{aligned} \text{Thus } b^2 &= a^2(1 - e^2) \\ &= a^2 - (ae)^2 \\ &= \frac{1}{9}g^2 - \frac{1}{4} \times \frac{4}{9}(g^2 - 3h^2) \\ &= \frac{1}{3}h^2 \end{aligned}$$

Hence the equation is

$$\frac{(x - \frac{2}{3}g)^2}{\frac{1}{9}g^2} + \frac{y^2}{\frac{1}{3}h^2} = 1$$

(v) We need to show that  $(\frac{1}{2}g, \frac{1}{2}h)$  lies on the ellipse, and that at this point the ellipse has gradient  $\frac{h}{g}$ .

Substituting in the equation for the ellipse:

$$\begin{aligned} LHS &= \frac{(\frac{1}{2}g - \frac{2}{3}g)^2}{\frac{1}{9}g^2} + \frac{(\frac{1}{2}h)^2}{\frac{1}{3}h^2} \\ &= \frac{\frac{1}{36}g^2}{\frac{1}{9}g^2} + \frac{3}{4} \\ &= \frac{1}{4} + \frac{3}{4} \\ &= 1 \end{aligned}$$

Hence the point lies on the ellipse.

By implicit differentiation of the equation of the ellipse:

$$\frac{2(x - \frac{2}{3}g)}{\frac{1}{9}g^2} + \frac{2yy'}{\frac{1}{3}h^2} = 0$$

$$\begin{aligned}
 \text{Thus } y' &= \frac{-2(x - \frac{2}{3}g)}{\frac{1}{9}g^2} \times \frac{\frac{1}{3}h^2}{2y} \\
 &= \frac{-2(\frac{1}{2}g - \frac{2}{3}g)}{\frac{1}{9}g^2} \times \frac{\frac{1}{3}h^2}{2\frac{1}{2}h} \\
 &= \frac{-2(-\frac{1}{6}g)}{\frac{1}{9}g^2} \times \frac{1}{3}h \\
 &= \frac{3}{g} \times \frac{1}{3}h \\
 &= \frac{h}{g}
 \end{aligned}$$

Which is the gradient of  $OA$ , and hence the ellipse is tangential to the triangle at the midpoint of  $OA$ .

A similar proof (not required) would show that the ellipse is tangential to the triangle at the midpoint of  $OB$  also.

- (vi) Angle  $AOB = 60^\circ$ , hence  $\frac{h}{g} = \tan 30^\circ = \frac{1}{\sqrt{3}}$ . Thus  $g^2 - 3h^2 = 0$ . The foci are both  $x = \frac{2}{3}g$  and the ellipse is a circle, with centre  $(\frac{2}{3}g, 0)$  and radius  $\frac{1}{3}g$ .

Note: At  $x = \frac{2}{3}g$  the polynomial has a double root of  $P'(x)$  and a point of inflexion – it defines a stationary point of inflexion.