

--	--	--	--	--	--	--	--	--	--

Student Number

**2016**

TRIAL HIGHER SCHOOL CERTIFICATE  
EXAMINATION

# Extension 1 Mathematics

\*\*\*\*\*

**General Instructions**

- Reading time – 5 minutes
- Working time - 2 hours
- Write using blue or black pen  
Black pen is preferred
- Approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11-14 show relevant mathematical reasoning and/or calculations
- Start a new booklet for each question

**Total Marks – 70**

**Section I** - Pages \*\*\*

**10 marks**

- Attempt Questions 1 – 10
- Allow about 15 minutes for this section

**Section II** - Pages \*\*\*\*

**60 marks**

- Attempt Questions 11 – 14
- Allow about 1 hour and 45 minutes for this section

Question	Marks
<b>1 - 10</b>	<b>/10</b>
<b>11</b>	<b>/15</b>
<b>12</b>	<b>/15</b>
<b>13</b>	<b>/15</b>
<b>14</b>	<b>/15</b>
<b>Total</b>	<b>/70</b>

***THIS QUESTION PAPER MUST NOT BE REMOVED FROM THE EXAMINATION ROOM***

*This assessment task constitutes 40% of the Higher School Certificate Course Assessment*

## Section I

10 marks

Attempt Questions 1 – 10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for questions 1 – 10 (Detach from paper)


---

1)	<p>In how many ways can 8 students be arranged if the tallest is first and the shortest is last?</p> <p>(A) <math>{}^8C_6</math></p> <p>(B) <math>{}^6C_6</math></p> <p>(C) <math>{}^8P_6</math></p> <p>(D) <math>{}^6P_6</math></p>
2)	<p>If <math>x = 1.6</math> is a close root of the equation <math>x^3 - 4x + 2 = 0</math>, find a better approximation to two decimal places.</p> <p>(A) 1.68</p> <p>(B) 13.71</p> <p>(C) 0.43</p> <p>(D) 4.96</p>
3)	<p>Evaluate <math>\sin^{-1}\left(-\frac{1}{2}\right)</math> as an exact answer.</p> <p>(A) <math>\frac{\pi}{6}</math></p> <p>(B) <math>\frac{5\pi}{6}</math></p> <p>(C) <math>-\frac{\pi}{6}</math></p> <p>(D) <math>-\frac{5\pi}{6}</math></p>

4) A cone has a base diameter of 16 cm and a perpendicular height of 12 cm. The angle the side of the cone makes with its base is:


(A)  $56^\circ$   
 (B)  $37^\circ$   
 (C)  $34^\circ$   
 (D)  $53^\circ$

5) Given  $\angle TPS = 94^\circ$ ,  $\angle RQU = 92^\circ$ ,  $\angle QSR = 48^\circ$ , find  $\angle SVR$







(A)  $82^\circ$   
 (B)  $92^\circ$   
 (C)  $94^\circ$   
 (D)  $96^\circ$

6)



The diagram shows a sketch of the curve  $y = \sin^2 x$  between  $x=0$  and  $x=2\pi$ . The shaded area equals:

(A)  $2\pi$  square units  
 (B)  $3\pi$  square units  
 (C)  $2(2\pi - 1)$  square units  
 (D)  $4\pi$  square units

7)	<p>The equation <math>\cos(2x) = \operatorname{cosec}\left(x - \frac{\pi}{2}\right)</math> <math>-\pi \leq x \leq \pi</math>, has how many solutions?</p> <p>(A) 0            (B) 1            (C) 2            (D) 3</p>
8)	<p>Which of the following is the graph of <math>\sin(\sin^{-1}x)</math>?</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> <p>(A)</p>  </div> <div style="text-align: center;"> <p>(B)</p>  </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;"> <p>(C)</p>  </div> <div style="text-align: center;"> <p>(D)</p>  </div> </div>
9)	<p>Using the substitution <math>u = \sqrt{x}</math>, <math>\int \frac{dx}{x+\sqrt{x}}</math> can be transformed to:</p> <p>(A) <math>\int \frac{2du}{u+1}</math>            (B) <math>\int \frac{du}{u^2+u}</math>            (C) <math>\int \frac{2du}{u^2+u}</math>            (D) <math>\frac{1}{2} \int \frac{du}{u^2+u}</math></p>

<p><b>10)</b></p>	<p>The speed <math>v</math> m/s of a point moving along the <math>x</math> axis is given by <math>v^2 = 36 + 6x - 2x^2</math>, where <math>x</math> is in meters.</p> <p>The period and amplitude of the motion are:</p> <p>(A) Period <math>\pi</math> and amplitude <math>\frac{9}{2}</math></p> <p>(B) Period <math>2\pi</math> and amplitude <math>\frac{\sqrt{63}}{2}</math></p> <p>(C) Period <math>\sqrt{2}\pi</math> and amplitude <math>\frac{9}{2}</math></p> <p>(D) Period <math>\sqrt{2}\pi</math> and amplitude <math>\frac{3}{2}</math></p>
<p><b>11)</b></p>	<p>A particle undergoes linear acceleration according to the equation <math>a = (x + 4)^3 \text{ m/s}^2</math>. Given that the particle commences motion at the origin with a velocity <math>4 \text{ m/s}</math>, what is the particle's displacement when <math>v = 10 \text{ m/s}</math>, given that <math>x &lt; 0</math>?</p> <p>(A) <math>-0.1313</math></p> <p>(B) <math>-7.7606</math></p> <p>(C) <math>-8.5378</math></p> <p>(D) <math>-20.0478</math></p>

## Section II

70 marks

Attempt Questions 11 – 14

Allow about 1 hours and 45 minutes for this section

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11 – 14, your responses should include relevant mathematical reasoning and/or calculations.

---

**Question 11** (15 marks) Use a SEPARATE writing booklet

(a) Solve  $\frac{3}{2x-1} < 2$  3

(b) Find  $\int_0^{\frac{\pi}{2}} \frac{dx}{9+4x^2}$  giving your answer in exact form 3

(c) Let  $A$  be the point  $(-2, 5)$  and  $B$  be the point  $(4, 1)$  2  
Find the coordinates of the point  $P$  that divides the interval externally in the ratio 4: 3.


(d) Evaluate  $\int \frac{x dx}{\sqrt{x-2}}$  using the substitution  $u^2 = x-2$  3

(e) Evaluate  $\lim_{x \rightarrow 0} \frac{\sin\left(\frac{x}{3}\right)}{2x}$  2

(f) Find  $\frac{d}{dx}\left(e^{\sin^{-1}(3x)}\right)$  2

**Question 12** (15 marks) Use a SEPARATE writing booklet

- (a) ST is a tangent at T, AT = 24cm, BT = 10cm, BS = 15cm and ST =  $h$  cm. O is the centre of the circle. 2




Find the value of  $h$  correct to one decimal place

- (b) i) Express  $\sqrt{3} \cos \theta - \sin \theta$  in the form  $R \cos(\theta + \alpha)$  where  $R > 0$  and  $0 < \alpha < 2\pi$  2
- ii) Hence or otherwise solve the equation  $\sqrt{3} \cos \theta - \sin \theta = 1$ , for  $0 \leq \theta \leq 2\pi$  2
- (c) Solve the equation  $x^3 - 21x^2 + 126x - 216 = 0$  given that the three roots form a geometric series 3
- (d) The acceleration of a particle moving in a straight line is given by  $\frac{d^2x}{dt^2} = -2e^{-x}$  where  $x$  is the displacement (in metres) from the origin. Initially the particle is at the origin and is moving with a velocity of  $2ms^{-1}$
- i) Prove that  $v = 2e^{\frac{x}{2}}$  2
- ii) Find an expression for the displacement  $x$  at any time  $t$ . 2

**Question 12 continues on page 8**

Question 12 (continued)

- (e) The diagram shows a conical wheat flue. The flue is being filled at the rate of  $2\text{m}^3 / \text{minute}$ . The height of wheat at any time, ' $t$ ' minutes, is ' $h$ ' metres, and the radius of the wheat's top surface is ' $r$ ' metres.



- i) Show that  $r = \frac{3h}{10}$  1
- ii) Find the rate at which the height is increasing when the height of wheat is 8 m 3  
(The volume of a cone is given by  $V = \frac{1}{3}\pi r^2 h$ )



**Question 13** (15 marks) Use a SEPARATE writing booklet

- (a) Consider the function  $f(x) = 2 \sin^{-1}(x-1)$ .
- iii) Find the domain and range of the function. 1
  - iv) Sketch the graph of the function. 1
  - v) Find the equation of the inverse function. 2
- (b) Four people go to a pizza festival, where four different gourmet pizzas A, B, C and D are offered. Each person chooses a pizza at random to try.
- i) Find the probability that they all choose different pizzas. 2
  - ii) Find the probability that exactly two of the people choose pizza A. 2
- (c)
- i)  $T(2at, at^2)$  is a point on the parabola  $x^2 = 4ay$ . Show that the normal to the parabola at  $T$  has equation  $x + ty - 2at - at^3 = 0$ . 2
  - ii)  $P$  and  $Q$  are points on the parabola  $x^2 = 4ay$  with parameter values  $t = 1$  and  $t = 2$  respectively. Show that the normal to the parabola at  $P$  and  $Q$  intersect at a point  $R$  on the parabola. 2
- (d) The formula for the  $n$ th term  $a_n$  of the Fibonacci sequence, 3

1, 1, 2, 3, 5, 8, 13, 21, 34, ...

is given by,


$$a_n = \begin{cases} 1 & \text{for } n = 1 \text{ and } 2 \\ a_{n-2} + a_{n-1} & \text{for } n > 2 \end{cases}$$

Prove by mathematical induction that,

$$a_n = \frac{(1 + \sqrt{5})^n - (1 - \sqrt{5})^n}{\sqrt{5} 2^n}$$

**Question 14** (15 marks) Use a SEPARATE writing booklet

(a)



In the diagram above,  $P$  is the midpoint of the chord  $AB$  in the circle with centre  $O$ . A second chord  $ST$  passes through  $P$ , and the tangents at the endpoints meet,  $AB$  produced at  $M$  and  $N$  respectively.


Join  $OS$ .

- i) Explain why  $OPNT$  is a cyclic quadrilateral. 2
- ii) Explain why  $OPSM$  is also cyclic. 1
- iii) Let  $\angle OTS = \theta$ . Show that  $\angle ONP = \angle OMP = \theta$ . 2
- iv) Hence, prove that  $AM = BN$ . 1

**Question 14 continues on page 11**

Question 14 (continued)

(b)



In the diagram above, a large number of projectiles are fired simultaneously from  $O$ , each with the same velocity  $V$  m/s, but different angles of projection  $\theta$ , at a wall  $d$  metres from  $O$ . The projectiles are fired so that they all lie in the same vertical plane perpendicular to the wall.

You may assume that the equations of motion at time  $t$  are given by:

$$x = Vt\cos\theta \quad \text{and} \quad y = -\frac{1}{2}gt^2 + Vt\sin\theta.$$

- i) Using these two equations of motion, prove that the relationship between the height  $y$  and time  $t$  is: 2

$$4y^2 + 4gt^2y + (g^2t^4 + 4x^2 - 4v^2t^2) = 0.$$

- ii) Show that the first impact at the wall occurs at time  $t = \frac{d}{V}$  and that this projectile was fired horizontally. 2

- iii) Hence, find where this projectile hits the wall. 1

- iv) Show that for  $t > \frac{d}{V}$ , there are *two* impacts at time  $t$ , and that the distance between these is: 2

$$2\sqrt{V^2t^2 - d^2}.$$

- v) Given that  $V = 10$  m/s and  $d = 10$  metres, what are the initial angles of projection of the two projectiles that will strike the wall simultaneously  $20\sqrt{3}$  metres apart. 2

End of Examination ☺

Multiple choices

① I ----- S  
 GP  
 6

(D)

②  $f(x) = 3x^2 - 4$

$$f'(1.6) = 3.68$$

$$f(1.6) = -0.304$$

(A)

$$x_2 = 1.6 - \frac{-0.304}{3.68} = 1.68$$

③

$$\sin^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{6}$$

(C)

④  $y = \frac{e^x - 2}{e^x + 2} = \frac{e^x + 2 - 4}{e^x + 2} = 1 - \frac{4}{e^x + 2}$

(B)

when  $x \rightarrow \infty$ ,  $y \rightarrow 1$ 

$$\text{if } x \rightarrow -\infty \quad y \rightarrow 1 - \frac{4}{0+2} = -1$$

⑤

(A)

⑥

$$A = 2 \times 2\pi - \int_0^{2\pi} \sin^2 x$$

$$= 4\pi - \frac{1}{2} \int_0^{2\pi} (1 - \cos 2x) dx$$


$$= 4\pi - \frac{1}{2} \left[ x - \frac{1}{2} \sin 2x \right]_0^{2\pi}$$

$$= 3\pi$$

(B)

$$\textcircled{7} \quad \cos 2x = -\frac{1}{\sin\left(\frac{\pi}{2} - x\right)}$$

$$= \frac{-1}{\cos x}$$



from the graph, there are 2 solutions or.

solve  $\cos 2x = -\frac{1}{\cos x}$

$$2\cos^2 x - \cos x + 1 = 0$$

$$\Rightarrow x = \pm \pi$$

$$\textcircled{8} \quad \sin(\sin^{-1} x) = x$$

domain  $-1 \leq x \leq 1$

range  $-1 \leq 1 \leq 1$

$\textcircled{C}$  ✓

$\textcircled{9}$

$$u = \sqrt{x}$$

$$\frac{du}{dx} = \frac{1}{2\sqrt{x}} \Rightarrow dx = 2u du$$

$$\int \frac{dx}{x + \sqrt{x}} = \int \frac{2u du}{u^2 + u} = 2 \int \frac{du}{u+1}$$

$\textcircled{A}$

$\textcircled{10}$

$$\frac{d}{dx} \frac{1}{2} v^2 = \frac{d}{dx} (18 + 3x - x^2)$$

$$= 3 - 2x = 2\left(\frac{3}{2} - x\right)$$

$$n = \sqrt{2} \quad T = \frac{2\pi}{\sqrt{2}} = \sqrt{2}\pi$$

$$x^2 - 3x - 18 = 0$$

$$x = 6$$

$$a = \frac{9}{2}$$

$$(x-6)(x+3) = 0$$

$$x = -3$$

$\textcircled{C}$  ✓

Q11


a)  $\frac{3}{2x-1} < 2$

$3(2x-1) = 2(2x-1)^2 < 0$

$(2x-1)(3=4x+2) < 0$

$(2x=1)(5=4x) < 0$

$x < 1/2$  and  $x > 5/4$



generally well done

b)  $\int_0^{\sqrt{3/2}} \frac{dx}{9+4x^2} = \int_0^{\sqrt{3/2}} \frac{dx}{3^2+(2x)^2}$

$= \frac{1}{2} \left[ \tan^{-1} \frac{2x}{3} \right]_0^{\sqrt{3/2}}$

$= \frac{1}{6} \times \frac{\pi}{6} = \frac{\pi}{36}$

generally well done

c)  $x = \frac{-4 \times 4 + 3 \times -2}{-1} = 22$

(22, -11)

$y = \frac{-4 \times 1 + 3 \times 5}{-1} = 11$

generally well done

(d)  $u^2 = x-2 \Rightarrow x = u^2 + 2$   
 $2u du = dx$

generally, well done

$$\int \frac{x dx}{\sqrt{x-2}} = \int \frac{u^2+2}{u} \cdot 2u du$$

$$= \int 2u^2 + 4 du$$

$$= \frac{2}{3} u^3 + 4u + C$$

$$= \frac{2}{3} (x-2)^{3/2} + 4(x-2) + C$$

few students did not express the answer in terms of  $x$ .

e)  $\lim_{x \rightarrow 0} \frac{\sin \frac{x}{3}}{\frac{x}{3}} = \frac{1/3}{2} = \frac{1}{6}$   
 $= \frac{1}{6}$

Working must be shown correctly to obtain full marks

generally well done

f)  $\frac{d}{dx} e^{\sin(3x)} = e^{\sin(3x)} \cdot 3$   
 $= \frac{3e^{\sin(3x)}}{\sqrt{1-9x^2}}$

generally well done

$$= \frac{3e^{\sin(3x)}}{\sqrt{1-9x^2}}$$

2)  $R = \sqrt{3+1} = 2$

$\alpha = \tan^{-1} \frac{1}{\sqrt{3}} = \frac{\pi}{6}$

$\therefore \sqrt{3} \cos \theta - \sin \theta = 2 \cos(\theta + \frac{\pi}{6})$

$2 \cos(\theta + \frac{\pi}{6}) = 1$

$\theta + \frac{\pi}{6} = \frac{\pi}{3}, \frac{5\pi}{3}$

$\therefore \theta = \frac{\pi}{6}$  and  $\frac{3\pi}{2}$

c) let the 3 roots be  $\frac{a}{r}, a$  and  $ar$

3 roots =  $\frac{a}{r} + a + ar = 21$  (1)

Prod of roots  $\frac{a}{r} \cdot a \cdot ar = 216$

$a^3 = 216$

$a = 6$  Sub  $\Rightarrow$  (1)

$\Rightarrow \frac{6}{r} + 6 + 6r = 21$

$6r^2 = 15r + 6 = 0$

$(3r-1)(r-2) = 0$

$\Rightarrow r = \frac{1}{3}$  or  $2$

$\therefore$  The 3 roots are 3, 6 and 12

generally well done

generally well done

your student to use polynomial division to find the answers.

few students had no idea how to approach this problem

✓ correct answers

✓ correct working

✓ correct value of a



c)  $\frac{d}{dt} \frac{1}{2} v^2 = -2e^{-x}$

$\frac{1}{2} v^2 = 2e^{-x} + c$

when  $x=0, v=2$

$\Rightarrow c = 2 - 2 = 0$

$\frac{1}{2} v^2 = 2e^{-x}$

$v^2 = 4e^{-x}$

$v = \pm 2e^{-x/2}$

when  $t=0, x_0=0, v^2 = 2m/s > 0$

$v = 2e^{-x/2}$

(ii)  $\frac{dx}{dt} = 2e^{-x/2}$

$dt = \frac{1}{2} e^{x/2}$

$\Rightarrow t = e^{x/2} + c$

when  $t=0, x=0 \Rightarrow c = -1$

$e^{x/2} = t + 1$

hence  $x = 2 \ln(t + 1)$

or  $x = \ln(t + 1)^2$

✓ correct expression of  $\frac{1}{2} v^2$

✓ correct answer of  $v^2$

✓ final answer with reason

✓ correct expression for  $t$


✓ form of answer

Many students assumed  
th. of  $v = 2e^{-x/2}$  without  
giving reason

generally well done

Q12 cont

a)



using similar  $\Delta$ 's'

$$\frac{r}{3} = \frac{h}{10}$$

$$\Rightarrow r = \frac{3h}{10}$$

well done

$$ii) V = \frac{1}{3} \pi r^2 h = \frac{1}{3} \pi \left(\frac{3h}{10}\right)^2 h$$

$$= \frac{3\pi h^3}{100}$$

$$\frac{dV}{dh} = \frac{9\pi h^2}{100}$$

$$\frac{dh}{dt} = \frac{dV}{dV} \times \frac{dV}{dt} \quad \text{where } \frac{dV}{dt} = 2\text{cm}^3/\text{min.}$$

$$= \frac{100}{9\pi h^2} \times 2 = \frac{200}{9\pi h^2}$$

Sol<sup>n</sup>  $h = 8$

$$\Rightarrow \frac{dh}{dt} = \frac{200}{9\pi \times 8^2}$$

$$= \frac{25}{72\pi} \text{ cm}$$

$$= 0.11 \text{ cm}$$

many students finding

$\frac{d}{dt} \frac{1}{3} \pi r^2 h$  without

sub.  $h = \frac{3h}{10}$  before

hand. This gave

incorrect answer

for  $\frac{dh}{dt}$ .

Some didn't know

way using  $\frac{dh}{dt} = \frac{dV}{dV} \times \frac{dV}{dh} \times \frac{dr}{dr} \times \frac{dh}{dh}$


and find these derivative functions.

✓ correct expression for  $\frac{dV}{dt}$

✓ correct expression for  $\frac{dV}{dh}$

✓ correct substitution &

answer.

(a)	Consider the function $f(x) = 2\sin^{-1}(x-1)$ .							
	i) Find the domain and range of the function.	1						
	<table border="1" style="width: 100%;"> <thead> <tr> <th style="width: 80%;">Criteria</th> <th style="width: 20%;">Marks</th> </tr> </thead> <tbody> <tr> <td>• Provides a correct solution</td> <td>1</td> </tr> </tbody> </table>	Criteria	Marks	• Provides a correct solution	1			
Criteria	Marks							
• Provides a correct solution	1							
	<p><i>Sample answer:</i>  Domain: <math>\{-1 \leq x-1 \leq 1\}</math>  : <math>\{0 \leq x \leq 2\}</math>  Range: <math>\{-\pi \leq y \leq \pi\}</math></p>	<p><i>Mostly done well.</i></p>						
	ii) Sketch the graph of the function.	2						
	<table border="1" style="width: 100%;"> <thead> <tr> <th style="width: 80%;">Criteria</th> <th style="width: 20%;">Marks</th> </tr> </thead> <tbody> <tr> <td>• Draws a correct sketch</td> <td>2</td> </tr> <tr> <td>• Draws a graph showing the correct shape, or equivalent merit</td> <td>1</td> </tr> </tbody> </table>	Criteria	Marks	• Draws a correct sketch	2	• Draws a graph showing the correct shape, or equivalent merit	1	
Criteria	Marks							
• Draws a correct sketch	2							
• Draws a graph showing the correct shape, or equivalent merit	1							
	<p><i>Sample solution:</i></p>  <p><i>be careful the shape is right.</i>  <u>(2, pi) is a horizontal tangent</u></p>							

	iii) Find the equation of the inverse function.	2						
	<table border="1"> <thead> <tr> <th data-bbox="169 288 796 312">Criteria</th> <th data-bbox="796 288 958 312">Marks</th> </tr> </thead> <tbody> <tr> <td data-bbox="169 312 796 336">• Provides a correct solution</td> <td data-bbox="796 312 958 336">2</td> </tr> <tr> <td data-bbox="169 336 796 387">• Swaps variables and makes some progress towards a correct solution, or equivalent merit.</td> <td data-bbox="796 336 958 387">1</td> </tr> </tbody> </table>	Criteria	Marks	• Provides a correct solution	2	• Swaps variables and makes some progress towards a correct solution, or equivalent merit.	1	
Criteria	Marks							
• Provides a correct solution	2							
• Swaps variables and makes some progress towards a correct solution, or equivalent merit.	1							
	<p><b>Sample solutions:</b>  Swap unknowns  <math>y = 2 \sin^{-1}(x-1)</math>  <math>\frac{1}{2}y = \sin^{-1}(x-1)</math>  <math>\sin\left(\frac{y}{2}\right) = x-1</math>  <math>x = 1 + \sin\left(\frac{y}{2}\right)</math>  hence  <math>f^{-1}(x) = 1 + \sin\left(\frac{x}{2}\right)</math> for <math>-\pi \leq x \leq \pi</math></p> <p style="text-align: right;"><i>well done</i></p>							
(b)	Four people go to a pizza festival, where four different gourmet pizzas A, B, C and D are offered. Each person chooses a pizza at random to try.							
	i) Find the probability that they all choose different pizzas.	1						
	<table border="1"> <thead> <tr> <th data-bbox="169 948 796 971">Criteria</th> <th data-bbox="796 948 958 971">Marks</th> </tr> </thead> <tbody> <tr> <td data-bbox="169 971 796 995">• Provides a correct solution</td> <td data-bbox="796 971 958 995">1</td> </tr> </tbody> </table>	Criteria	Marks	• Provides a correct solution	1			
Criteria	Marks							
• Provides a correct solution	1							
	<p><b>Sample solution:</b>  <math>P(\text{all different}) = \frac{4!}{4^4}</math>  <math>= \frac{3}{32}</math></p>							
	ii) Find the probability that exactly two of the people choose pizza A.	2						
	<table border="1"> <thead> <tr> <th data-bbox="169 1197 796 1220">Criteria</th> <th data-bbox="796 1197 958 1220">Marks</th> </tr> </thead> <tbody> <tr> <td data-bbox="169 1220 796 1244">• Provides a correct solution</td> <td data-bbox="796 1220 958 1244">2</td> </tr> <tr> <td data-bbox="169 1244 796 1295">• Makes progress towards an answer with at most one option missed or one element unconsidered, or equivalent merit.</td> <td data-bbox="796 1244 958 1295">1</td> </tr> </tbody> </table>	Criteria	Marks	• Provides a correct solution	2	• Makes progress towards an answer with at most one option missed or one element unconsidered, or equivalent merit.	1	
Criteria	Marks							
• Provides a correct solution	2							
• Makes progress towards an answer with at most one option missed or one element unconsidered, or equivalent merit.	1							
	<p><b>Sample solution:</b>  <math>P(\text{exactly 2 choose A}) = {}^4C_2 \left(\frac{1}{4}\right)^2 \left(\frac{3}{4}\right)^2</math>  <math>= \frac{27}{128}</math></p> <p style="text-align: right;"><i>Make sure to show working</i></p>							

(c)	$P$ and $Q$ are points on the parabola $x^2 = 4ay$ with parameter values $t = 1$ and $t = 2$ respectively.	
	i) Given the equation of the normal at $T$ is $x + ty - 2at - at^3 = 0$ (Do not prove). Find the equation of the normal at $P$ and at $Q$ .	1
	Criteria	Marks
	<ul style="list-style-type: none"> <li>Provides a correct solution</li> </ul>	1
	<p><i>Sample solution:</i>  when <math>t</math> is 1 the normal at <math>P</math>:  <math>x + 1.y - 2.a.1 - a.1^3 = 0</math>  <math>x + y = 3a</math></p> <p>when <math>t</math> is 2 the normal at <math>Q</math>:  <math>x + 2.y - 2.a.2 - a.2^3 = 0</math>  <math>x + 2y = 12a</math></p> <p><i>Many students confused <math>p</math> &amp; <math>q</math> as variables instead of substituting in 1 &amp; 2</i></p>	
	ii) Show that the normal to the parabola at $P$ and $Q$ intersect at a point $R$ on the parabola.	2
	Criteria	Marks
	<ul style="list-style-type: none"> <li>Provides a correct solution</li> </ul>	2
	<ul style="list-style-type: none"> <li>Solve the normal equations simultaneously and attempt to demonstrate solution lies on the parabola, or equivalent merit</li> </ul>	1
	<p><i>Sample solution:</i>  <math>x + y = 3a</math> --- 1  <math>x + 2y = 12a</math> -- 2  subtracting 2 from 1  <math>-y = -9a</math>  <math>y = 9a</math>  substituting into 1  <math>x + 9a = 3a</math>  <math>x = -6a</math>  giving <math>R(-6a, 9a)</math> as the point of intersection  <math>x^2 = 36a^2</math>  <math>= 4a(9a)</math>  <math>= 4ay</math>  <math>\therefore R</math> lies on the parabola</p> <p><i>Some students forget to show the point lies on the parabola</i></p>	

(d)	i) Prove the following by the process of mathematical induction.  $\ln(2) + \ln\left(\frac{3}{2}\right) + \ln\left(\frac{4}{3}\right) + \dots + \ln\left(\frac{n+1}{n}\right) = \ln(n+1)$	3
-----	---	---

Criteria	Marks
• Provides a correct solution	3
• Demonstrate that k implies k+1	2
• Demonstrate the first case	1

**Sample solution:**  
 when  $n = 1$   

$$\ln\left(\frac{1+1}{1}\right) = \ln(2)$$

$$= \ln(1+1)$$
 $\therefore$  statement true for  $n = 1$   
 Assume true for  $n = k$   
 that is:  

$$\ln(2) + \ln\left(\frac{3}{2}\right) + \ln\left(\frac{4}{3}\right) + \dots + \ln\left(\frac{k+1}{k}\right) = \ln(k+1)$$
 when  $n = k+1$   

$$\ln(2) + \ln\left(\frac{3}{2}\right) + \ln\left(\frac{4}{3}\right) + \dots + \ln\left(\frac{k+1}{k}\right) + \ln\left(\frac{k+1+1}{k+1}\right)$$

$$= \ln(k+1) + \ln\left(\frac{k+1+1}{k+1}\right)$$

$$= \ln((k+1)+1)$$
 as required  
 $\therefore$  as  $n=1$  is true and  $n=k$  true proves  $n=k+1$  is true  
 the statement is true by the principle of mathematical induction.

*Some students made logically false statements. learn the process of induction*

	ii) Hence find $p$ for which $\sum_{n=1}^p \ln\left(\frac{n+1}{n}\right) \geq \pi$	1
--	--	---

Criteria	Marks
• Provides a correct solution	1

**Sample solution:**  

$$\sum_{n=1}^p \ln\left(\frac{n+1}{n}\right) \geq \pi$$

$$\ln(p+1) \geq \pi$$

$$p+1 \geq e^\pi$$

$$p \geq e^\pi - 1$$

$$p \geq 22.14\dots$$
 $\therefore p = 23$

*Mostly done well*

*sum of*

<p>14a) (i)</p>	<p>From the equations of motion,</p> $V \cos \theta = \frac{x}{t}$ $V \sin \theta = \frac{y}{t} + \frac{1}{2}gt$ $\left(\frac{x}{t}\right)^2 + \left(\frac{y}{t} + \frac{1}{2}gt\right)^2 = V^2(\cos^2 \theta + \sin^2 \theta)$ $(x)^2 + \left(y + \frac{1}{2}gt^2\right)^2 = V^2t^2$ $x^2 + y^2 + ygt^2 + \frac{1}{4}g^2t^4 = V^2t^2$ $4y^2 + 4ygt^2 + (4x^2 + g^2t^4 - 4V^2t^2) = 0$	<p>2 marks: correct proof</p> <p>1 mark: substantial progress is made in trying to eliminate <math>\sin \theta</math> and <math>\cos \theta</math></p>	<p>Students who could understand, "eliminate <math>\theta</math> was the purpose of the question did it correctly. Well done by many students</p>
<p>(ii)</p>	<p><math>t = \frac{x}{V \cos \theta}</math> where <math>x = d</math> and <math>V</math> are constants. Hence, <math>t</math> is minimum when <math>\cos \theta</math> is maximum which occurs when <math>\theta = 0</math>. i.e. The projectile is fired horizontally. 1 mark</p> <p><math>\theta = 0</math>, then <math>\cos \theta = 1</math>. Hence, <math>t = \frac{x}{V \cos \theta} = \frac{d}{V}</math> seconds. 1 mark</p>	<p>1 mark: proves angle of projection <math>\theta = 0</math>.</p> <p>1 mark: proves the result with correct working and reasoning.</p>	<p>Very few students realised that "first" impact means, you need to minimise <math>t</math>.</p> <p>Most of the students <u>used</u> <math>\theta = 0</math> to prove <math>t = \frac{d}{V}</math> and vice versa. These students were awarded 0 marks. They have not proved any of the required results.</p>
<p>(iii)</p>	<p><math>y = -\frac{1}{2}gt^2 + Vt \sin \theta</math> <math>\theta = 0</math>, then <math>\sin \theta = 0</math>.</p> <p><math>y = -\frac{1}{2}g\left(\frac{d}{V}\right)^2 = -\frac{gd^2}{2V^2}</math> (hits below the horizontal.)</p>	<p>1 mark: substitutes <math>\theta = 0</math> into the equation and gives the correct result.</p>	<p>Most students got this mark, except those who did not realise <math>\theta = 0</math>.</p>
<p>(iv)</p>	<p>To have two impacts on the wall, there need to prove that here are two solutions for <math>y</math> when <math>t &gt; \frac{d}{V}</math>.</p> $4y^2 + 4gt^2y + (g^2t^4 + 4x^2 - 4V^2t^2) = 0$ <p><math>x = d</math>,</p> $\Delta = (4gt^2)^2 - 4 \times 4(g^2t^4 + 4d^2 - 4V^2t^2)$ $= 16g^2t^4 - 16(g^2t^4 + 4d^2 - 4V^2t^2)$ $= 64(V^2t^2 - d^2)$ <p>When <math>t &gt; \frac{d}{V}</math>, <math>Vt &gt; d</math>.</p> <p>Hence, <math>V^2t^2 - d^2 &gt; 0</math>, and <math>\Delta &gt; 0</math>.</p> <p>i.e. There are two solutions for <math>y</math>. This means that there are two real and</p>	<p>1 mark: proves <math>\Delta &gt; 0</math> for the quadratic in <math>y</math>.</p>	<p>Many students did not attempt this question. Those who realised two impacts at given <math>t</math> meant that there are two values for <math>y</math>.</p> <p>Majority of the students who attempted this question got 1 mark for setting <math>\Delta &gt; 0</math>.</p>

distinct roots for  $y$  and so, there are two impacts at the same  $t$ . 1 mark.

Now the distance between the impacts equals the difference between the roots.

$$\text{The roots are } \frac{-b \pm \sqrt{\Delta}}{2a}$$

$$\text{Hence the difference} = \frac{2\sqrt{\Delta}}{2a} = \frac{\sqrt{\Delta}}{a}$$

$$= \frac{\sqrt{64(v^2t^2 - d^2)}}{4}$$

$$= 2\sqrt{(v^2t^2 - d^2)} \quad \text{1 mark}$$

**Another approach:**

$$y_1 + y_2 = -gt^2$$

$$y_1y_2 = \frac{g^2t^4 + 4x^2 - 4v^2t^2}{4}$$

$$y_1 + y_2 = \sqrt{(y_1 + y_2)^2 - 4y_1y_2}$$

$$= \sqrt{g^2t^4 - g^2t^4 - 4x^2 + 4v^2t^2}$$

$$= 2\sqrt{v^2t^2 - x^2}$$

$$= 2\sqrt{v^2t^2 - d^2}$$

1 mark: correct answer from correct working

(v)

$$\text{Distance} = 2\sqrt{(v^2t^2 - d^2)}$$

$$= 2\sqrt{(100t^2 - 100)}$$

$$= 20\sqrt{(t^2 - 1)}$$

$$20\sqrt{(t^2 - 1)} = 20\sqrt{3}$$

$$t^2 - 1 = 3$$

$$t = 2, \quad t > 0$$


$$\cos\theta = \frac{x}{Vt} = \frac{10}{10 \times 2} = \frac{1}{2}$$

Hence,  $\theta = 60^\circ$  and  $-60^\circ$  are the angles of projection.

*Students who realised this question is a follow up of (iv) got the mark.*

*Many set  $y = 20\sqrt{3}$  in  $y = V \sin\theta - \frac{1}{2}gt^2$  and tried to solve for  $t$  unsuccessfully.*



<p>14b) (i)</p>	<p><math>\angle OTN = 90^\circ</math> (<math>OT \perp TN</math>, angle between tangent and radius is <math>90^\circ</math>)  <math>\angle OPN = 90^\circ</math> (the line from the centre that <b>bisects the chord</b> is perpendicular to the chord) 1 mark  <math>OPNT</math> is cyclic as the opposite angles add to <math>180^\circ</math> 1 mark</p>	<p>1 mark: both reasoning statements are correct   1 mark: gives the reason for the quadrilateral to be cyclic.</p>	<p><i>Well done.</i>  A very common error the statement:  <b>Line from the centre to the tangent makes <math>90^\circ</math>.</b></p>  <p><i>Clery the diagram above doesn't.</i>  Also, as shown in the answers, only if the chord is bisected, it is perpendicular.</p> <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p><b>Note: you need to learn and present the statements of the theorems accurately.</b></p> </div>
<p>(ii)</p>	<p>Similarly, <math>\angle OPM = 90^\circ</math> (from (i) <math>OP \perp PN</math> and hence <math>MN</math>.)  <math>\angle OSM = 90^\circ</math> (angle between tangent and radius is <math>90^\circ</math>)  <math>OPSM</math> is cyclic as the angles subtended by arc <math>OM</math> in the same segment are equal.) 1 mark</p>	<p>1 mark: gives the correct reasoning for the quadrilateral to be cyclic.</p>	
<p>(iii)</p>	<p><math>\angle OTS = \angle OST = \theta</math> (<math>OT = OS</math> radii, angles opposite equal sides of isosceles <math>\triangle OTS</math> 1 mark  Now, <math>\angle OTP = \angle ONP = \theta</math> and <math>\angle OSP = \angle OMP = \theta</math> angles in the same segment of cyclic quadrilaterals <math>OTNP</math> and <math>OPSM</math>. 1 mark   Hence, <math>\angle ONP = \angle OMP = \theta</math>.</p>	<p>2 marks: both reasoning correct   1 mark only: if substantial progress</p>	<p><i>A common error was in the interpretation of the diagram. Some students misunderstood the notation on the diagram meant <math>MP = PN</math>, rather than <math>AP = PB</math>. You always need to verify the diagram from the description of facts stated in the question.</i></p>
<p>(iv)</p>	<p><math>\triangle OMN</math> is isosceles (<math>OM = MN</math> as <math>\angle ONP = \angle OMP = \theta</math>, sides opposite equal angles are equal)  <math>OP</math> is the altitude.  Hence, <math>MP = PN</math> (perpendicular bisects the opposite side)  Hence, <math>MP - AP = PN - PB</math> (equals subtracted from equals)  ie. <math>AM = BN</math></p>	<p>1 mark: correct reasoning</p>	<p><i>Well done</i></p>