

Name:

Maths Teacher:

SYDNEY TECHNICAL HIGH SCHOOL



Year 12

Mathematics Extension 2

TRIAL HSC

2016

Time allowed: 3 hours plus 5 minutes reading time

General Instructions:

- Reading time – 5 minutes
- Working time – 3 hours
- Marks for each question are indicated on the question.
- Board - Approved calculators may be used
- In Questions 11- 16, show relevant mathematical reasoning and/or calculations.
- Full marks may not be awarded for careless work or illegible writing
- *Begin each question on a **new** page*
- Write using black pen
- All answers are to be in the writing booklet provided
- A BOSTES reference sheet is provided at the rear of this Question Booklet, and may be removed at any time.

Total marks – 100

Section I Multiple Choice

10 Marks

- Attempt Questions 1-10
- Allow 15 minutes for this section

Section II

90 Marks

- Attempt Questions 11-16
- Allow 2 hours 45 minutes for this section

Section I

10 marks

Attempt Questions 1- 10

Allow about 15 minutes for this section

Use the multiple- choice answer sheet located in your answer booklet for Questions 1 -10

1. Which conic has eccentricity $\frac{\sqrt{3}}{3}$?

(A) $\frac{x^2}{3} + \frac{y^2}{2} = 1$

(B) $\frac{x^2}{3^2} + \frac{y^2}{2^2} = 1$

(C) $\frac{x^2}{3} - \frac{y^2}{2} = 1$

(D) $\frac{x^2}{3^2} - \frac{y^2}{2^2} = 1$

2. What value of z satisfies; $z^2 = 20i - 21$?





(A) $-2 + 5i$

(B) $2 - 5i$

(C) $2 + 5i$

(D) $5 - 2i$

3. Which graph represents the curve, $y = (x+3)^2(x-1)^3$?



4. The polynomial $2x^4 - 17x^3 + 45x^2 - 27x - 27$ has a triple root at $x = \alpha$.

What is the value of α ?

(A) $-\frac{1}{2}$

(B) $\frac{1}{2}$

(C) -3

(D) 3

5. If $z_1 = 1 + 2i$ and $z_2 = 3 - i$ then $z_1 + \overline{z_2}$ is,

(A) $\frac{1}{2} - \frac{1}{2}i$

(B) $\frac{1}{2} + \frac{1}{2}i$

(C) $4 + 3i$

(D) $\frac{5}{8} + \frac{5}{8}i$

6. Which expression is equal to, $\int \frac{x^2}{\cos^2 x} dx$?

(A) $2x \tan x - 2 \int \tan x dx$

(B) $\frac{1}{3}(x^3 \sec^2 x - \int x^3 \tan x dx)$

(C) $x^2 \tan x - 2 \int x \tan x dx$

(D) $x^2 \tan x - 2 \int x \sec^2 x dx$

7. What is the natural domain of the function $f(x) = \frac{1}{2}(x\sqrt{x^2-1} - \ln(x + \sqrt{x^2-1}))$?

(A) $x \leq -1$ or $x \geq 1$

(B) $-1 \leq x \leq 1$

(C) $x \geq 1$

(D) $x \leq -1$

8. If α, β, δ are the roots of $x^3 + x - 1 = 0$, then an equation with roots

$$\frac{(\alpha+1)}{2}, \frac{(\beta+1)}{2}, \frac{(\delta+1)}{2} \text{ is?}$$

- (A) $x^3 - 3x^2 + 4x - 3 = 0$
- (B) $x^3 + 3x^2 + 4x + 1 = 0$
- (C) $x^3 - 6x^2 + 16x - 24 = 0$
- (D) $8x^3 - 12x^2 + 8x - 3 = 0$


9. The complex number Z satisfies $|Z+2|=1$

What is the smallest positive value of the $\arg(z)$ on the Argand diagram?

- (A) $\frac{\pi}{3}$
- (B) $\frac{5\pi}{6}$
- (C) $\frac{2\pi}{3}$
- (D) $\frac{\pi}{6}$

10. The base of a solid is the region bounded by the parabola $x = 4y - y^2$ and the y axis.

Vertical cross sections are right angled isosceles triangles perpendicular to the x -axis as shown.



Which integral represents the volume of this solid?

- (A) $\int_0^4 2\sqrt{4-x} dx$
- (B) $\int_0^4 \pi(4-x) dx$
- (C) $\int_0^4 (8-2x) dx$
- (D) $\int_0^4 (16-4x) dx$

Question 11 (15 marks)

(a) Express $\frac{18+4i}{3-i}$ in the form, $x+iy$, where x and y are real. 2

(b) Consider the complex numbers $z = -1 + \sqrt{3}i$ and $w = \sqrt{2} \left(\cos\left(\frac{-\pi}{4}\right) + i \sin\left(\frac{-\pi}{4}\right) \right)$

(I) Evaluate $|z|$ 1

(II) Evaluate $\arg(z)$ 1

(III) Find the argument of $\frac{w}{z}$ 2


(c) (i) Find A, B and C such that 3

$$\frac{1}{x(x^2+4)} = \frac{A}{x} + \frac{Bx+C}{x^2+4}$$

(ii) Hence, or otherwise, find;

$$\int \frac{dx}{x(x^2+4)} \quad \text{2}$$

(d)



A particle moves along the x - axis. At time, $t=0$, the particle is at $x=0$.

Its velocity v at time t is shown on the graph above.

Copy or trace this graph onto your answer page.

(i) At what time is the acceleration the greatest? Explain your answer. 1

(ii) At what time does the particle first return to $x=0$? Explain your answer. 1

(iii) Sketch the displacement time graph for the particle in the interval, $0 \leq t \leq 9$. 2

Question 12 (15 marks) START THIS QUESTION ON A NEW PAGE.

(a) Find $\int x\sqrt{x+1}dx$ 2

(b) Evaluate

(i) $\int_0^{\frac{\pi}{4}} \sin x \cos 2x dx$ 2

(ii) $\int_1^e \frac{\ln x}{x^2} dx$ 2

(c) Find the equation of the normal to the curve, $3x^2y^3 + 4xy^2 = 6 + y$ at the point (1,1). 4

(d)

(i) Prove that,

$$\cos(A-B)x - \cos(A+B)x = 2 \sin Ax \sin Bx$$
 1

(ii) Using the above result, express the equation $\sin 3x \sin x = 2 \cos 2x + 1$,
as a quadratic equation in terms of $\cos 2x$ 2

(iii) Hence, solve, $\sin 3x \sin x = 2 \cos 2x + 1$ for $0 \leq x \leq 2\pi$ 2

Question 13 (15 marks) START THIS QUESTION ON A NEW PAGE.

(a) The function $y = f(x)$ is defined by the equation;

$$f(x) = \frac{x(x-4)}{4}$$

Without the use of calculus, draw sketches of each of the following, clearly labelling any intercepts, asymptotes and turning points.

- | | | |
|-------|------------------------|---|
| (i) | $y = f(x)$ | 1 |
| (ii) | $y^2 = f(x)$ | 2 |
| (iii) | $y = \frac{x x-4 }{4}$ | 2 |
| (iv) | $y = \tan^{-1} f(x)$ | 2 |
| (v) | $y = e^{f(x)}$ | 2 |

(b) Sketch the locus of z satisfying

- | | | |
|------|-----------------------------------|---|
| (i) | $Re(z) = z $ | 2 |
| (ii) | $Im(z) \geq 2$ and $ z-1 \leq 2$ | 2 |

(c) Write down the domain and range of $y = 2 \sin^{-1} \sqrt{1-x^2}$ 2

Question 14 (15 marks) **START THIS QUESTION ON A NEW PAGE.**

(a) Use the substitution $t = \tan \frac{x}{2}$ to find

4

$$\int_0^{\pi/2} \frac{1}{5 + 4 \cos x + 3 \sin x} dx$$

(b) The area enclosed by the curves $y = \sqrt{x}$ and $y = x^2$ is rotated about the y -axis.


Use the method of *cylindrical shells* to find the volume of the solid formed.

4

Question 14 continues on the next page....

Question 14 continued....

(c)




$P(a \sec \theta, b \tan \theta)$ lies on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$.

The tangent at P meets the line $x = -a$ and $x = a$ at R and Q respectively.

- | | | |
|-------|---|---|
| (i) | Show that the equation of the tangent is given by $\frac{x \sec \theta}{a} - \frac{y \tan \theta}{b} = 1$. | 2 |
| (ii) | Find the coordinates of Q and R . | 1 |
| (iii) | Show that QR subtends a right angle at the focus $S(ae, 0)$. | 2 |
| (iv) | Deduce that Q, S, R, S' are concyclic. | 2 |

Question 15 (15 marks) **START THIS QUESTION ON A NEW PAGE.**

- (a) In the diagram, AB and AC are tangents from A to the circle with centre O , meeting the circle at B and C respectively. ADE is a secant of the circle. G is the midpoint of DE . CG produced meets the circle at F .



- (i) Copy the diagram, using about one third of the page, into your answer booklet and prove that $ABOC$ and $AOGC$ are cyclic quadrilaterals 3
- (ii) Explain why $\angle OGF = \angle OAC$. 1
- (iii) Prove that $BF \parallel AE$ 3

(b)

(i) Let $I_n = \int_0^1 x^n \sqrt{1-x^3} dx$ for $n \geq 2$.

Show that: $I_n = \frac{2n-4}{2n+5} I_{n-3}$ for $n \geq 5$ 3

(ii) Hence find I_8 2

- (c) A sequence of numbers is given by $T_1 = 6$, $T_2 = 27$ and $T_n = 6T_{n-1} - 9T_{n-2}$ for $n \geq 3$.

Prove by Mathematical Induction that:

$$T_n = (n+1) \times 3^n \text{ for } n \geq 1 \quad \text{3}$$

Question 16 (15 marks) **START THIS QUESTION ON A NEW PAGE.**

(a) Show that the minimum value of $ae^{mx} + be^{-mx}$ is $2\sqrt{ab}$

if a, b and m are all positive constants.

4

(b) A particle of mass 1 kilogram is projected upwards under gravity (g) with a speed of $2k$

in a medium in which resistance to motion is $\frac{g}{k^2}$ times the square of the speed, where k

is a positive constant.

(i) Show that the maximum height (H) reached by the particle is

$$H = \frac{k^2}{2g} \ln 5$$

3


(ii) Show that the speed with which the particle returns to its starting point

$$\text{is given by } V = \frac{2k}{\sqrt{5}}$$

4

(c) The shaded region in the diagram is bounded by the curves $y = \sin x$, $y = \cos x$ and the line $y = 1$.

This region is rotated around the y -axis.



Calculate the volume of the solid formed, using the process of **Volume by Slicing**.

4


SIHS - Ext 2 Trial - Suggestion Solution

Section 1

1. A 2. C 3. C 4. D 5. B
 6. C * 7. C 8. D 9. B 10. C
 (ans given)

Section 2

Question 11

a) $\frac{18+4i}{3-i} \times \frac{3+i}{3+i}$	equating: $0 = A + B$
$= \frac{54 + 18i + 12i - 4}{10}$	$B = -1/4$
$= 50 + 30i$	$0 = C$
$= 5 + 3i$	$1e \ A = 1/4 \ B = -1/4 \ C = 0$
b) $\omega = \sqrt{2} \text{cis}(-\pi/4)$ $z = -1 + \sqrt{3}i$	Now $\int \frac{dx}{x(x^2+4)} = \int \frac{1}{4x} - \frac{1/4x}{x^2+4} dx$
i) $ z = \sqrt{1+3} = 2$	d) i. $t = 6.5$ (point of inflection on vel. curve is greatest acc)
ii) $\text{arg}(z) = 2\pi/3$	ii. When the area above the t-axis equals area below \therefore at $t = 4$
iii) $\text{arg}\left(\frac{\omega}{z}\right) = \text{arg}(\omega) - \text{arg}(z)$	iii. 
c) $\frac{1}{x(x^2+4)} = \frac{A}{x} + \frac{Bx+C}{x^2+4}$	
$\therefore 1 = A(x^2+4) + x(Bx+C)$	
let $x=0$	
$1 = A(4) \rightarrow A = 1/4$	

Question 12

a) $\int x\sqrt{x+1} dx$	ii. $\int_1^e \frac{\ln x}{x^2} dx$
one method:	$= \int_1^e x^{-2} \ln x dx$
let $u = x+1$	$= \ln x \cdot \frac{x^{-1}}{-1} - \int 1/x \cdot \frac{-1}{x} dx$
$du = 1 \therefore du = dx$	$= -\ln x \Big _1^e + \int_1^e x^{-2} dx$
$= \int (u-1)\sqrt{u} du$	$= -\left[\frac{1}{e} - 0\right] + \left[-\frac{1}{x}\right]_1^e$
$= \int u\sqrt{u} - \sqrt{u} du$	$= -1/e + \left[-1/e - (-1)\right]$
$= \int u^{3/2} - u^{1/2} du$	$= 1 - 2/e$
$= \frac{2}{5} u^{5/2} - \frac{2}{3} u^{3/2}$	e) $3x^2y^3 + 4xy^2 = 6 + y$ @ (1,1)
$= \frac{2}{5}(x+1)^{5/2} - \frac{2}{3}(x+1)^{3/2} + C$	$3x^2 \left[3y^2 \frac{dy}{dx}\right] + y^3 \cdot 6x + 4x \cdot 2y \frac{dy}{dx} + 4y^2 = \frac{dy}{dx}$
b) i. $\int_0^{\pi/4} \sin x \cos 2x dx$	$6xy^3 + 4x^2 = \frac{dy}{dx}(1 - 9x^2y^2 - 8xy)$
$= \int_0^{\pi/4} \sin x (2\cos^2 x - 1) dx$	at (1,1)
$= \int_0^{\pi/4} 2\sin x \cos^2 x^2 - \sin x dx$	$\frac{dy}{dx} = \frac{10}{-16}$
$= \left[-\frac{2}{3} \cos^3 x + \cos x\right]_0^{\pi/4}$	$M_T = -5/8 \therefore M_N = 8/5$
$= \frac{-2}{3} \left(\frac{1}{\sqrt{2}}\right)^3 + \frac{1}{\sqrt{2}} - \left[-\frac{2}{3}(1)^3 + 1\right]$	$y-1 = 8/5(x-1)$
$= \frac{2}{3\sqrt{2}} - 1$	$5y - 5 = 8x - 8$
$3\sqrt{2} \quad 3$	$8x - 5y - 3 = 0$

Question 12 - con't.

a)

1. Prove that

$$\cos(A-B)x - \cos(A+B)x = 2\sin A x \sin B x$$

$$\text{LHS} = \cos A x \cos B x + \sin A x \sin B x - [\cos A x \cos B x - \sin A x \sin B x]$$

$$= 2\sin A x \sin B x$$

= RHS.

ii. $\sin 3x \sin x = 2\cos 2x + 1$

$A=3$
 $B=1$

$$\therefore [\cos(3-1)x - \cos(3+1)x] \div 2 = \cos 2x + 1$$

$$\cos 2x - \cos 4x = 2\cos 2x + 2$$

$$\cos 2x - [2\cos^2 2x - 1] = 2\cos 2x + 2$$

$$\cos 2x - 2\cos^2 2x + 1 = 2\cos 2x + 2$$

$$2\cos^2 2x + 3\cos 2x + 1 = 0$$

iii. hence,

$$(2\cos 2x + 1)(\cos 2x + 1) = 0$$






$$\cos 2x = -\frac{1}{2} \quad \text{or} \quad \cos 2x = -1$$

$$2x = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}, \frac{10\pi}{3} \quad \text{or} \quad 2x = \pi, 3\pi$$


$$x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3} \quad \text{or} \quad x = \frac{\pi}{2}, \frac{3\pi}{2}$$

Question 13


a) $f(x) = x(x-4)$



b) $R(z) = |z|$
let $z = x + iy$
 $x = \sqrt{x^2 + y^2} \quad x \geq 0$
 $x^2 = x^2 + y^2$
 $y^2 = 0$
 $y = 0$ but $x \geq 0$



ii. $\text{Im}(z) \geq 2 \quad |z-1| \leq 2$
 $\therefore y \geq 2$ circle centre $(1, 0)$



c) $y = 2\sin^{-1}\sqrt{1-x^2}$
 $y/2 = \sin^{-1}\sqrt{1-x^2}$

O: $-1 \leq x \leq 1$

R: $0 \leq y \leq \pi$

Question 14.

a) $x^2 - y^2 = 1$

$x = \pi/2 \quad t = \tan \pi/4 = 1$

$a^2 - b^2$

$x=0 \quad t = \tan 0 = 0$

$2x/a^2 - 2y/b^2 \cdot \frac{dy}{dx} = 0$

$\therefore \int_0^1 \frac{1}{5+4\left(\frac{1-t^2}{1+t^2}\right) + 3\left(\frac{2t}{1+t^2}\right)} \cdot t^2 + 1 \cdot t^2 dt$

$\rho(\operatorname{asec}\theta, b\tan\theta)$

$\int_0^1 \frac{1}{2} dt \quad M_T = b/\operatorname{asin}\theta$

$\int_0^1 5(1+t^2) + 4(1-t^2) + 6t$

$\int_0^1 \frac{1}{2} dt \quad \text{Now } y - b\sin\theta = \frac{b}{a\sin\theta} \left(x - \frac{a}{\cos\theta}\right)$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$\int_0^1 \frac{1}{2} dt \quad \operatorname{asin}\theta - b\operatorname{asin}\theta = bx - ab$

$= 1/6$

ii. at $x = a$

$y = x^2 \quad y = \sqrt{x} \quad \frac{a\operatorname{sec}\theta - \tan\theta y}{a} = 1$

$\Delta v = 2\pi x (\sqrt{x} - x^2) \Delta x \quad \frac{1}{\cos\theta} - \frac{y}{a} \frac{\sin\theta}{\cos\theta} = \frac{\cos\theta}{\cos\theta}$

$V = \lim_{\Delta x \rightarrow 0} \sum 2\pi x (\sqrt{x} - x^2) \Delta x \quad 1 - y/a \sin\theta = \cos\theta$

$= 2\pi \int_0^1 x^{3/2} - x^3 dx \quad y = b(1 - \cos\theta)$

$= \left[\frac{2}{5} x^{5/2} - \frac{1}{4} x^4 \right]_0^1 \quad \text{or } y = b \left(\frac{\operatorname{sec}\theta - 1}{\tan\theta} \right)$

$= 2\pi \left[\frac{2}{5} - \frac{1}{4} - 0 \right] = \frac{3\pi}{10} u^3$

$\left[a, b(1 - \cos\theta) \right] \frac{1}{\sin\theta}$

at $R = -a$

$-a\operatorname{sec}\theta - y\tan\theta = 1$

$\frac{a}{a} - \frac{y}{b} \sin\theta = \cos\theta$

$-1 - \frac{y}{b} \sin\theta = \cos\theta$

$y = \frac{-b - b\cos\theta}{\sin\theta} \quad \therefore M_{QR} \times M_{RS} = \frac{b^2(1 - \cos^2\theta)}{-a^2(e^2 - 1)\sin^2\theta}$

$\text{or } R \quad y = -b(1 + \operatorname{sec}\theta) \quad = \frac{b^2}{-b^2}$

iii) $S(ae, 0) \quad = -1$

$M_{SQ} = 0 - \frac{b(1 - \cos\theta)}{\sin\theta} \quad \therefore \angle OSR = 90^\circ$

$ae - a \quad \text{and } \angle OSR + \angle OSR = 180^\circ$

$= -b(1 - \cos\theta) \quad \text{making } OSR \text{ a cyclic quad. as opposite angles are supplementary.}$

$\frac{\sin\theta}{a(e-1)} \quad M_{RS} = \frac{b(1 + \cos\theta)}{\sin\theta}$

$a(e+1) \quad \text{Now } M_{SQ} \times M_{RS} = \frac{-b(1 - \cos\theta)}{\sin\theta} \times \frac{b(1 + \cos\theta)}{\sin\theta}$

$\frac{a(e-1)}{a(e+1)} \quad = \frac{-b^2(1 - \cos^2\theta)}{\sin^2\theta}$

$\frac{a^2(e^2 - 1)}{-b^2} \quad \text{but } a^2(e^2 - 1) = b^2$

$= -1 \quad \therefore \angle OSR = 90^\circ$

$\frac{a^2(e^2 - 1)}{-b^2} \quad \text{but } a^2(e^2 - 1) = b^2$

$= -1 \quad \therefore \angle OSR = 90^\circ$

$\frac{a^2(e^2 - 1)}{-b^2} \quad \text{but } a^2(e^2 - 1) = b^2$

$= -1 \quad \therefore \angle OSR = 90^\circ$

$\frac{a^2(e^2 - 1)}{-b^2} \quad \text{but } a^2(e^2 - 1) = b^2$

$= -1 \quad \therefore \angle OSR = 90^\circ$

$\frac{a^2(e^2 - 1)}{-b^2} \quad \text{but } a^2(e^2 - 1) = b^2$

$= -1 \quad \therefore \angle OSR = 90^\circ$

Question 15.

Join AO, BF BC

as BO = OC radii

$$1. \angle ABD = \angle OCA = 90^\circ$$

$$\angle DCB = \angle CBO \text{ (equal angles)}$$

$$= \alpha \text{ (opposite equal sides)}$$

(radii to tangent at point of contact is 90°)

Now in $\triangle BOC$

$$\angle BOC = 180 - 2\alpha \text{ (angle sum)}$$

\therefore opposite angles in $\therefore \angle BFC = 90 - \alpha$

ABOC are supplementary and

(angle at the circumference is

ABOC is a cyclic quadrilateral.

half the angle at the centre on arc BC)

$$\text{Now, } \angle ABO = 90^\circ$$

(AO is a diameter or line from

$\therefore \angle BFC = \angle FGE$ ($90 - \alpha$)

midpt to centre is perpendicular)

and the alternate angles

$$\angle OCA = \angle OCB$$

(angles at circumference of circle OCA)

are equal

$$= 90^\circ$$

$\therefore BF \parallel AE$

\therefore AOCB is a cyclic quad

as opposite angles are

supplementary.

$$\text{ii. } \angle OGF = \angle OAC$$

exterior angle of a cyclic

quadrilateral equals opposite

interior angle (AOCB).

$$\text{iii. let } \angle OGF = \angle OAC = \alpha$$

$\therefore \angle FGE = 90 - \alpha$ (straight line)

and

$$\angle OBC = \angle OAC \text{ (angles in the same segment)}$$

$$= \alpha \text{ of } \triangle OBC$$

Question 15 con't

$$I_n = \int_0^1 x^n \sqrt{1-x^3} dx \quad n \geq 2$$

$$= \int_0^1 x^{n-2} \cdot x^2 \sqrt{1-x^3} dx$$

$$= \left[x^{n-2} (1-x^3)^{3/2} \cdot \frac{-2}{9} - \int_0^1 (n-2)x^{n-3} \cdot \frac{-2}{9} (1-x^3) \sqrt{1-x^3} dx \right]_0^1$$

$$I_n = 0 + \frac{2(n-2)}{9} \int_0^1 x^{n-3} \sqrt{1-x^3} dx - \frac{2(n-2)}{9} \int_0^1 x^n \sqrt{1-x^3} dx$$

$$I_n \left[1 + \frac{2(n-2)}{9} \right] = \frac{2(n-2)}{9} \int_0^1 x^{n-3} \sqrt{1-x^3} dx$$

$$I_n \left[\frac{9+2n-4}{9} \right] = \frac{2n-4}{9} I_{n-3}$$

$$I_n = \frac{2n-4}{2n+5} I_{n-3}$$

ii)

$$I_8 = \left(\frac{16-4}{16+5} \right) I_5$$

$$= \frac{12}{21} \left[\frac{(10-4)}{10+5} \right] I_2 \quad \text{Now } I_2 = \int_0^1 x^2 \sqrt{1-x^3} dx$$

$$= \frac{12}{21} \cdot \frac{6}{15} \cdot \frac{2}{9} = \frac{-2}{9} \left[(1-x^3)^{3/2} \right]_0^1$$

$$= \frac{16}{315} = \frac{-2}{9} \left[0 - 1^{3/2} \right]$$

$$= \frac{2}{9}$$

7

8

Question 15 con 1

c)

$$T_1 = 6 \quad T_2 = 27$$

$$T_n = 6T_{n-1} - 9T_{n-2} \quad n \geq 3$$

$$T_n = (n+1)3^n \quad \text{for } n \geq 1$$

Test $n=1$

$$T_1 = (1+1) \times 3^1$$

$$= 2 \times 3$$

= 6 which is given

\therefore True for $n=1$

Assume true for $n=k$

$$\text{ie } T_k = (k+1)3^k$$

where

$$T_k = 6T_{k-1} - 9T_{k-2}$$

Prove true for $n=k+1$

aim to prove

$$T_{k+1} = 6T_k - 9T_{k-1} = (k+1+1) \cdot 3^{k+1}$$

$$T_{k+1} = 6T_k - 9T_{k-1}$$

$$= 6(k+1) \cdot 3^k - 9 \left[k(3^{k-1}) \right]$$

By assumption

$$= 2(k+1) \times 3 \times 3^k - 3^2 \cdot k \cdot 3^{k-1}$$

$$= 2(k+1) \cdot 3^{k+1} - k \cdot 3^{k+1}$$

$$= (2k+2-k) \cdot 3^{k+1}$$

$$= (k+2) \cdot 3^{k+1}$$

$$= (k+1+1) \cdot 3^{k+1}$$

as required

(statement required).

9.

Question 16

let $P = ae^{mx} + be^{-mx}$

b) $\begin{matrix} \uparrow & \downarrow & \downarrow \\ +ve & g & R \\ & & m=1/g \end{matrix}$

$$\frac{dP}{dx} = mae^{mx} - mbe^{-mx} = 0$$

$$ae^{mx} = be^{-mx} \quad \therefore m^2 x^2 = -mg - g \sqrt{v^2} \quad m=1$$

$$ae^{mx} = b$$

$$\therefore x = -g - \frac{g}{k^2} \sqrt{v^2}$$

$$e^{2mx} = b/a$$

$$2mx = \ln(b/a) \quad v \frac{dv}{dx} = -g \left(\frac{k^2 + v^2}{k^2} \right)$$

$$x = \frac{1}{2m} \ln \left(\frac{b}{a} \right)$$

$$\frac{dv}{dx} = -g \left(\frac{k^2 + v^2}{\sqrt{k^2 + v^2}} \right)$$

$$\text{test} \quad \frac{dx}{dv} = -1 \frac{vk^2}{k^2 + v^2}$$

$$\frac{d^2P}{dx^2} = m^2 ae^{mx} + m^2 be^{-mx} \quad \therefore \frac{dv}{dv} = g \frac{k^2 + v^2}{k^2 + v^2}$$

$$\text{at } x = \frac{1}{2m} \ln(b/a) \quad x = -k^2 \int \frac{v}{k^2 + v^2} dv$$

$$\frac{d^2P}{dx^2} > 0 \text{ as } e^{-m} x > 0 \quad x = -k^2 \cdot \frac{1}{2} \ln(k^2 + v^2) + C_1$$

$$\frac{dx^2}{dx^2} > 0 \quad \text{and } a, b, m > 0 \quad x = 0 \quad v = 2k$$

$$\therefore \text{min value is when } v = 2k$$

\therefore min value is when

$$x = \frac{1}{2m} \ln(b/a) \quad \therefore C_1 = k^2/2g \ln(5k^2)$$

$$P = m \left(\frac{1}{2m} \ln(b/a) \right) - m \left(\frac{1}{2m} \ln(b/a) \right)$$

$$ae^{mx} + be^{-mx} \quad x = -k^2 \ln(k^2 + v^2) + k^2 \ln(5k^2)$$

$$= ae^{1/2 \ln(b/a)} + b e^{-1/2 \ln(b/a)} \quad \text{max height } v=0$$

$$= a e^{\ln \sqrt{b/a}} + b e^{\ln \sqrt{a/b}}$$

$$= a \sqrt{\frac{b}{a}} + b \sqrt{\frac{a}{b}} \quad x = -k^2/2g \ln k^2 + k^2/2g \ln(5k^2)$$

$$= \sqrt{\frac{a^2 b}{a}} + \sqrt{\frac{b^2 a}{b}} \quad = k^2/2g \ln \left[\frac{5k^2}{k^2} \right]$$

$$= \sqrt{ab} + \sqrt{ba} = 2\sqrt{ab}$$

$$= \sqrt{ab} + \sqrt{ba} = 2\sqrt{ab}$$

10.

Question 16 con't

b) $x=0$ $t=0$ $v=0$ $\frac{K^2 \ln 5 = -K^2 \ln(K^2 - v^2) + K^2 \ln(K^2)}{2g}$



(m=1)

$\ln 5 = -\ln(K^2 - v^2) + \ln K^2$

$\ln 5 = \ln \left(\frac{K^2}{K^2 - v^2} \right)$

$5(K^2 - v^2) = K^2$

$\ddot{x} = g - R$

$\ddot{x} = g - \frac{g v^2}{K^2}$

$v \frac{dv}{dx} = g - \frac{g v^2}{K^2}$

$\frac{dv}{dx} = \frac{g}{v} - \frac{g v}{K^2}$

$= \frac{g K^2 - g v^2}{v K^2}$

$\frac{dx}{dv} = \frac{v K^2}{g K^2 - g v^2}$

$x = \int \frac{v K^2}{g K^2 - g v^2} dv$

$x = K^2 \int \frac{v}{K^2 - v^2} + C_2$

$\frac{x}{g} = \frac{K^2}{2g} \ln K^2$

$x = \frac{-K^2 \ln(K^2 - v^2) + K^2 \ln K^2}{2g}$

Now $x = K^2 \ln 5$

$\frac{K^2 \ln 5 = -K^2 \ln(K^2 - v^2) + K^2 \ln K^2}{2g}$

$\ln 5 = -\ln(K^2 - v^2) + \ln K^2$

$5(K^2 - v^2) = K^2$

$-5v^2 = -4K^2$

$v^2 = \frac{4K^2}{5}$

$\therefore v = \sqrt{\frac{4K^2}{5}}$

$v > 0$

$V = \frac{2K}{\sqrt{5}}$

$\frac{dx}{dv} = \frac{v K^2}{g K^2 - g v^2}$


$x = \int \frac{v K^2}{g K^2 - g v^2} dv$

$x = K^2 \int \frac{v}{K^2 - v^2} + C_2$

(11)

Question 16 con't

$V = \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2(0) + 0 - \left(\frac{\pi \cdot 1}{2\sqrt{2}} - 2 \frac{\pi}{\sqrt{4}} + \frac{2}{\sqrt{2}} \right) \right]$



x lies of

$y = \cos x$

$\therefore x = \cos^{-1} y$

(I used symmetry can do $\sin^{-1} y$ & $\cos^{-1} y$)

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - \sqrt{2} \right] u^3$

$R = \frac{\pi}{2} - x = \frac{\pi}{2} - \cos^{-1} y$

$\Delta V = \pi (R^2 - r^2) \Delta y$

$= \pi \left[\frac{\pi}{2} - \cos^{-1} y - \cos^{-1} y \right] \left[\frac{\pi}{2} - \cos^{-1} y + \cos^{-1} y \right] \Delta y$

$= \pi \left[\frac{\pi}{2} - 2 \cos^{-1} y \right] \left[\frac{\pi}{2} \right] \Delta y$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2 \cos^{-1} y \right] \Delta y$

other solutions such as

$\int \frac{\pi^2}{2} \int_{\frac{\pi}{2}}^1 2 \sin^{-1} y - \frac{\pi}{2} dy$

can be used.

∴

Total volume

$= \lim_{\Delta y \rightarrow 0} \sum_{y=\frac{1}{\sqrt{2}}}^1 \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2 \cos^{-1} y \right] \Delta y$

$= \frac{\pi^2}{2} \int_{\frac{1}{\sqrt{2}}}^1 \left[\frac{\pi}{2} - 2 \cos^{-1} y \right] dy$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} y - \int 2y \cos^{-1} y - \int \frac{1}{\sqrt{1-y^2}} dy \right]$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} y - 2y \cos^{-1} y + 2 \sqrt{1-y^2} \right]_{\frac{1}{\sqrt{2}}}^1$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2 \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) + 2 \sqrt{1 - \left(\frac{1}{\sqrt{2}} \right)^2} \right]$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - 2 \left(\frac{\pi}{4} \right) + 2 \left(\frac{\sqrt{2}}{2} \right) \right]$

$= \frac{\pi^2}{2} \left[\frac{\pi}{2} - \frac{\pi}{2} + \sqrt{2} \right]$

(12)