



# *Acceleration with Uniform Circular Motion*

Uniform circular motion is when a particle moves with constant angular velocity. ( $\therefore$  the magnitude of the linear velocity will also be constant)




A particle moves from  $A$  to  $P$  with constant angular velocity.

The acceleration of the particle is the change in velocity with respect to time.



This triangle of vectors is similar to  $\Delta OAP$



$$\frac{\Delta v}{AP} = \frac{v}{r}$$


$$\Delta v = \frac{v \cdot AP}{r}$$

$$a = \frac{\Delta v}{\Delta t}$$

$$a = \frac{v \cdot AP}{r \cdot \Delta t}$$

But, as  $\Delta t \rightarrow 0$ ,  $AP = \text{arc}AP$

$$\therefore a = \lim_{\Delta t \rightarrow 0} \frac{v \cdot \text{arc}AP}{r \cdot \Delta t}$$



distance = speed  $\times$  time

$$\text{arc}AP = v \cdot \Delta t$$

$$\therefore a = \lim_{\Delta t \rightarrow 0} \frac{v \cdot v \cdot \Delta t}{r \cdot \Delta t}$$

$$= \lim_{\Delta t \rightarrow 0} \frac{v^2}{r}$$

$$= \frac{v^2}{r}$$

$\therefore$  the acceleration in uniform circular

motion has magnitude  $\frac{v^2}{r}$  and is

directed towards the centre

Acceleration Involved in  
Uniform Circular Motion

$$a = \frac{v^2}{r}$$

OR

$$a = r\omega^2$$

Forces Involved in  
Uniform Circular Motion

$$F = \frac{mv^2}{r}$$

OR

$$F = mr\omega^2$$

e.g. (i) (2003)

A particle  $P$  of mass  $m$  moves with constant angular velocity  $\omega$  on a circle of radius  $r$ . Its position at time  $t$  is given by;

$$x = r \cos \theta$$

$$y = r \sin \theta, \quad \text{where } \theta = \omega t$$

a) Show that there is an inward radial force of magnitude  $mr\omega^2$  acting on  $P$ .

$$x = r \cos \theta$$

$$\dot{x} = -r \sin \theta \cdot \frac{d\theta}{dt}$$

$$= -r\omega \sin \theta$$

$$\ddot{x} = -r\omega \cos \theta \cdot \frac{d\theta}{dt}$$

$$= -r\omega^2 \cos \theta$$

$$= -\omega^2 x$$

$$a^2 = (\ddot{x})^2 + (\ddot{y})^2$$

$$= \omega^4 x^2 + \omega^4 y^2$$


$$= \omega^4 (x^2 + y^2)$$

$$= \omega^4 r^2$$

$$a = r\omega^2$$

$$F = ma$$

$$\therefore F = mr\omega^2$$



$$y = r \sin \theta$$

$$\dot{y} = r \cos \theta \cdot \frac{d\theta}{dt}$$

$$= r\omega \cos \theta$$

$$\ddot{y} = -r\omega \sin \theta \cdot \frac{d\theta}{dt}$$

$$= -r\omega^2 \sin \theta$$

$$= -\omega^2 y$$


$$\alpha = \tan^{-1} \frac{\ddot{y}}{\ddot{x}}$$

$$= \tan^{-1} \left( \frac{-\omega^2 y}{-\omega^2 x} \right)$$

$$= \tan^{-1} \frac{y}{x}$$

$$= \theta$$

$\therefore$  There is a force,  $F = mr\omega^2$ , acting towards the centre



b) A telecommunications satellite, of mass  $m$ , orbits Earth with constant angular velocity  $\omega$  at a distance  $r$  from the centre of the Earth. The gravitational force exerted by Earth on the satellite is  $\frac{Am}{r^2}$  where

$A$  is a constant. By considering all other forces on the satellite to be negligible, show that;

$$r = \sqrt[3]{\frac{A}{\omega^2}}$$

$$m\ddot{x} = mr\omega^2 \quad \begin{array}{c} \bullet \\ \downarrow \text{blue} \\ \downarrow \text{red} \end{array} \quad \frac{Am}{r^2}$$

$$mr\omega^2 = \frac{Am}{r^2}$$

$$r^3 = \frac{Am}{m\omega^2}$$

$$= \frac{A}{\omega^2}$$


$$r = \sqrt[3]{\frac{A}{\omega^2}}$$


---

(ii) A string is 50cm long and it will break if a mass exceeding 40kg is hung from it.

A mass of 2kg is attached to one end of the string and it is revolved in a circle.

Find the greatest angular velocity which may be imparted without breaking the string.




$$m\ddot{x} = mg - T$$

$$0 = mg - T$$

$$T = (40)(9.8)$$

$$= 392N$$



$$T = mr\omega^2$$

$$392 = (2)(0.5)\omega^2$$

$$\omega^2 = 392$$

$$\omega = \sqrt{392}$$

$$\omega = 2\sqrt{98}\text{rad/s}$$

**Patel Exercise 9B; all**

**“Cambridge” Exercise 9A; 1 to 4, 6, 9, 12, 14**