## Dilations

**horizontal dilation (stretch)** replace x with  $\frac{x}{a}$  $y = f\left(\frac{x}{a}\right)$ 

#### curve is stretched horizontally by a factor of *a* (*if a* > 1, *curve is shallower*)


domain may be altered, range will be unchanged

## vertical dilation (stretch) replace y with $\frac{y}{a}$


$$\frac{y}{a} = f(x) \ OR \ y = af(x)$$

curve is stretched vertically by a factor of *a* (*if a* > 1, *curve is steeper*)

domain unchanged, range may be altered



stretch vertically by a factor of 2



# Enlargements

An enlargement is when the **same** dilation factor is applied both horizontally and vertically.

$$f(x,y) \Rightarrow f\left(\frac{x}{a}, \frac{y}{a}\right)$$

e.g. The circle  $(x - 1)^2 + (y + 2)^2 = 1$  is enlarged by a factor of 2. Using the origin as the centre of enlargement, find the circle's new equation


$$(x-1)^{2} + (y+2)^{2} = 1 \implies \left(\frac{x}{2} - 1\right)^{2} + \left(\frac{y}{2} + 2\right)^{2} = 1$$
$$(x-2)^{2} + (y+4)^{2} = 4$$

### Not all transformations commute A mathematical operation commutes if the order of the objects being

operated on does not matter

addition and multiplication commute 3 + 4 = 4 + 3 $3 \times 4 = 4 \times 3$  subtraction and division do not commute  $3-4 \neq 4-3$  $3 \div 4 \neq 4 \div 3$ 

Transformations can be done in any order, with the exception of a dilation and a translation in the **same** direction



e.g. Determine the equation after  $y = x^2$  has been;

(i) shifted up 1 unit then reflected vertically

$$x^{2} \rightarrow x^{2} + 1 \rightarrow -(x^{2} + 1) \implies y = -x^{2} - 1$$

(ii) reflected vertically the shifted up 1 unit

$$x^2 \rightarrow -x^2 \rightarrow -x^2 + 1 \Rightarrow y = 1 - x^2$$

Exercise 3H; 1aceg, 2bd, 3, 4, 5a, 6a, 8, 9ad, 10, 13, 14, 15a, 17

**Exercise 3I; 1 to 5, 6bdfh, 8ab, 12**