

Dot (Scalar) Product

$$\text{Let } \underset{\sim}{u} = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} \text{ and } \underset{\sim}{v} = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}$$

then;

$$\underset{\sim}{u} \cdot \underset{\sim}{v} = x_1 x_2 + y_1 y_2$$

NOTE: the result is a scalar

$$\text{e.g. } \begin{pmatrix} i \\ -7j \end{pmatrix} \cdot \begin{pmatrix} 6i \\ 4j \end{pmatrix} = 1 \times 6 + (-7) \times 4 \\ = \underline{\underline{-22}}$$


$$\lambda \underset{\sim}{u} \cdot \underset{\sim}{v} = \lambda(\underset{\sim}{u} \cdot \underset{\sim}{v})$$

$$\underset{\sim}{u} \cdot \underset{\sim}{u} = x_1^2 + y_1^2 \\ = |\underset{\sim}{u}|^2$$

$$\underset{\sim}{a} \cdot (\underset{\sim}{u} + \underset{\sim}{v}) = \underset{\sim}{a} \cdot \underset{\sim}{u} + \underset{\sim}{a} \cdot \underset{\sim}{v}$$

$$(\underset{\sim}{u} + \underset{\sim}{v}) \cdot (\underset{\sim}{u} - \underset{\sim}{v}) = \underset{\sim}{u} \cdot \underset{\sim}{u} - \underset{\sim}{v} \cdot \underset{\sim}{v} \\ = |\underset{\sim}{u}|^2 - |\underset{\sim}{v}|^2$$

$$\underset{\sim}{u} \cdot \underset{\sim}{v} = \underset{\sim}{v} \cdot \underset{\sim}{u}$$



By the cosine rule;

$$|\underline{u} - \underline{v}|^2 = |\underline{u}|^2 + |\underline{v}|^2 - 2|\underline{u}||\underline{v}| \cos \theta$$

$$(x_1 - x_2)^2 + (y_1 - y_2)^2 = x_1^2 + x_2^2 + y_1^2 + y_2^2 - 2|\underline{u}||\underline{v}| \cos \theta$$

$$-2x_1 x_2 - 2y_1 y_2 = -2|\underline{u}||\underline{v}| \cos \theta$$

$$2(\underline{u} \cdot \underline{v}) = 2|\underline{u}||\underline{v}| \cos \theta$$

$$\underline{u} \cdot \underline{v} = |\underline{u}||\underline{v}| \cos \theta$$

Let $\underline{u} = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$ and $\underline{v} = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}$

then;

$$\underline{u} \cdot \underline{v} = x_1 x_2 + y_1 y_2$$

$$= |\underline{u}||\underline{v}| \cos \theta$$

NOTE: θ is acute or obtuse

e.g. Find, to the nearest degree, the angle between the two vectors

$$\underline{\underline{a}} = 3\underline{\underline{i}} - 2\underline{\underline{j}} \text{ and } \underline{\underline{b}} = 4\underline{\underline{i}} + \underline{\underline{j}}$$

$$\underline{\underline{a}} \cdot \underline{\underline{b}} = |\underline{\underline{a}}| |\underline{\underline{b}}| \cos \theta$$

$$\cos \theta = \frac{\underline{\underline{a}} \cdot \underline{\underline{b}}}{|\underline{\underline{a}}| |\underline{\underline{b}}|}$$

$$\cos \theta = \frac{3 \times 4 + (-2) \times 1}{\sqrt{13} \times \sqrt{17}}$$

$$= \frac{10}{\sqrt{221}}$$

$$\underline{\underline{\theta = 48^\circ}} \text{ (to nearest degree)}$$

Consequences of the Dot Product

$$\underline{u} \cdot \underline{v} = 0 \iff \underline{u} \perp \underline{v}$$

$$\therefore \underline{i} \cdot \underline{j} = \underline{j} \cdot \underline{i} = 0$$

$$\underline{u} \cdot \underline{v} = \pm |\underline{u}| |\underline{v}| \iff \underline{u} \parallel \underline{v}$$

$|\underline{u}| |\underline{v}| > 0 \Rightarrow \underline{u}$ and \underline{v} have the same direction

$|\underline{u}| |\underline{v}| < 0 \Rightarrow \underline{u}$ and \underline{v} have opposite directions

$$\therefore \underline{i} \cdot \underline{i} = \underline{j} \cdot \underline{j} = 1$$

**Exercise 8C; 1ac, 2a, 3a, 4b, 5ac, 6, 8, 9b, 10,
11 abc (i, iv, vi), 12, 13, 15, 17, 20, 21**