

# *Geometric Proofs*

## **Things to keep in mind when using vectors in a geometric proof**

- \* choose a vertex or other key point to represent the origin
- \* sum (or difference) of vectors creates a triangle
- \* parallel vectors are multiples of each other
- \* the angle between vectors can be found using the dot product
- \* perpendicular vectors have a dot product equal to zero
- \* vectors can be written as their position vector using “*head minus tail*”

e.g. Prove that the diagonals of a rhombus are perpendicular

$OABC$  is a rhombus

Let  $\vec{OA} = \vec{a}$  and  $\vec{OC} = \vec{c}$

Diagonals are  $OB$  and  $AC$

$$\vec{OB} = \vec{a} + \vec{c}$$


$$\vec{AC} = \vec{c} - \vec{a}$$

$$\begin{aligned}\vec{OB} \cdot \vec{AC} &= (\vec{a} + \vec{c}) \cdot (\vec{c} - \vec{a}) \\ &= \vec{c} \cdot \vec{c} - \vec{a} \cdot \vec{a} \\ &= |\vec{c}|^2 - |\vec{a}|^2\end{aligned}$$

however  $|\vec{a}| = |\vec{c}|$

so  $\vec{OB} \cdot \vec{AC} = 0$

$\therefore OB \perp AC$



(sides in a rhombus are =)


(ii) Prove Pythagoras' Theorem

$OAB$  is a right angled triangle

Let  $\vec{OA} = \underline{a}$  and  $\vec{OB} = \underline{b}$

$$\vec{AB} = \underline{b} - \underline{a}$$

$$\begin{aligned} OA^2 + OB^2 \\ = |\underline{a}|^2 + |\underline{b}|^2 \end{aligned}$$



$$\begin{aligned} AB^2 &= |\underline{b} - \underline{a}|^2 \\ &= (\underline{b} - \underline{a}) \cdot (\underline{b} - \underline{a}) \\ &= \underline{b} \cdot \underline{b} - 2\underline{a} \cdot \underline{b} + \underline{a} \cdot \underline{a} \end{aligned}$$

however  $OA \perp OB$

$$\text{so } \underline{a} \cdot \underline{b} = 0$$

$$\begin{aligned} AB^2 &= \underline{b} \cdot \underline{b} + \underline{a} \cdot \underline{a} \\ &= |\underline{b}|^2 + |\underline{a}|^2 \end{aligned}$$

$$\underline{\therefore AB^2 = OA^2 + OB^2}$$

(iii)  $ABCD$  is a quadrilateral,  $P$ ,  $Q$ ,  $R$  and  $S$  are midpoints of the lines  $AC$ ,  $BD$ ,  $AD$  and  $BC$  respectively.

What type of quadrilateral is  $PRQS$ ?

treat  $A$  as the origin


$$\text{Let } \vec{AB} = \underline{u}, \vec{AC} = \underline{v}, \vec{AD} = \underline{w}$$

$$P = \frac{1}{2} \vec{AC} = \frac{1}{2} \underline{v}$$

$$Q = \frac{1}{2} \vec{BD} = \frac{1}{2} (\underline{w} - \underline{u})$$

$$R = \frac{1}{2} \vec{AD} = \frac{1}{2} \underline{w}$$

$$S = \frac{1}{2} \vec{BC} = \frac{1}{2} (\underline{v} - \underline{u})$$



$$\vec{PR} = \frac{1}{2} \underline{v} - \frac{1}{2} \underline{w} = \frac{1}{2} (\underline{v} - \underline{w})$$

$$\vec{SQ} = \frac{1}{2} (\underline{v} - \underline{u}) - \frac{1}{2} (\underline{w} - \underline{u}) = \frac{1}{2} (\underline{v} - \underline{w})$$

$$\therefore \vec{PR} = \vec{SQ}$$

**Exercise 8D; 1 to 4, 6 to 8, 10, 11, 12**

Thus  $PRQS$  is a parallelogram as a pair of opposite sides are both equal and parallel