

Integration By Partial Fractions

To find; $\int \frac{A(x)}{P(x)} dx$

(1) If $\deg A(x) \geq \deg P(x)$, perform a division

(2) If $\deg A(x) < \deg P(x)$, factorise $P(x)$

a) for linear factor $(x - a)$, write $\frac{A}{x - a}$

b) for multiple linear factors $(x - a)^n$, write

$$\frac{A}{(x - a)} + \frac{B}{(x - a)^2} + \dots + \frac{C}{(x - a)^n}$$

c) for polynomial factors e.g. $ax^2 + bx + c$, write $\frac{Ax + B}{ax^2 + bx + c}$

$$\begin{aligned} \text{e.g. } (i) & \int \frac{x^2}{x+1} dx \\ &= \int \left[x - 1 + \frac{1}{x+1} \right] dx \\ &= \frac{x^2}{2} - x + \log|x+1| + c \end{aligned}$$

$$\begin{array}{r} x - 1 \\ x + 1 \end{array} \overline{) x^2 + 0x + 0} \\ \underline{x^2 + x} \\ -x + 0 \\ \underline{-x - 1} \\ 1$$

$$\begin{aligned} (ii) & \int \frac{3dx}{x^2 - x} \\ &= \int \frac{3dx}{x(x-1)} \\ &= \int \left[\frac{-3}{x} + \frac{3}{(x-1)} \right] dx \\ &= -3\log|x| + 3\log|x-1| + c \\ &= 3\log \left| \frac{x-1}{x} \right| + c \end{aligned}$$

$$\begin{aligned} \frac{A}{x} + \frac{B}{x-1} &= \frac{3}{x(x-1)} \\ A(x-1) + Bx &= 3 \end{aligned}$$

$$\begin{array}{ll} \xrightarrow{x=0} & \xrightarrow{x=1} \\ -A = 3 & B = 3 \\ A = -3 & \end{array}$$

$$\begin{aligned}
 (iii) & \int \frac{x+5}{x^2 - 3x - 10} dx \\
 &= \int \frac{x+5}{(x-5)(x+2)} dx \\
 &= \int \left[\frac{10}{7(x-5)} - \frac{3}{7(x+2)} \right] dx \\
 &= \frac{10}{7} \log|x-5| - \frac{3}{7} \log|x+2| + c
 \end{aligned}$$

$$\begin{aligned}
 \frac{A}{(x-5)} + \frac{B}{(x+2)} &= \frac{x+5}{(x-5)(x+2)} \\
 A(x+2) + B(x-5) &= x+5 \\
 \underline{x = -2} & \qquad \qquad \underline{x = 5} \\
 -7B &= 3 \qquad \qquad 7A = 10 \\
 B &= \frac{-3}{7} \qquad \qquad A = \frac{10}{7}
 \end{aligned}$$

$$\begin{aligned}
 (iv) & \int \frac{dx}{x^3 + x} \\
 &= \int \frac{dx}{x(x^2 + 1)} \\
 &= \int \left[\frac{1}{x} - \frac{x}{x^2 + 1} \right] dx \\
 &= \log|x| - \frac{1}{2} \log(x^2 + 1) + c
 \end{aligned}$$

$$\begin{aligned}
 \frac{A}{x} + \frac{Bx+C}{x^2+1} &= \frac{1}{x(x^2+1)} \\
 A(x^2+1) + (Bx+C)x &= 1 \\
 \underline{x = 0} & \qquad \qquad \underline{x = i} \\
 A &= 1 \qquad \qquad -B + Ci = 1 \\
 & \qquad \qquad B = -1 \qquad C = 0
 \end{aligned}$$

$$\begin{aligned}
 & (v) \int \frac{x dx}{(x+1)^2(x^2+1)} \quad \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{Cx+D}{(x^2+1)} = \frac{x}{(x+1)^2(x^2+1)} \\
 & A(x+1)(x^2+1) + B(x^2+1) + (Cx+D)(x+1)^2 = x \\
 & = \int \left[\frac{-1}{2(x+1)^2} + \frac{1}{2(x^2+1)} \right] dx \quad \begin{array}{l} \underline{x = -1} \\ 2B = -1 \end{array} \quad \begin{array}{l} \underline{x = i} \\ -2C + 2Di = i \end{array} \\
 & = \frac{1}{2} \left(\frac{1}{x+1} + \tan^{-1} x \right) + c \quad \begin{array}{l} B = \frac{-1}{2} \\ C = 0 \quad D = \frac{1}{2} \end{array} \\
 & \quad \underline{x = 0} \quad 2A + B + D = 0 \\
 & \quad 2A - \frac{1}{2} + \frac{1}{2} = 0 \\
 & \quad A = 0
 \end{aligned}$$

*Alternative method for finding constants
when denominator is a product of distinct linear factors*

$$\frac{A}{(x-a)} + \frac{B}{(x-b)} + \frac{C}{(x-c)} + \dots = \frac{f(x)}{(x-a)(x-b)(x-c)\dots}$$

- 1) To find A substitute a into $\frac{f(x)}{(x-b)(x-c)\dots}$ i.e. $A = \frac{f(a)}{(a-b)(a-c)\dots}$
- 2) To find B substitute b into $\frac{f(x)}{(x-a)(x-c)\dots}$ i.e. $B = \frac{f(b)}{(b-a)(b-c)\dots}$
- 3) To find C substitute c into $\frac{f(x)}{(x-a)(x-b)\dots}$ i.e. $C = \frac{f(c)}{(c-a)(c-b)\dots}$
and so on for all of the constants

e.g. (i) $\frac{A}{x} + \frac{B}{x-1} = \frac{3}{x(x-1)}$

$$\begin{aligned} A &= \frac{3}{(0-1)} \\ &= -3 \end{aligned} \qquad \begin{aligned} B &= \frac{3}{1} \\ &= 3 \end{aligned}$$

(ii) $\frac{A}{(x-5)} + \frac{B}{(x+2)} = \frac{x+5}{(x-5)(x+2)}$

$$\begin{aligned} A &= \frac{5+5}{(5+2)} \\ &= \frac{10}{7} \end{aligned} \qquad \begin{aligned} B &= \frac{-2+5}{(-2-5)} \\ &= -\frac{3}{7} \end{aligned}$$

using complex numbers, the idea can be applied to quadratic factors

$$(iii) \frac{A}{x} + \frac{Bx+C}{x^2+1} = \frac{1}{x(x^2+1)}$$

$$\begin{aligned} A &= \frac{1}{(0^2+1)} & Bi+C &= \frac{1}{i} \\ &= 1 & &= -i \\ B &= -1 & C &= 0 \end{aligned}$$

$$(iv) \frac{A}{(x-1)} + \frac{Bx+C}{4x^2+1} = \frac{x^2+4x}{(x-1)(4x^2+1)}$$

$$\begin{aligned} A &= \frac{1^2+4(1)}{(4(1)^2+1)} & \frac{Bi}{2} + C &= \frac{\left(\frac{i}{2}\right)^2+4\left(\frac{i}{2}\right)}{\left(\frac{i}{2}-1\right)} \\ &= 1 \end{aligned}$$

$$\begin{aligned} &= \frac{-\frac{1}{4}+2i}{\frac{i}{2}-1} \\ &= \frac{-1+8i}{2i-4} \times \frac{2i+4}{2i+4} \\ &= \frac{-2i-4-16+32i}{-4-16} \end{aligned}$$

$$B = -3 \quad C = 1 \quad = 1 - \frac{3}{2}i$$

multiple factors require just a little bit more work

$$(v) \frac{A}{x+2} + \frac{B}{x-1} + \frac{C}{(x-1)^2} = \frac{9x}{(x+2)(x-1)^2}$$

the constant of the higher power is found the same way

$$A = \frac{9(-2)}{(-2-1)^2} \\ = -2$$

$$C = \frac{9(1)}{(1+2)} \\ = 3$$

the third constant is found via a simple substitution

let $x = 0$

$$\frac{A}{0+2} + \frac{B}{0-1} + \frac{C}{(0-1)^2} = \frac{9(0)}{(0+2)(0-1)^2}$$
$$-1 - B + 3 = 0$$
$$B = 2$$

$$(vi) \quad \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{Cx+D}{(x^2+1)} = \frac{x}{(x+1)^2(x^2+1)}$$

$$B = \frac{-1}{((-1)^2 + 1)} \quad Ci + D = \frac{i}{(i+1)^2} \quad \text{let } x = 0 \quad A + B + D = 0$$

$$= -\frac{1}{2} \quad \quad \quad = \frac{i}{2i} \quad \quad \quad A - \frac{1}{2} + \frac{1}{2} = 0$$

$$= \frac{1}{2} \quad \quad \quad \quad \quad \quad \quad \quad \quad A = 0$$

$$C = 0 \quad D = \frac{1}{2}$$

quadratic denominator that can't be factorised

$$(vii) \frac{A}{x+1} + \frac{Bx+C}{x^2+4x+5} = \frac{2}{(x+1)(x^2+4x+5)}$$

Complete the square
to find
an appropriate
substitution

$$A = \frac{2}{((-1)^2 + 4(-1) + 5)}$$
$$= 1$$

$$x^2 + 4x + 5 = (x+2)^2 + 1$$

$$\text{let } x = -2 + i \quad B(-2+i) + C = \frac{2}{(-2+i+1)}$$

$$\begin{aligned}-2B + C + Bi &= \frac{2}{-1+i} \times \frac{-1-i}{-1-i} \\&= \frac{-2-2i}{2} \\&= -1-i\end{aligned}$$

$$B = -1$$

$$-2B + C = -1$$

$$2 + C = -1$$

$$C = -3$$

**Exercise 4D; 1, 2, 3, 4bc, 5ac,
6b, 7, 8ab (i), 9, 10, 11b,
12, 13bcef , 14**