

Definite Integral

Fundamental Theorem of Calculus

$$F'(x) = \frac{d}{dx} \left[\int_a^x f(t) dt \right] = f(x)$$

Properties of the Definite Integral

$$(1) \int_a^b x^n dx = \left[\frac{x^{n+1}}{n+1} \right]_a^b$$

$$(2) \int_a^b kf(x)dx = k \int_a^b f(x)dx \quad (\text{can only factorise constants})$$

$$(3) \int_a^b \{f(x) \pm g(x)\} dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

$$(4) \int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

$$(5) \int_a^b f(x) dx > 0 \quad , \text{if } f(x) > 0 \text{ for } a < x < b$$
$$< 0 \quad , \text{if } f(x) < 0 \text{ for } a < x < b$$

$$(6) \int_a^b f(x)dx < \int_a^b g(x)dx \quad , \text{if } f(x) < g(x) \text{ for } a < x < b$$

$$(7) \int_a^b f(x)dx = -\int_b^a f(x)dx$$

$$(8) \int_{-a}^a f(x) = 0, \quad \text{if } f(x) \text{ is odd}$$

$$(9) \int_{-a}^a f(x) = 2 \int_0^a f(x), \quad \text{if } f(x) \text{ is even}$$

NOTE :

odd \times *odd* = *even*

odd \times *even* = *odd*

even \times *even* = *even*

$$\begin{aligned} \text{e.g. (i)} \quad \int_1^2 6x^2 dx &= 6 \left[\frac{1}{3} x^3 \right]_1^2 \\ &= 2(2^3 - 1^3) \\ &= \underline{14} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \int_0^5 \sqrt[3]{x} dx &= \int_0^5 x^{\frac{1}{3}} dx \\ &= \frac{3}{4} \left[x^{\frac{4}{3}} \right]_0^5 \\ &= \frac{3}{4} \left[x^{\sqrt[3]{x}} \right]_0^5 \\ &= \frac{3}{4} (5^{\sqrt[3]{5}} - 0) \\ &= \underline{\frac{15^{\sqrt[3]{5}}}{4}} \end{aligned}$$


$$(iii) \int_{-2}^2 \sin^5 x dx = 0 \quad (\text{odd function})^5 = \text{odd function}$$

$$\begin{aligned}(iv) \int_{-1}^1 (x^3 + 2x^2 + x + 1) dx &= 2 \int_0^1 (2x^2 + 1) dx \\ &= 2 \left[\frac{2}{3} x^3 + x \right]_0^1 \\ &= 2 \left\{ \frac{2}{3} (1)^3 + 1 \right\} - 0 \\ &= \frac{10}{3}\end{aligned}$$

Sketching the Signed Area Function (primitive function)

$$A(x) = \int_a^x f(t) dt$$

$$= F(x) - F(a)$$



$$A(a) = 0 \quad \text{NOTE: } a \text{ is a stationary point as } A'(x) = 0$$

$$A(0) = \int_a^0 f(t) dt = - \int_0^a f(t) dt = -A$$

$$A(b) = -A$$


$$\text{NOTE: } b \text{ is a possible inflection point as } A''(x) = f'(x) = 0$$

$$A(c) = -2A$$

$$\text{NOTE: } c \text{ is a stationary point as } A'(x) = 0$$

$$A(d) = -A$$

$$\text{NOTE: in the interval } [c, \infty) A(x) \text{ is increasing as } A'(x) > 0$$



$$\text{Area } 0 \rightarrow a = \text{Area } a \rightarrow b = \text{Area } b \rightarrow c = \text{Area } c \rightarrow d = A$$

Exercise 5C; 5, 6a, 7, 10, 11, 12a, 16, 18

Exercise 5D; 3, 4b, 7bc, 9