


# Graphs with Asymptotes

**Asymptotes** are a geometrical way of describing the behaviour of a function at its extremities

## Types of Asymptotes



**Vertical asymptotes** occur if  $\lim_{x \rightarrow a} f(x) = \pm\infty$

Functions **do not** touch/cut vertical asymptotes

**Horizontal asymptotes** occur if  $\lim_{x \rightarrow \pm\infty} f(x) = b$

**Oblique asymptotes** occur if  $\lim_{x \rightarrow \pm\infty} f(x) = \pm\infty$


*Notes: oblique asymptotes are straight lines, asymptotes could be any function that approaches infinity  
functions **can** touch/cut horizontal and oblique asymptotes*

# *Rectangular Hyperbolic Function*

$$y = \frac{1}{x}$$

OR

$$xy = 1$$



Any data that demonstrates **inverse variation** will lie on a rectangular hyperbola.

Rectangular hyperbolas have two asymptotes that are perpendicular

All rectangular hyperbolas can be transformed from the basic equation  $y = \frac{1}{x}$  using translations, rotations, reflections or a combination of all three.


## Recognising the hyperbolic function

$$y = \frac{1}{x}$$

- one variable is in the numerator of a fraction, the other is in the denominator of another fraction


# *Exponential Functions*

The orientation of the basic **exponential function** is determined by the base



exponentials with a base  $> 1$  start shallow and increase more rapidly as the dependent variable increases

exponentials with a base  $< 1$  start steep and decrease less rapidly as the dependent variable increases



## Recognising the exponential function

$$y = a^x$$

- one variable is in the power (or **exponent**)
- the base is positive (not equal to 1)

**Exercise 3H; 1, 2, 3,  
4, 10, 11, 13b,  
14a, 15, 18, 19**