


Euler's Formula


Maclaurin Series

A Maclaurin series is a representation of a function as an infinite sum

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$




$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$



$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$y = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!}$$



Euler's Formula

Euler's formula uses imaginary numbers to convert between exponential and trigonometric functions

Let $x = \theta$ in the Maclaurin series expressions for $\cos x$ and $\sin x$

$$\cos \theta = 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \dots \quad \sin \theta = \theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} + \dots$$

Let $x = i\theta$ in the expression for e^x

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \dots \\ &= 1 + i\theta - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} + \dots \\ &= \left(1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots \right) + i \left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \dots \right) \\ &= \cos \theta + i \sin \theta \end{aligned}$$

Euler's Formula

$$e^{i\theta} = \cos\theta + i\sin\theta$$

for real θ

This now gives a third way for expressing complex numbers

$$z = x + iy = r(\cos\theta + i\sin\theta) = re^{i\theta}$$

Cartesian
form

polar
form

exponential
form

e.g. Write $4 - 4i$ in exponential form

$$\begin{aligned} 4 - 4i &= 4\sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right) \\ &= \underline{4\sqrt{2} e^{-\frac{i\pi}{4}}} \end{aligned}$$

Using Euler's Formula for multiplication and division

e.g. if $z = \sqrt{3} + i$ and $w = 1 - \sqrt{3}i$, find;

$$(i) zw = (\sqrt{3} + i)(1 - \sqrt{3}i)$$

$$= 2e^{\frac{i\pi}{6}} \times 2e^{-\frac{i\pi}{3}}$$

$$= 4e^{-\frac{i\pi}{6}}$$

$$= 4 \left(\cos\left(-\frac{\pi}{6}\right) + i \sin\left(-\frac{\pi}{6}\right) \right)$$

$$= \underline{2\sqrt{3} - 2i}$$

$$(ii) \frac{z}{w} = \frac{2e^{\frac{i\pi}{6}}}{2e^{-\frac{i\pi}{3}}}$$

$$= e^{\frac{i\pi}{2}}$$

$$= \underline{i}$$

$$(iii) z^5 = 2^5 e^{\frac{5i\pi}{6}}$$

$$= 32 \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i \right)$$

$$= \underline{-16 - 16\sqrt{3}i}$$

**Exercise 3D; 1ac, 2cd, 3abf,
4ade, 5, 6bd, 7, 8, 11ad,
12ab (i, ii, iv), 14bd**